

Intrinsic risk factors for acute ankle injuries among male soccer players – a prospective cohort study

*A. H. Engebretsen, MD¹ **

G. Myklebust, PT, PhD¹

I. Holme, PhD¹

L. Engebretsen, MD PhD^{1,2}

R. Bahr, MD PhD¹

¹Oslo Sports Trauma Research Center
Norwegian School of Sports Sciences
PO Box 4014 Ullevål Stadion
N-0806 Oslo
Norway

²Department of Orthopaedic Surgery
Ullevål University Hospital and Medical School
University of Oslo
N-0407 Oslo
Norway

Corresponding Author:
***Engebretsen AH, MD**
Oslo Sports Trauma Research Center
Norwegian School of Sports Science
PO Box 4014 Ullevål Stadion
N-0806 Oslo
Norway
Tel: (+47) 23 26 23 57
Fax: 23 26 23 07
E-mail: Anders.Engebretsen@nih.no

Running title: "Risk factors for ankle injuries in soccer"

Key terms: Ankle injuries; football; risk factors; prospective cohort study; previous injuries.

Abstract

This prospective cohort study was conducted to identify risk factors for acute ankle injuries among male soccer players. A total of 508 players representing 31 amateur teams were tested during the 2004 preseason through a questionnaire on previous injury and function score (Foot and Ankle Outcome Score; FAOS), functional tests (balance tests on the floor and a balance mat), and a clinical examination of the ankle. Generalized estimating equations were used in univariate analyses to identify candidate risk factors, and factors with a p-value <0.10 were then examined in a multivariate model. During the season, 56 acute ankle injuries, affecting 46 legs (43 players), were registered. Univariate analyses identified a history of previous acute ankle injuries (OR per previous injury: 1.25, 95% CI 1.09 to 1.43) and the FAOS sub-score "Pain" (OR for a 10-point difference in score: 0.81, 95% CI 0.62 to 1.04) as candidate risk factors. In a multivariate analysis, only the number of previous acute ankle injuries proved to be a significant (adjusted OR per previous injury: 1.23; 95 % CI 1.06 to 1.41, $P=0.005$) predictor of new injuries. Function scores, functional tests and clinical examination could not independently identify players at increased risk in this study.

Introduction

The ankle joint is one of the most common injury locations in sports in general and soccer in particular. The injury incidence ranges from 1.7 to 4.5 injuries per 1000 playing hours, accounting for 11-25% of all acute injuries (Ekstrand and Tropp, 1990, Árnason et al., 1996, Andersen et al., 2004, Hawkins and Fuller, 1999, Junge et al., 2004, Juma, 1998). An ankle sprain may leave an athlete out of play for several weeks, and in many cases full recovery takes much longer. Injuries to the ankle are therefore a concern.

To possibly prevent new injuries, the specific intrinsic and extrinsic risk factors for the injury type in question must be known (Meeuwisse WH, 1994). Regarding intrinsic risk factors, it has been suggested that previous injury, especially when rehabilitation is inadequate, places an athlete at increased risk of suffering an injury to the ankle (Tropp et al., 1985, Árnason et al., 2004, Kofotolis et al., 2007, Ekstrand and Gillquist, 1983). Several other potential risk factors have been tested and suggested as possible predictors of increased risk, however, with limited data on male soccer players. These include slow reaction time (Taimela et al., 1990, Árnason et al., 2004), personality factors (Taimela et al., 1990, Árnason et al., 2004, Lysens et al., 1989, Taerk, 1977, Junge et al., 2000), age (Backous et al., 1988, Lindenfeld et al., 1994, Ostenberg and Roos, 2000), general joint laxity (Baumhauer et al., 1995, Beynnon et al., 2001, Ostenberg and Roos, 2000), ankle joint laxity (Beynnon et al., 2001) and balance tests (Trojian and McKeag, 2006). Regarding body size measures such as height, weight and body mass index (BMI), the literature is also inconclusive (Backous et al., 1988, Tyler et al., 2006, Baumhauer et al., 1995, Beynnon et al., 2001). Some risk factors have been tested further in intervention studies, and balance training (Tropp et al., 1985) and orthoses (Tropp et al., 1985, Surve et al., 1994) have resulted in significantly fewer ankle sprains, indicating that reduced neuromuscular control is an important risk factor for ankle injuries.

To examine the contribution of the various risk factors of injuries and etiology and to explore their interrelationship, it is necessary to include all in a multivariate analysis (Meeuwisse WH, 1994). Even though a large number of risk factor studies have been carried out, only a few of them have included multivariate analyses. We therefore planned the present prospective cohort study on soccer players to screen for several potential risk factors for ankle injuries, some of which have not been studied in depth earlier.

Elite players only constitute a small portion of all soccer players, and advanced resources for screening tests are not available for the majority of players. Therefore, one goal of this study was to investigate if simple screening tests, which are easy to do and not require advanced equipment, can be used to identify individuals at risk. In this way, if the questionnaire and balance tests in this study prove useful, teams and players with no medical staff can test themselves in the pre-season to find out whether they have an increased risk of injuries.

We hypothesized that previous ankle injuries, reduced function scores and abnormalities on a clinical examination or balance tests indicating reduced neuromuscular control, could predict increased risk of new ankle injuries. In addition, we included clinical examination and player information such as age, height, weight, BMI and player position to investigate if there were any correlations between these variables and injury risk.

Hence, the aim of this study was to examine potential intrinsic risk factors for injuries to the ankle in a prospective cohort study among subelite male soccer players.

Methods

Design and participants

This study is based on data from a randomized trial on male amateur soccer players examining the effect of a training program designed to prevent injuries. The design, the

intervention program, and the results of the study have previously been described in detail in a separate paper (Engebretsen et al., 2008). Because no differences were seen in injury rates between the intervention and control groups (Engebretsen et al., 2008), the entire cohort could be used to assess the effect of a number of risk factors assessed at baseline.

A total of 35 teams (n=769 players) from the Norwegian 1st, 2nd or 3rd division of soccer for men, geographically located in the proximity of Oslo, were invited to participate in the study. The 3rd division teams either won their league or finished as first runners up the previous season, resulting in a relatively homogenous group of teams, even if they competed in three different divisions. Three of the teams (n=60 players) declined the invitation to participate, 177 players did not report for testing, three players did not speak Norwegian and could therefore not complete the questionnaire and four players were excluded for other reasons (Figure 1). Hence, 244 of the players invited could not be included. In addition, one team (n=17 players) was later excluded because the physiotherapist did not record injuries, resulting in a final sample of 508 players representing 31 teams from three divisions (1st division, n=7, 122 players; 2nd division, n=16, 260 players; and 3rd division, n=8; 126 players).

Risk factor screening

The teams were tested for potential risk factors for ankle injuries during the 2004 pre-season, January through March, at the Norwegian School of Sport Sciences. Every player capable (not injured at the time) completed single leg balance tests for both legs, both on a balance mat and on the floor, a clinical examination and a questionnaire.

For the balance tests players were asked to stand barefoot on one straight leg, keeping his arms crossed across the chest and his other leg bent 90° at the knee, and only using the ankle

joint to correct his balance. Both balance tests (Figure 2A and 2B) were scored in the same manner (quantitatively and qualitatively), in 5 categories:

- 5 points (maximum score): The player can keep his balance for 60 s with eyes open and for an additional 15 s with eyes closed, always using an ankle strategy only to keep his balance

- 4 points: The player can keep his balance for 60 s with eyes open, using an ankle strategy only for at least 45 s of this period

- 3 points: The player is able to keep his balance for 60 s with eyes open, but needs to use other body parts than the ankle joint (knee, hip, torso, arms) to correct his balance for more than 15 s of this period

- 2 points: The player can balance for 60 s but needs to use the upper body and touch the floor with his other foot at times to correct imbalance

- 1 point: The player can not manage to balance on one leg for more than short periods of time.

The clinical testing of the players was performed by a group of ten sports physical therapists and sports physicians who were blinded for any injury history. Both legs were examined for foot type (normal, pes planus, pes cavus, splayed forefoot), standing rearfoot alignment (normal, valgus), hallux position (normal, valgus), anterior drawer (normal, pathologic) and range of motion for supination, pronation (measured in degrees with the ankle in 10° of plantar flexion) and dorsiflexion.

The players also completed a questionnaire in two parts, where the first part covered general player information (age, height, body mass index, position on the field, number of junior or senior national team matches played, level of play this season, and level of play the previous

season), and history of previous injuries (number, severity, nature and number of months since the most recent ankle injury, use of protective gear such as tape or brace, and if the most recent injury had caused the player to miss matches). The second part was a function score for the ankle (Foot and Ankle Outcome Score; FAOS) (Roos et al., 2001) translated into Norwegian. This form consists of five major parts (symptoms, pain, activities of daily living, function in sports and recreation, quality of life) and is scored by calculating the mean value of the five parts in percent of the total possible score, where 100% is the maximal and 0% the lowest score.

In addition, a similar screening was done for risk factors for hamstring, knee and groin injuries. The data from these tests will be reported in separate papers.

Injury reporting

Each team was supplied with a physiotherapist who was responsible for reporting injuries for all the players on the team throughout the preseason and the season. An injury was defined as any physical complaint sustained by a player that resulted from a soccer match or soccer training, forcing the player to miss or being unable to take full part in future soccer training or match play (“time-loss” injury). Acute injuries were defined as injuries with a sudden onset associated with a known trauma, whereas overuse injuries were those with a gradual onset without any known trauma. Two of the authors were blinded to all other information regarding risk factors and categorized all injuries based on the injury reports from the physiotherapist. For the purpose of the present paper, an injury was classified as ankle sprain if it was recorded as an acute injury of the ankle ligaments. Injuries were classified into three severity categories according to the time it took until the player was fully fit to take part in all types of organized soccer play: minor (1-7 days), moderate (8-28 days) and major (>28 days).

The head coach for every team registered each player's participation in training and the number of minutes played in matches.

Most of the teams from the 1st and 2nd division already had a physical therapist working with the team. In cases where there was no physical therapist attached to the team, we provided them with one. However, the physiotherapist was not required to be present at every training session and match; the degree of follow-up therefore varied from team to team participating in the study.

Reliability testing

Interobserver reliability tests were carried out by different test personnel for both the clinical examination and the single leg balance test by having the same player repeat the same test with different personnel after he had completed the first test. Each examiner was blinded to the other's results. The same scoring system/clinical forms were used at both stations.

Interobserver reliability for the categorical variables in the interpretation of the balance tests and the clinical examination was computed using kappa statistics.

Statistical methods

Exposure to matches and training was calculated by adding the individual duration of all training and match play during the season.

For the continuous dependent variable risk factor analyses, where each leg was the unit of analysis, generalized estimating equations (STATA, version 8; STATA, Texas, U.S.A.) were used, accounting for total individual exposure during the soccer season and for the fact that the left and right foot belonged to the same player. Logistic regression analyses were used to analyse the relationships between per subject first occurrence of calculated dichotomous injury variables and their risk factors.

All risk factor variables were examined in univariate analyses, and those with a P value <0.10 were investigated further in a multivariate model.

Results

The total incidence of injuries during the season was 4.7 injuries per 1000 playing hours (95% CI 4.3 to 5.1), 12.1 (95% CI 10.5 to 13.7) for match injuries and 2.7 (95% CI 2.4 to 3.1) for training injuries. The total exposure to match play and training was 108 111 player hours. A total of 56 acute ankle injuries were reported, affecting 46 legs on 43 (8.5%) of the 508 players in the study. The total incidence of acute ankle injuries was 0.5 injuries per 1000 playing hours (95% CI 0.4 to 0.7), 0.3 injuries per 1000 training hours (95% CI 0.2 to 0.4) and 1.5 injuries per 1000 match hours (95% CI 0.9 to 2.0). A total of 34 players sustained one ankle injury, while 6 and 2 players sustained two and three injuries, respectively. One player sustained four ankle injuries throughout the season. Of the 56 injuries, 34 occurred on the right side, while 22 were on the left. There were 26 minor injuries (time loss 1 to 7 days), 22 moderate injuries (8 to 28 days) and 5 severe injuries (>28 days). In 3 cases, information on the duration of time loss was missing.

Interobserver reliability for the categorical variables, computed using kappa statistics, were 0.40 and 0.19 for balance tests on the floor and mat, respectively. For the clinical examination, kappa values were 0.45 (anterior drawer), 0.84 (foot type), 0.91 (standing rearfoot alignment), 1.00 (hallux position), and 1.00 (toe deformity).

Univariate analyses revealed the number of previous acute ankle injuries and the FAOS sub score "Pain" as potential leg-dependent risk factors for acute ankle injuries (Table 1). None of the balance tests, floor or balance mat, or clinical tests were candidates for predicting increased risk of ankle injury. Additionally, none of the player-dependent factors (age, height, body mass index, position on the field, having played at the junior national team or at the

senior national team level, level of play this season or level of play the previous season) were significantly associated with the risk of ankle injury (Table 2).

Risk factors with p-value of <0.10 were then considered as candidates to predict which players are more prone to sustain an acute injury to the ankle. Since these factors may be inter-correlated, a multivariate analysis was performed, and only previous acute ankle injury was found to be a significant risk factor for new acute ankle (Table 3). The importance of this risk factor increases with number of previous injuries (test of trend, $P=0.001$), and seems to decrease with time since the last injury (test of trend, $P=0.06$).

Discussion

The main finding of this cohort study investigating potential risk factors for ankle injuries in soccer was that previous ankle injury was the only significant predictor we could identify for new acute ankle injuries. The risk increases with the number of previous injuries and is highest during the first 6 months after injury. Other candidates for identification of players with increased risk of acute ankle injuries, such as function scores, balance tests, other player characteristics or a clinical examination were not significantly associated with injury risk.

Several authors have found previous ankle injuries to be a significant risk factor for new injuries, both in male soccer (Ekstrand and Gillquist, 1983, Árnason et al., 2004, Tropp et al., 1985, Kofotolis et al., 2007) and among male athletes in other sports (Bahr and Bahr, 1997, McGuine and Keene, 2006, McKay et al., 2001, McHugh et al., 2006, Tyler et al., 2006). Árnason et al. (Árnason et al., 2004) found previous ankle injury to be the only significant risk factor for a new injury to the same ankle in a large cohort study investigating risk factors for soccer injuries. In the same study, lateral instability and a positive anterior drawer test were also correlated with previous injury. In contrast to these findings, Trojian and McKeag (Trojian and McKeag, 2006) and Häggglund et al. (Häggglund et al., 2006) did not find a

history of previous ankle injury to be associated with future ankle sprains. However, a limited number of acute ankle injuries were included in these studies (Trojian and McKeag, 2006, Árnason et al., 2004, Häggglund et al., 2006).

Ankle injuries have been prevented effectively through neuromuscular training, either on a balance board or balance mat, in soccer (Tropp et al., 1985, Árnason et al., 1996) and in other sports (Garrick and Requa, 2005, Bahr et al., 1997, Olsen et al., 2005, McHugh et al., 2007). It therefore seemed reasonable to suggest that a similar exercise could be used as a screening test to identify players at risk. The literature is limited on the topic, and only two publications have looked at whether single leg balance tests can predict risk of new ankle injuries in male soccer (McHugh et al., 2006, Trojian and McKeag, 2006). Trojian and McKeag (Trojian and McKeag, 2006) found a predictive value of balance tests, while McHugh (McHugh et al., 2006) did not. However, several publications looking at balance, measured in different ways, as a predictor of increased risk of injury among male athletes do exist from other sports (Hrysomallis et al., 2007, Tropp et al., 1984, McGuine et al., 2000, Wang et al., 2006, Willems et al., 2005). In the present study, none of the balance tests, on the floor or a balance mat, turned out to be significant predictors. There are several potential explanations for this apparent discrepancy. First, even though this study is one of the largest cohort studies on risk factors for injuries to date, with as many as 56 acute ankle injuries, the statistical power is limited for multivariate tests. Nevertheless, the strength of the candidate risk factors studied does not indicate that any of these would be helpful as screening tools. As pointed out by Bahr & Holme (Bahr and Holme, 2003) in their review, to detect moderate to strong associations 20–50 injury cases are needed, whereas small to moderate associations would need about 200 injured subjects. However, for a risk factor to be clinically relevant with sufficient sensitivity and specificity, strong associations are needed. Second, the results indicate that the intertester reliability for the balance tests used is low, with kappa values of

0.40 and 0.19. This shows that the same player will not necessarily be scored the same way from two different tests of the same ankle, a factor which clearly influences the ability to identify players with reduced ankle control. Third, the floor test has a ceiling effect in this player population, with 97.4% of the subject obtaining a normal or supranormal test score. Because we suspected that this test could be too easy, we also included the balance mat test. For this, the test distribution was better (34.6%, 34.5% and 25.8% in categories 2, 3 and 4, respectively), and the main problem may be that the balance mat test is inconsistent, as indicated by the low kappa value. Also, data from Australian football suggest that balance deficits do not necessarily persist among previously injured athletes (Hrysomallis et al., 2005). To identify athletes at risk based on tests measuring balance and ankle control, we clearly need to develop new methodology with better test properties and reliability. One limitation of the current study is that we had to rely on the coaches for the exposure registration. We had no way to check their figures, but there should be no reason to misreport. If a game or practice session was missed, it would affect all players on the team, which is unlikely to influence the analysis regarding any specific risk factor. The same should be the case for the physiotherapists registering injuries.

Using multivariate methods where we have controlled for significant risk factors as well as player exposure, this study confirms the consistent finding from previous studies that players with a history of ankle sprains are at increased risk (Tropp et al., 1985, Arnason et al., 2004, Ekstrand and Gillquist, 1983, Kofotolis et al., 2007). The high risk period is the first 6 months after a previous injury, as also shown in a study among volleyball players (Bahr and Bahr, 1997). It seems reasonable to recommend that injured players complete a program of balance training on a wobble board for 10 weeks, as first described by Tropp et al. (Tropp et al., 1985), and that they use tape or a brace during high risk activities until their rehabilitation is completed (Ekstrand et al., 1983, Tropp et al., 1985). Studies have shown that taping

(Ekstrand et al., 1983, Tropp et al., 1985) or using an orthotic device (Surve et al., 1994) prevents reinjury in athletes with a history of ankle sprain, but that neither of these methods appear to have any effect on athletes who have not been injured before. This may be due to the manner in which taping and braces apparently work; that is, they improve the ability of the ankle to react quickly to an inversion stress, but not as a passive mechanical support. Following these guidelines may prevent the athlete from entering a vicious circle with repeated ankle sprains and chronic ankle instability problems.

Perspectives

A history of previous acute ankle injury proved to be the only significant risk factor for new injuries to the same ankle in this prospective cohort study among male soccer players. Players with multiple and/or recent injuries are at high risk. For practical use, the sensitivity of previous injury (yes or no) as a predictor for new ankle sprains was 74%, which means 74% of the players who sustained an ankle injury during the season had a history of ankle sprains. However, the positive predictive value was only 6%, which means that only 6% of previously injured players suffered a new ankle sprain during the season. This figure increases gradually with the number of previous injuries to 10%, if the player has had five or more previous acute ankle injuries. The same is the case if there is a history of a recent sprain, i.e. during the last 6 months (9%). Based on these results, it does not seem possible to target preventive measures based on a history of ankle sprains alone. The results from this study also show that additional information such as balance tests, player interviews or clinical examination does not increase our ability to identify players at risk.

Figure 1. Flow chart showing movement of numbers of players participating.




Figure 2. Front (a) and side (b) view of balance test on floor and balance mat respectively.

Players were asked to stand barefoot on one straight leg, keeping his arms crossed across the chest and his other leg bent 90° at the knee, and only using the ankle joint to correct his balance.

Table 1. Risk factor analyses where each leg was the unit of analysis, including both continuous (Mean \pm SEM) and categorical (yes/no) dependent variables. Comparisons of risk factors between ankles that sustained at least one injury during the following season (“Injured”) and ankles that did not (“Uninjured”). P-values are the results from univariate analyses in STATA using generalized estimating equations taking into account the individual exposure and the fact that the left and the right leg belong to the same player.

	Current injury				SD	OR	95% CI	p-value
	Uninjured (n=970)		Injured (n=46)					
	n	n/Mean \pm SEM	n/Mean \pm SEM	% injured				
Previous ankle injury								
Yes	616	582	34	5.5%		1.95	[0.99-3.84]	0.05
No	399	387	12	3.0%		1.00		
Missing	1							
Number of previous injuries ¹								
Average number		1.6 \pm 0.1	2.5 \pm 0.3			1.25	[1.09-1.43]	0.001
No previous injury						1.00		
1 injury	219	210	9	4.1%		0.92	[0.44-1.95]	0.84
2 injuries	145	140	5	3.4%		0.74	[0.29-1.91]	0.54
3 injuries	87	83	4	4.6%		1.02	[0.34-2.97]	0.97
4 injuries	45	41	4	8.9%		2.34	[0.78-7.01]	0.13
5 injuries	25	22	3	12.0%		2.58	[0.69-9.59]	0.16
>5 injuries	95	86	9	9.5%		2.55	[1.17-5.56]	0.02

Time since previous injury (n=1016) ²								0.06	
Never	399	387	12	3.0%		1.00			
0-6 months	137	124	13	9.5%		2.81	[1.42-5.54]	0.003	
6-12 months	114	109	5	4.4%		0.96	[0.37-2.50]	0.93	
1-2 years	141	134	7	5.0%		1.10	[0.47-2.56]	0.83	
>2 years	218	209	9	4.1%		0.89	[0.42-1.90]	0.77	
Missing	7								
FAOS ³ function score									
Total score	902	93 ± 0.3	91 ± 1.7			9.7	0.83	[0.65-1.06]	0.14
Symptoms	931	88 ± 0.4	86 ± 2.2			12.9	0.87	[0.71-1.07]	0.19
Pain	956	96 ± 0.3	93 ± 1.5			9.2	0.81	[0.62-1.04]	0.10
Activities of daily life	957	98 ± 0.2	97 ± 1.3			6.4	0.89	[0.60-1.32]	0.58
Sport	961	94 ± 0.4	92 ± 2.3			13.2	0.92	[0.75-1.11]	0.38
Quality of life	960	90 ± 0.5	87 ± 3.0			15.3	0.88	[0.75-1.04]	0.13
Testing ⁴									
Balance test, floor	999	4.6 ± 0.02	4.7 ± 0.1			0.55	1.08	[0.79-1.48]	0.64
Balance test, mat	999	3.0 ± 0.02	3.2 ± 0.1			0.90	1.14	[0.84-1.54]	0.41
Clinical examination									
Any pathological findings (n=817)									
Yes	427	407	20	4.7%		1.03	[0.75-1.42]	0.85	
No	390	374	16	4.1%		1.00			
Foot type (n=886)								0.78	
Normal	568	543	25	4.4%		1.00			
Pes planus	228	221	7	3.1%		0.69	[0.29-1.61]	0.39	
Pes cavus	73	68	5	6.8%		1.60	[0.59-4.31]	0.36	
Splayed forefoot	17	16	1	5.9%		1.36	[0.17-10.6]	0.77	
Standing rearfoot alignment (valgus) (n=864)									
Yes	134	131	3	2.2%		1.00			
No	730	697	33	4.5%		1.86	[0.56-6.24]	0.31	
Hallux position (valgus) (n=873)									
Yes	76	72	4	5.3%		1.46	[0.49-4.34]	0.50	
No	797	763	34	4.3%		1.00			

Anterior drawer (pathologic) (n=876)								
Yes	138	129	9	6.5%	1.83	[0.85-3.98]	0.13	
No	738	698	29	3.9%	1.00			
Supination (degrees) ⁴	886	28.8° ± 0.6 (848)	35.0° ± 4.5 (38)		19.2	1.21	[0.93-1.57]	0.15
Pronation (degrees) ⁴	884	9.2° ± 0.2 (846)	9.5° ± 0.6 (38)		9.2	0.98	[0.48-2.00]	0.95
Dorsal extention (degrees) ⁴	865	10.4° ± 7.3 (827)	10.1° ± 5.3 (38)		10.3	0.94	[0.60-1.48]	0.79

^a The number of legs in the uninjured and injured groups reflect the number of legs that completed each of the tests.

¹ Results (OR and 95% CI) are presented per previous injury.

² Results (OR and 95% CI) are presented per category increase.

³ FAOS (Foot and Ankle Outcome Score). (Roos et al., 2001) All results (OR and 95% CI) are presented for a change of 10 in FAOS-score.

⁴ Results (OR and 95% CI) are presented per increase of one standard deviation.

Range (mean,min-max) of continuous variables: FAOS (total score: 93.3, 37.2-100.0), (Symptoms: 88.4, 28.6-100.0), (Pain: 95.6, 38.9-100.0), (Activities of daily life: 98.2, 45.6-100.0), (Sport: 94.1, 25.0-100.0), (Quality of Life: 90.1, 6.3-100.0), Balance test on floor (4.6, 1.0-5.0), Balance test on mat (3.1, 1.0-5.0), Supination (29.1, 0-150), Pronation (9.2, 0-30) and Dorsal extention (10.3, 0-90).

Table 2. Risk factor analyses where each player was the unit of analysis, including both continuous (Mean \pm SEM) and categorical (yes/no) dependent variables. Comparison between the players who sustained at least one ankle injury during the following season (“Injured”) and the players who did not (“Uninjured”).

Factor	Current injury				SD	OR	95% CI	p-value
	n	Uninjured (n=465) n/Mean \pm SEM	Injured (n=43) n/Mean \pm SEM	% injured				
Age [†] (years)	500	24.0 \pm 0.2 (458)	24.0 \pm 0.6 (42)		4.2	1.00	[0.85-1.18]	0.99
Height [†] (cm)	497	181.4 \pm 0.3 (455)	181.0 \pm 1.0 (42)		6.3	0.93	[0.68-1.27]	0.66
Weight [†] (kg)	493	78.0 \pm 1.1 (450)	77.9 \pm 0.4 (43)		8.0	1.01	[0.74-1.38]	0.94
BMI [†] (kg * m ⁻²)	486	23.7 \pm 0.1 (444)	23.8 \pm 0.2 (42)		2.1	1.13	[0.76-1.68]	0.56
Player position	485							0.51
Forward	84	78	6	7.1		1.00		
Winger	70	65	5	7.1		1.00	[0.29-3.43]	1.00
Attacking midfielder	62	54	8	12.9		1.93	[0.63-.5.87]	0.25
Central midfielder	66	61	5	7.6		1.07	[0.31-3.66]	0.92
Wingback	87	77	10	11.5		1.69	[0.59-4.87]	0.33
Center back	71	65	6	8.5		1.20	[0.37-3.90]	0.76
Goalkeeper	45	44	1	2.2		0.30	[0.03-2.53]	0.27
Level of play	508							0.89
1st division	119	109	10	8.4		1.00		
2nd division	256	233	23	9.0		1.08	[0.50-2.34]	0.85
3rd division	133	123	10	7.5		0.89	[0.36-2.21]	0.80

Level of play last season	485							0.71
Elite division	4	3	1	25.0	1.00			
1st division	126	115	11	8.7	0.29	[0.03-3.00]		0.30
2nd division	154	141	13	8.4	0.28	[0.03-2.85]		0.28
3rd division or lower	201	184	17	8.5	0.28	[0.03-2.81]		0.28
Junior or senior national team matches	508							
Yes	92	86	6	6.5	0.72	[0.29-1.75]		0.46
No	416	379	37	8.9	1.00			

^a The number of players in the uninjured and injured groups reflect the number of players who completed each of the tests.

¹ Results (OR and 95% CI) are presented per increase of one standard deviation.

Range (mean, min-max) of continuous variables: Age (24.0, 16.2-37.7), Height (181.4, 153-198), Weight (77.9, 56.0-105.0), BMI (23.7, 19.4-29.8).

Table 3. Multivariate analyses of the potential risk factors with $p < 0.10$ in univariate analyses.

Adjusted odds ratio (OR) and 95% confidence interval (CI) of number of previous ankle injuries as continuous variable and per difference of 10 in FAOS (Foot and Ankle Outcome Score) (Roos et al., 2001) sub-score "Pain".

Risk factors	Adjusted OR	95% CI	p-value
Previous ankle injury Per previous ankle injury	1.23	[1.06-1.41]	0.005
FAOS ¹ sub-score "Pain"	0.89	[0.67-1.18]	0.41

¹FAOS (Foot and Ankle Outcome Score) (Roos et al., 2001) (OR and 95% CI) are presented for a change of 10 in FAOS-score

Reference List

- Andersen T E, Floerenes T W, Arnason A, Bahr R. Video analysis of the mechanisms for ankle injuries in football. *Am J Sports Med* 2004; (32): 69S-79S.
- Arnason A, Gudmundsson A, Dahl H A, Johannsson E. Soccer injuries in Iceland. *Scand J Med Sci Sports* 1996; (6): 40-45.
- Arnason A, Sigurdsson S B, Gudmundsson A, Holme I, Engebretsen L, Bahr R. Risk factors for injuries in football. *Am J Sports Med* 2004; (32): 5S-16S.
- Backous D D, Friedl K E, Smith N J, Parr T J, Carpine W D, Jr. Soccer injuries and their relation to physical maturity. *Am J Dis Child* 1988; (142): 839-842.
- Bahr R, Bahr I A. Incidence of acute volleyball injuries: a prospective cohort study of injury mechanisms and risk factors. *Scand J Med Sci Sports* 1997; (7): 166-171.
- Bahr R, Holme I. Risk factors for sports injuries--a methodological approach. *Br J Sports Med* 2003; (37): 384-392.
- Bahr R, Lian O, Bahr I A. A twofold reduction in the incidence of acute ankle sprains in volleyball after the introduction of an injury prevention program: a prospective cohort study. *Scand J Med Sci Sports* 1997; (7): 172-177.
- Baumhauer J F, Alosa D M, Renstrom A F, Trevino S, Beynnon B. A prospective study of ankle injury risk factors. *Am J Sports Med* 1995; (23): 564-570.
- Beynnon B D, Renstrom P A, Alosa D M, Baumhauer J F, Vacek P M. Ankle ligament injury risk factors: a prospective study of college athletes. *J Orthop Res* 2001; (19): 213-220.
- Ekstrand J, Gillquist J. Soccer injuries and their mechanisms: a prospective study. *Med Sci Sports Exerc* 1983; (15): 267-270.
- Ekstrand J, Gillquist J, Liljedahl S O. Prevention of soccer injuries. Supervision by doctor and physiotherapist. *Am J Sports Med* 1983; (11): 116-120.
- Ekstrand J, Tropp H. The incidence of ankle sprains in soccer. *Foot Ankle* 1990; (11): 41-44.
- Engebretsen A H, Myklebust G, Holme I, Engebretsen L, Bahr R. Prevention of injuries among male soccer players: a prospective, randomized intervention study targeting players with previous injuries or reduced function. *Am J Sports Med* 2008; (36): 1052-1060.
- Garrick J G, Requa R. Structured exercises to prevent lower limb injuries in young handball players. *Clin J Sport Med* 2005; (15): 398.
- Hagglund M, Walden M, Ekstrand J. Previous injury as a risk factor for injury in elite football: a prospective study over two consecutive seasons. *Br J Sports Med* 2006; (40): 767-772.

- Hawkins R D, Fuller C W. A prospective epidemiological study of injuries in four English professional football clubs. *Br J Sports Med* 1999; (33): 196-203.
- Hrysomallis C, McLaughlin P, Goodman C. Does a balance deficit persist in Australian Football players with previous lower limb ligament injury? *J Sci Med Sport* 2005; (8): 85-91.
- Hrysomallis C, McLaughlin P, Goodman C. Balance and injury in elite Australian footballers. *Int J Sports Med* 2007; (28): 844-847.
- Juma A H. Outline of sport injuries in the V World Youth Championship for FIFA Cup in Saudi Arabia. *Indian J Med Sci* 1998; (52): 433-437.
- Junge A, Dvorak J, Rösch D, Graf-Baumann T, Chomiak J, Peterson L. Psychological and Sport-Specific Characteristics of Football Players. *Am J Sports Med* 2000; (28): 22-28.
- Junge A, Dvorak J, Graf-Baumann T, Peterson L. Football injuries during FIFA tournaments and the Olympic Games, 1998-2001: development and implementation of an injury-reporting system. *Am J Sports Med* 2004; (32): 80S-89S.
- Kofotolis N D, Kellis E, Vlachopoulos S P. Ankle sprain injuries and risk factors in amateur soccer players during a 2-year period. *Am J Sports Med* 2007; (35): 458-466.
- Lindenfeld T N, Schmitt D J, Hendy M P, Mangine R E, Noyes F R. Incidence of injury in indoor soccer. *Am J Sports Med* 1994; (22): 364-371.
- Lysens R J, Ostyn M S, Vanden A Y, Lefevre J, Vuylsteke M, Renson L. The accident-prone and overuse-prone profiles of the young athlete. *Am J Sports Med* 1989; (17): 612-619.
- McGuine T A, Greene J J, Best T, Levenson G. Balance as a predictor of ankle injuries in high school basketball players. *Clin J Sport Med* 2000; (10): 239-244.
- McGuine T A, Keene J S. The effect of a balance training program on the risk of ankle sprains in high school athletes. *Am J Sports Med* 2006; (34): 1103-1111.
- McHugh M P, Tyler T F, Mirabella M R, Mullaney M J, Nicholas S J. The effectiveness of a balance training intervention in reducing the incidence of noncontact ankle sprains in high school football players. *Am J Sports Med* 2007; (35): 1289-1294.
- McHugh M P, Tyler T F, Tetro D T, Mullaney M J, Nicholas S J. Risk factors for noncontact ankle sprains in high school athletes: the role of hip strength and balance ability. *Am J Sports Med* 2006; (34): 464-470.
- McKay G D, Goldie P A, Payne W R, Oakes B W. Ankle injuries in basketball: injury rate and risk factors. *Br J Sports Med* 2001; (35): 103-108.
- Meeuwisse WH. Assessment of causation in sport injury: A multifactorial model. *Clin J Sport Med* 4, 166-170. 1994.
Ref Type: Journal (Full)
- Olsen O E, Myklebust G, Engebretsen L, Holme I, Bahr R. Exercises to prevent lower limb injuries in youth sports: cluster randomised controlled trial. *BMJ* 2005; (330): 449.

- Ostenberg A, Roos H. Injury risk factors in female European football. A prospective study of 123 players during one season. *Scand J Med Sci Sports* 2000; (10): 279-285.
- Roos E M, Brandsson S, Karlsson J. Validation of the foot and ankle outcome score for ankle ligament reconstruction. *Foot Ankle Int* 2001; (22): 788-794.
- Surve I, Schweltnus M P, Noakes T, Lombard C. A fivefold reduction in the incidence of recurrent ankle sprains in soccer players using the Sport-Stirrup orthosis. *Am J Sports Med* 1994; (22): 601-606.
- Taerk G S. The injury-prone athlete: a psychosocial approach. *J Sports Med Phys Fitness* 1977; (17): 186-194.
- Taimela S, Kujala U M, Osterman K. Intrinsic risk factors and athletic injuries. *Sports Med* 1990; (9): 205-215.
- Trojian T H, McKeag D B. Single leg balance test to identify risk of ankle sprains. *Br J Sports Med* 2006; (40): 610-613.
- Tropp H, Askling C, Gillquist J. Prevention of ankle sprains. *Am J Sports Med* 1985; (13): 259-262.
- Tropp H, Ekstrand J, Gillquist J. Stabilometry in functional instability of the ankle and its value in predicting injury. *Med Sci Sports Exerc* 1984; (16): 64-66.
- Tyler T F, McHugh M P, Mirabella M R, Mullaney M J, Nicholas S J. Risk factors for noncontact ankle sprains in high school football players: the role of previous ankle sprains and body mass index. *Am J Sports Med* 2006; (34): 471-475.
- Wang H K, Chen C H, Shiang T Y, Jan M H, Lin K H. Risk-factor analysis of high school basketball-player ankle injuries: a prospective controlled cohort study evaluating postural sway, ankle strength, and flexibility. *Arch Phys Med Rehabil* 2006; (87): 821-825.
- Willems T M, Witvrouw E, Delbaere K, Mahieu N, De B, I, De C D. Intrinsic risk factors for inversion ankle sprains in male subjects: a prospective study. *Am J Sports Med* 2005; (33): 415-423.