

Bloomquist, K. L., Langberg, H. C., Karlsen, S., Madsgaard, S., Boesen, M., Raastad, T. (2013). Effect of range of motion in heavy load squatting on muscle and tendon adaptations. *European Journal of Applied Physiology*, 113, 2133-2142.

Dette er siste tekst-versjon av artikkelen, og den kan inneholde små forskjeller fra forlagets pdf-versjon. Forlagets pdf-versjon finner du på link.springer.com: <http://dx.doi.org/10.1007/s00421-013-2642-7>

This is the final text version of the article, and it may contain minor differences from the journal's pdf version. The original publication is available at link.springer.com: <http://dx.doi.org/10.1007/s00421-013-2642-7>

Effect of range of motion in heavy load squatting on muscle and tendon adaptations

Bloomquist K³, Langberg H², Karlsen S¹, Madsgaard S¹, Boesen M, Raastad T¹

¹Norwegian School of Sport Sciences, Oslo, Norway

²CopenRehab, Institute of Social Medicine, Department of Public Health and Centre for Healthy Ageing, Faculty of Health Sciences, University of Copenhagen, Denmark

³The University Hospitals Centre for Health Research, Copenhagen University Hospital, Denmark

Short title:

Squat range of motion and musculotendinous adaptations

Key Words:

Resistance training, hypertrophy, patellar tendon, jumping performance

Corresponding author:

Truls Raastad

Email truls.raastad@nih.no

Norwegian School of Sport Sciences

P.O. Box 4014, U.S.

0806 Oslo, Norway

Fax: +47 22234220

Tlf: +47 23262328

Abstract

Purpose Manipulating joint range of motion during squat training may have differential effects on adaptations to strength training with implications for sports and rehabilitation. Consequently, the purpose of this study was to compare the effects of squat training with a short vs. a long range of motion.

Methods Male students (n=17) were randomly assigned to 12 weeks of progressive squat training (repetition matched, repetition maximum sets) performed as either a) deep squat (0-120° of knee flexion); n=8 (DS) or (b) shallow squat (0-60° of knee flexion); n=9 (SS). Strength (1 RM and isometric strength), jump performance, muscle architecture and cross-sectional area (CSA) of the thigh muscles, as well as CSA and collagen synthesis in the patellar tendon, were assessed before and after the intervention.

Results The DS group increased 1RM in both the SS and DS with $\sim 20 \pm 3\%$, while the SS group achieved a $36 \pm 4\%$ increase in the SS, and $9 \pm 2\%$ in the DS ($P < 0.05$). However, the main finding was that DS training resulted in superior increases in front thigh muscle CSA (4-7%) compared to SS training, whereas no differences were observed in patellar tendon CSA. In parallel with the larger increase in front thigh muscle CSA, a superior increase in isometric knee extension strength at $75^\circ (6 \pm 2\%)$ and $105^\circ (8 \pm 1\%)$ knee flexion, and squat-jump performance ($15 \pm 3\%$), were observed in the DS group compared to the SS group.

Conclusion Training deep squats elicited favourable adaptations on knee extensor muscle size and function compared to training shallow squats.

Introduction

Strength training is associated with improvements in muscle strength through adaptations in neural control (Aagaard 2002; Del Balso and Cafarelli 2007), muscle cross-sectional area (CSA) (Wickiewicz et al. 1984; Kawakami et al. 1995; Aagaard et al. 2001), muscle architecture (Blazevich et al. 2003; Aagaard et al. 2001; Alegre et al. 2006), fibre-type transformation (Andersen and Aagaard 2000) and alterations in the length-force characteristics (Abe et al. 2000). Furthermore, muscular adaptations appear to be dependent on loading parameters, volume of exercise, velocity of exercise, and movement intent of the exercises used in training (Hakkinen et al 1985; Thepaut-Mathieu et al. 1988; Weir et al. 1994; Rimmer 2000; Blazevich et al. 2003; Lockie et al. 2003; Markovic et al. 2007).

Improved muscle strength increases the forces distributed from the muscles through the tendons (Kannus 2000) and increases the stress on the connective tissue within the muscle, as well as on the tendons in series with the muscles. It is thus likely that the biomechanical properties of the connective tissue are influenced by the force-generating capacity of the muscles, although little research has been conducted to address this (Kongsgaard et al. 2007; Coupe et al. 2008).

Exercise has been shown to increase the turnover of tissue within tendons both acutely and as a result of a prolonged training intervention (Langberg et al. 1999; Langberg et al 2000; Miller et al. 2005; Kongsgaard et al. 2007; Langberg et al. 2007). Animal studies show that exercise improves the physical properties of tendons, e.g. maximal tensile strength (Elliot 1965; Woo et al. 1981) and newly published data indicate that this also is the case in humans (Haraldsson et al. 2005; Kongsgaard et al. 2007; Coupe et al. 2008)..

Strength training utilizing the squat exercise can be performed in various ways, among those being a full range deep squat (DS) or a limited range shallow squat (SS). To our knowledge, only one

study has explored the effect of squat training at different joint angles (Weiss 2000). Hypothetically adaptations to squat training performed using full- or limited range of motion could lead to differential adaptations, with implications for e.g. power sport performance or during preventative rehabilitation programs against certain musculotendinous injuries.

Theoretically, in maximal lifts the forces on the knee extensors and patellar tendon are the same in both the DS and SS even though the SS can be performed with substantially more load (Figure 1). This is because the external moment arm is approximately twice as long when the femur is parallel to the ground (DS) compared to a limited range of motion of 60° of knee flexion (SS). Thus, assuming that the force on the muscle-tendon system is the same in both ranges of motion, the only difference is the length at which the working muscles contract.

However, changes in the patellar tendon moment arm with increasing knee angles also need to be considered, as the moment arm of the knee extensor muscles is described by the moment arm of the patellar tendon. Peak values for the patellar tendon moment arm are estimated to be near 45° of knee flexion (Krevolin et al. 2004). Interestingly, at 90° of knee flexion the patellar tendon moment arm seems to decrease by approximately 50%, and most likely continues to decrease with increasing knee angles (Krevolin et al. 2004). Assuming this to be true, the DS would thus elicit higher tendon and muscles forces compared to the SS, and hypothetically be a catalyst for patellar tendon hypertrophy and collagen synthesis. This hypothesis is supported by findings reported by Tsaopoulos et al. (2006).

The purpose of this randomised study was therefore to explore whether the DS and SS exercise had a differential effect on specific adaptations in the front thigh muscles and patellar tendon, as well as on jump performance. It was hypothesized that SS training would be superior in eliciting increased strength in the SS. In contrast, DS training would be superior in increasing strength in the DS and front thigh muscle CSA as well as increasing patellar tendon CSA and collagen synthesis.

Furthermore, it was hypothesized that these superior muscle and tendon adaptations with DS training would translate into a more positive effect on jumping performance compared to the SS.

Materials and methods

Subjects It was calculated that ten subjects in each training group would give a statistical power of 90%. Normally, the drop-out rate in training interventions is around 10-20%. Therefore, twenty four males were recruited for the study (Table 1). All subjects were sports science students. If they had been squat training more than once weekly during the preceding six months, or if they were engaged in strength- or power sports, they were excluded from the study. During the intervention, subjects were requested not to participate in endurance sports more than three times per week, or to engage in strength training of the lower extremities. After a one week familiarization period, subjects were tested and paired according to their initial DS strength. From each pair one subject was drawn, by envelope, into either the DS or SS group with the other member of the pair allotted to the opposite group. Four subjects withdrew preceding training. This left 20 subjects, where an additional two subjects withdrew due to illness and injury. Training attendance was set at $\geq 80\%$ and one additional subject was excluded due to a lack of attendance. This left 17 subjects with 9 in the SS group, and 8 in the DS group.

Training Both groups engaged in strength training, three times per week for twelve weeks. Each session started with a ten minute general warm up, followed by a specialized warm- up consisting of 1-3 submaximal squats (shallow or deep according to training group). Both groups performed barbell squat free weight exercises. The SS group performed the squat from complete knee extension (0°) to 60° of knee flexion, and back to extended knee, while the DS group performed a full range of motion squat, with the femur parallel to the floor in the lowest position (120° of knee

flexion) (Figure 1). Both squat variations were executed with an eccentric phase lasting two to four seconds followed by a maximal effort in the concentric phase with the subjects' feet staying on the ground. The training program was periodized and loads progressively increased during the 12 weeks (Table 2). All training sessions were supervised to ensure correct range of motion and safety. The study complied with the Declaration of Helsinki and was approved by the Regional Ethics Committee of Southern Norway.

(Table 1 near here)

(Table 2 near here)

(Figure 1 near here)

Testing procedures Microdialysis and ultrasonography were carried out during the familiarization week, while the remainder of the pretests were carried out the following week. All tests were carried out at pre-intervention and after 12 weeks. Testers were blinded in regard to training group.

1 RM strength All subjects were tested using 1 RM for both the DS and SS after a general and specialized warm-up consisting of a series of 10-6-3-1- repetitions, without subjects reaching fatigue. Based on the last sub-maximal series of 1 repetition, a plausible load was chosen. Hereafter loads were increased with a minimum of 5 or 10 kg and a maximum of 15 or 30 kg, for DS and SS respectively, until the subjects failed to lift the load with correct technique.

Isometric strength Isometric strength of the knee extensors on the right leg were measured in a dynamometer (Technogym REV 9000, Gambettola, Italy) at knee angles of 40°, 75° and 105° (full knee extension at 0°). After a specific warm up with four isokinetic knee extensions with increasing intensity, two maximal contractions of 5 seconds were performed at each knee angle with a 30

second rest between attempts. Peak torque at each knee angle was used for analysis (coefficient of variation (CV) <5%).

Cross Sectional Area (CSA) The CSA of the front thigh muscles (m.sartorius and quadriceps (and adductors in the most proximal sections)), back thigh muscles (hamstrings) and the patellar tendon were obtained using magnetic resonance imaging ((MRI), GE Signa 1.5 Tesla EchoSpeed, GE Medical Systems, Milwaukee, WI). A total of nine slices were analysed for muscle CSA from both legs. The first slice was placed 10 cm proximal to the lateral femoral epicondyle and was defined as the most distal slice. The remaining eight slices were then placed proximally to this reference point with 10 mm between each slice. To measure the CSA of the patellar tendon seven slices per leg were taken. The first slice was placed 5 mm distal to the tibial plateau (reference point). The second slice was placed on the tibial plateau, and the remaining slices were taken proximal to the tibial plateau with 5 mm between each slice.

A line was manually drawn along the perimeter of the muscle bellies and the tendon on each slice, and the CSA was automatically generated in the software (OsiriX 3.9.3, Pixmeo, Bernex, Switzerland).

Lean body mass (LBM) of the legs and body composition were measured using Dual energy x-ray absorption ((DEXA), Lunar Prodigy densitometer, GE Medical Systems, Madison, WI). Subjects were requested not to eat or drink during the two hours preceding the scanning and to eat identical meals at identical times at both pre- and posttest.

Muscle architecture Using ultrasound imaging (Toshiba Sonolayer Just Vision 400) pennation angle and muscle thickness of the right m.vastus lateralis was measured. Ultrasound was performed with subjects relaxed and lying in supine with the knees fully extended. Using a point midway between the greater trochanter and the lateral condyle, isolated muscle thickness and pennation

angle were measured *in vivo* using the ultrasonograph and pictures were stored and blinded. Muscle thickness was determined by measuring the distance between the superficial and deep aponeurosis whereas the pennation angle was defined as the angle between the fascicle and the deep aponeurosis (Alegre et al. 2006). Five measurements were taken. The highest and lowest values were withdrawn, and a mean value was determined from the three remaining measurements. Reliability measurements were not conducted in the present trial, however CV of 3% for muscle thickness and 5% for pennation angle have hence been performed utilizing same analysis procedures and ultrasonograph technicians. The posttest was performed 1 day after a submaximal training session during the last training week.

Collagen synthesis in the patellar tendon Pretest microdialysis was performed, with no exercise or testing done during the preceding two days (Langberg et al. 1999). The post sampling was executed the day after the last training session (16-28 hours after exercise). On each leg one microdialysis catheter, custom-made in the laboratory, was placed in front of the patellar tendon before and after the intervention. The active part of the membrane was 30 mm long covering the width of the patellar tendon. The sterilized (ETO) fibres were all high molecular mass cut-off fibres (3000kDa, membrane length 30mm, catheter outer diameter 0.05mm). *In vivo* recovery was determined using labelled glucose (Glucose D-[3-³H], 250 μ Ci in 2.5 ml ethanol/water (9:1), as no radioactively labelled procollagen type 1 N-propeptide (PINP) was commercial available. The catheters were perfused (CMA 100) at a rate of 2 μ l/min. After the microdialysis catheters had been positioned the subjects rested for at least 90 min before starting the sampling (4 hours), to ensure that the insertion trauma was minimised (Langberg et al. 1999). The samples were immediately frozen at -80° C until analyses were done. Collagen synthesis was analysed as the peritendinous concentration of PINP in the microdialysis samples by a sandwich ELISA (Christensen et al. 2008).

The dialysate samples were diluted: 1:9, 1:10 or 1:20 before the analysis, based on previous analysis of the sample. The detection level was 41 pg/ml and the intra-assay variation (CV) of 4.9 % at 4.2 ng/ml. All the samples from the same subject were analysed on the same ELISA plate.

Jump performance Squat jumps (SJ) and counter movement jumps (CMJ) were performed on a forceplate (SG-9, Advanced Mechanical Technologies, Newton, MA, USA) and low-pass filtered at 1050 Hz. SJ were performed with no counter-movement from a knee angle of 90° with hands fixed at the hip. The CMJ started from a standing position with the hands fixed at the hip. Jump height was calculated from the impulse during takeoff. At each test the subjects did three to six jumps and the best result was used for analysis (CV<5%).

Statistical methods A per protocol-analysis was applied, thus all results are based on the 17 subjects who completed the training intervention. Means, standard deviations (SD) and standard errors (SE) were calculated and all values are presented as means and standard errors unless otherwise noted. The paired t-test was performed in order to assess changes over time within each training group, whereas the un-paired t-test was performed to assess the statistical significance of between group differences. Pearson's correlation coefficient (r) was used to calculate SS and DS collapsed variables. Significance was set at 5% ($P \leq 0.05$).

Results

The pretest characteristics of the subjects are presented in Table 1 with no between group differences.

Strength Both training groups increased 1 RM when tested in the SS and DS (Figure 2). For the DS group an increase of $\sim 20 \pm 3\%$ was observed in both squat ranges ($P < 0.05$), while the SS group achieved a $36 \pm 4\%$ increase in the SS and $9 \pm 2\%$ in the DS ($P < 0.05$). The SS group increased 1 RM in the SS more than the DS group, and the DS group increased 1 RM in the DS more than the SS group ($P < 0.05$). Maximal isometric knee extensor torque increased in the DS group at knee angles of 75° ($6 \pm 2\%$) and 105° ($8 \pm 1\%$) ($P < 0.05$). At 105° the increased torque in the DS group was larger than in the SS group ($8 \pm 1\%$ vs. $0 \pm 5\%$, respectively) ($P < 0.05$) (Figure 3).

(Figure 2 and 3 near here)

Thigh muscle CSA The three most distal slices were a mix of muscle and tendon tissue (especially in the tallest subjects) and were therefore not analysed. The muscle CSA of the front thigh was increased at all measured sites in the DS group (4-7%), while increases at the two most proximal sites were observed in the SS group ($P < 0.05$) (Figure 4, upper panel). The change in muscle CSA at all measured sites was greater in the DS group ($P < 0.05$). The muscle CSA of the back thigh was increased at the second most proximal site in the DS group ($P < 0.05$) (Figure 4, lower panel).

(Figure 4 near here)

Lean body mass LBM of the legs increased by $2.0 \pm 0.8\%$ ($P < 0.05$) in the DS group while no increase was observed in the SS group ($1.5 \pm 0.9\%$) (Figure 5). The DS group achieved increases in body mass (1.7 ± 0.6 kg ($2.2 \pm 0.6\%$) ($P < 0.05$) and total LBM 1.2 ± 0.4 kg ($1.8 \pm 0.8\%$) ($P < 0.05$). No

changes in body mass were detected in the SS group. There were no differences between training groups in body composition, and no changes in fat percent for either group.

(Figure 5 near here)

Muscle architecture No changes in muscle thickness were observed in either training group. However, increases in pennation angle were observed for both the SS and DS group ($23\pm 5\%$ and $22\pm 6\%$) ($P < 0.05$), respectively. There were no differences between groups for changes in muscle thickness or pennation angle (Table 3).

(Table 3 near here)

Tendon CSA and collagen synthesis We found no changes in the CSA of the patellar tendon at any of the measured sites in any group. Collagen synthesis, indicated by the PINP content measured with microdialysis, did not increase after training in either group.

Jump height CMJ height increased by $7\pm 4\%$ in the SS group and $13\pm 2\%$ in the DS group ($P < 0.05$) (Figure 6). SJ height increased in the DS group ($15\pm 3\%$) ($P < 0.05$) and was greater than in the SS group ($P < 0.05$).

(Figure 6 near here)

Correlations.

Front thigh muscle CSA/LBM legs and strength: Correlations between muscle CSA of the front thigh and 1RM DS strength were detected at both pretest and posttest, whereas correlations between muscle CSA of the front thigh and 1RM SS strength only were observed at pretest (Table 4). No correlations were detected between isometric knee extension strength and front thigh muscle CSA.

However, correlations between isometric knee extension strength at both 75° and 105° and LBM of the legs were found at pretest.

Muscle strength and architecture: Correlations between isometric strength at 105° and pennation angle were found at pretest (Table 4).

Front thigh muscle CSA and patellar tendon CSA: No correlations between front thigh muscle CSA and patellar tendon CSA were observed. However, at pretest, a correlation was found between the mean patellar tendon CSA and LBM of the legs. Furthermore, there was a correlation between mid-patellar tendon CSA and maximal isometric knee extensor strength at 105° (Table 4).

Jump height and muscle architecture/strength: A correlation was found between SJ height and isometric knee extensor strength at 105° both at pretest and posttest (Table 4). No correlations between jump performance and muscle architecture were observed. However, a negative correlation was found between pennation angle and the difference between the CMJ and SJ height.

(Table 4 near here)

Discussion

We found that 12 weeks of progressive heavy load squat training, regardless of range of motion, resulted in increases in 1RM strength and pennation angle, as well as increases in CMJ height. However, only the DS group increased SJ height, LBM of the legs, isometric strength at 75° and 105°, and front thigh muscle CSA at all measured points. The SS group elicited front thigh muscle CSA increases only at the two most proximal sites.

Maximal muscle strength In accordance with our hypothesis, the SS exercise elicited larger strength gains in the SS, while the DS exercise resulted in larger strength gains in the DS. It is, however, worth noting that DS training elicited similar results in both the DS and SS. In contrast, isometric knee extension strength measurements revealed no increases of strength in the SS group despite higher loads of training, whereas the DS group achieved increases at both 75° and 105° knee flexion. These results are similar to those reported by Weiss et al. (2000) who concluded that the DS was superior to the SS in regard to strength and squat performance.

In the SS group, the CSA of the front thigh muscles was increased only at the two most proximal sites. In conjunction, no isometric strength gains were found in the SS group, as well as no increases in leg LBM or in back thigh muscle CSA. This indicates that the increases in 1 RM strength observed in the SS group likely were due to neural adaptations and/or to muscular adaptations in muscles not analysed. The two most proximal sites included adductor muscles. Substantially higher loads were lifted by the SS group, and it seems reasonable to assume that this potentially resulted in an increased load on the adductor muscles, with hypertrophy to follow as well as favourable adaptations to other muscles working over the hip. Indeed, it is a study limitation that other muscles (e.g. hip extensors) were not measured, as these may have been affected differentially by the two squat training modes.

The DS group increased front thigh muscle CSA at all measured sites in accordance with our hypothesis. These findings are supported by the increases of leg LBM, and increases in both isometric and 1RM strength in the DS group as well as pre- and posttest correlations between 1RM strength and front thigh CSA. Estimations of the external moment arms in the DS and SS showed that in the deepest position the external moment arm was approximately twice as long in the DS as in the SS (Figure 1). This difference in external moment arm corresponded with the observed doubling of external load that could be lifted in the SS compared to the DS exercise. However, the patellar tendon moment arm is reduced when knee flexion is above 60° (deepest position in the SS) (Kremlin et al. 2004). Consequently, although the knee joint torque was similar in the deepest positions, the muscle force (and tendon force) working in the DS was probably 50-100% higher than in the SS training due to the 25-50% shorter patellar tendon moment arm in this position (Kremlin et al. 2004). Furthermore, when body mass is taken into consideration, the difference in muscle and tendon force between the two conditions may have been even larger. The greater hypertrophy observed in the extensor muscles in the DS group may therefore partly be explained by the larger muscle force developed in the DS training.

Muscle Architecture Both training groups achieved increases in muscle thickness and pennation angle of the m.vastus lateralis. There were no differences between groups, indicating no differential muscle architecture adaptations in response to the two different ranges of motion. As stated above, front thigh muscle CSA increases were larger in the DS than in the SS group, indicating that other muscle bellies could have been more affected by DS training than the m. vastus lateralis.

In the present study a correlation between pennation angle and isometric strength at 105° was observed at pretest, and though no posttest correlation was found, it is reasonable to assume that the increases in pennation angle in both training groups may have had a positive influence on the observed strength development.

Patellar tendon properties We had expected to find increases of patellar tendon CSA and collagen synthesis as a result of the squat training. However, neither group elicited gains in patellar tendon CSA or collagen synthesis. At pretest, correlations were found between the patellar tendon CSA and leg LBM and strength consistent with previous studies that have shown a relationship between muscle strength/size and patellar CSA (Elliot 1965; Kongsgaard et al. 2007). However, no change in patellar tendon CSA was observed in either group despite the increase in strength and CSA of the adjacent muscle, nor was any correlations found. Though not expected, these results are in accordance with several resistance training studies that have shown that increases in strength were not accompanied by increases in tendon CSA (Reeves et al. 2003; Kubo et al. 2007; Seynnes et al. 2009). Rather, a markedly altered elastic modulus was found in these studies, implying a change in the composition of the tendon structure instead of the size. However, in the present study no changes in collagen synthesis were observed which may indicate that 12 weeks of single-mode squat training is not sufficient time to generate a detectable change in patellar tendon CSA, nor collagen synthesis, in well trained subjects. Future studies are warranted to better understand the coordinated response of muscle and tendon as this knowledge could be of significant value in preventing overuse injuries of the tendon.

Jump height CMJ height increased in both groups, however only the DS group achieved gains in the SJ. As summarized by Earp et al. (2010), CMJ is a movement that involves a stretch shortening cycle (SSC) that allows the body to store and redirect energy through an eccentric movement quickly followed by a concentric movement. Due to the SSC, more force and power can be generated during the concentric phase of the jump, than if no eccentric phase was involved. This is seen when comparing the CMJ height to the SJ height.

We found no correlations between front thigh muscle CSA or strength gains and CMJ jump height in the present study. However improvements in jumping performance could be caused by changes

in muscle CSA only in variable individual combinations with changes in pennation angle, as these two variables are governing factors for the training-induced change in physiological CSA, and hence muscle force. As the SJ lacks the eccentric phase, and therefore the benefits of the SSC, the SJ is more dependent on maximal muscle strength. In the present study we observed no between group differences in CMJ height. In contrast, the DS group was superior to the SS group in the SJ. In addition, increases in isometric strength were largest in the DS group and a correlation both at pre- and posttest, were found between SJ height and isometric strength at 105°. Thus, the increases in strength achieved by the DS group, could account for the superior SJ performance. This positive relationship between increases in SJ height and strength underlines that increases in isolated muscle strength are beneficial for jump performance despite the increase in LBM and body mass due to muscle hypertrophy.

Conclusion

In conclusion, we found that 12 weeks of heavy load DS training was superior to the SS in regard to increases in front thigh muscle CSA and leg LBM, with no changes in patellar tendon CSA. Parallel with these adaptations, superior changes in isometric knee extension strength and SJ performance were observed with DS training. In both groups increases in 1RM squat strength and CMJ height were observed. We suggest that larger muscle-tendon forces over the knee joint, more internal (patellar tendon) work produced, and longer muscle length of the knee extensors during the DS compared to the SS exercise, were the main explanations for the superior adaptations observed with DS training in this study. Adaptations in muscles involved in hip extension and stabilization were not measured in this study, and possible favourable adaptations with SS training in these areas can therefore not be ruled out.

Acknowledgements

The authors express their thanks to the subjects for their time and effort, and a special thank goes to Oliver Faul and Tron Krosshaug for the construction of the DS and the SS figures.

Conflict of Interest

No conflicts of interest, financial or otherwise are declared by the authors(s).

References

- Aagaard P, Andersen JL, Dyhre-Poulsen P, Leffers AM, Wagner A, Magnusson SP, Halkjaer-Kristensen J, Simonsen EB (2001) A mechanism for increased contractile strength of human pennate muscle in response to strength training: Changes in muscle architecture. *J Physiol* 534:613-623
- Aagaard P, Simonsen EB, Andersen JL, Magnusson P, Dyhre-Poulsen P (2002) Neural adaptation to resistance training: Changes in evoked v-wave and h-reflex responses. *J Appl Physiol* 92:2309-2318
- Abe T, Kumagai K, Brechue WF (2000) Fascicle length of leg muscles is greater in sprinters than distance runners. *Med Sci Sports Exerc* 32:1125-1129
- Alegre LM, Jimenez F, Gonzalo-Orden JM, Martin-Acero R, Aguado X (2006) Effects of dynamic resistance training on fascicle length and isometric strength. *J Sports Sci* 24:501-508
- Andersen JL, Aagaard P (2000) Myosin heavy chain iix overshoot in human skeletal muscle. *Muscle Nerve* 23:1095-1104
- Blazevich AJ, Gill ND, Bronks R, Newton RU (2003) Training-specific muscle architecture adaptation after 5-wk training in athletes. *Med Sci Sports Exerc* 35:2013-2022
- Christensen B, Dyrberg E, Aagaard P, Kjaer M, Langberg H (2008) Short-term immobilization and recovery affect skeletal muscle but not collagen tissue turnover in humans. *J Appl Physiol* 105:1845-1851
- Coupe C, Kongsgaard M, Aagaard P, Hansen P, Bojsen-Moller J, Kjaer M, Magnusson SP (2008) Habitual loading results in tendon hypertrophy and increased stiffness of the human patellar tendon. *J Appl Physiol* 105:805-810
- Del Balso C, Cafarelli E (2007) Adaptations in the activation of human skeletal muscle induced by short-term isometric resistance training. *J Appl Physiol* 103:402-411

- Earp JE, Joseph M, Kraemer WJ, Newton RU, Comstock BA, Fragala MS, Dunn-Lewis C, Solomon-Hill G, Penwell ZR, Powell MD, Volek JS, Denegar CR, Hakkinen K, Maresh CM (2010) Lower-body muscle structure and its role in jump performance during squat, countermovement, and depth drop jumps. *J Strength Cond Res* 24:722-729
- Elliott (1965) Structure and function of mammalian tendon. *Biol Rev* 40:392-421
- Hakkinen K, Komi PV, Alen M (1985) Effect of explosive type strength training on isometric force- and relaxation-time, electromyographic and muscle fibre characteristics of leg extensor muscles. *Acta Physiol Scand* 125:587-600
- Haraldsson BT, Aagaard P, Krogsgaard M, Alkjaer T, Kjaer M, Magnusson SP (2005) Region-specific mechanical properties of the human patella tendon. *J Appl Physiol* 98:1006-1012
- Kannus P (2000) Structure of the tendon connective tissue. *Scand J Med Sci Sports* 10:312-320
- Kawakami Y, Abe T, Kuno SY, Fukunaga T (1995) Training-induced changes in muscle architecture and specific tension. *Eur J Appl Physiol Occup Physiol* 72:37-43
- Kongsgaard M, Reitelseder S, Pedersen TG, Holm L, Aagaard P, Kjaer M, Magnusson SP (2007) Region specific patellar tendon hypertrophy in humans following resistance training. *Acta Physiol (Oxf)* 191:111-121
- Krevolin JL, Pandy MG, Pearce JC (2004) Moment arm of the patellar tendon in the human knee. *J Biomech* 37:785-788
- Kubo K, Morimoto M, Komuro T, Tsunoda N, Kanehisa H, Fukunaga T (2007) Influences of tendon stiffness, joint stiffness, and electromyographic activity on jump performances using single joint. *Eur J Appl Physiol* 99:235-243
- Langberg H, Ellingsgaard H, Madsen T, Jansson J, Magnusson SP, Aagaard P, Kjaer M (2007) Eccentric rehabilitation exercise increases peritendinous type I collagen synthesis in humans with achilles tendinosis. *Scand J Med Sci Sports* 17:61-66

- Langberg H, Skovgaard D, Asp S, Kjaer M (2000) Time pattern of exercise-induced changes in type I collagen turnover after prolonged endurance exercise in humans. *Calcif Tissue Int* 67:41-44
- Langberg H, Skovgaard D, Petersen LJ, Bulow J, Kjaer M (1999) Type I collagen synthesis and degradation in peritendinous tissue after exercise determined by microdialysis in humans. *J Physiol* 521 Pt 1:299-306
- Lockie RG, Murphy AJ, Spinks CD (2003) Effects of resisted sled towing on sprint kinematics in field-sport athletes. *J Strength Cond Res* 17:760-767
- Markovic G, Jukic I, Milanovic D, Metikos D (2007) Effects of sprint and plyometric training on muscle function and athletic performance. *J Strength Cond Res* 21:543-549
- Miller BF, Olesen JL, Hansen M, Dossing S, Crameri RM, Welling RJ, Langberg H, Flyvbjerg A, Kjaer M, Babraj JA, Smith K, Rennie MJ (2005) Coordinated collagen and muscle protein synthesis in human patella tendon and quadriceps muscle after exercise. *J Physiol* 567:1021-1033
- Reeves ND, Narici MV, Maganaris CN (2003) Strength training alters the viscoelastic properties of tendons in elderly humans. *Muscle Nerve* 28:74-81
- Rimmer E, G. S (2000) Effects of a plyometrics intervention program on sprint performance. *J Strength Cond Res* 14:295-301
- Seynnes OR, Erskine RM, Maganaris CN, Longo S, Simoneau EM, Grosset JF, Narici MV (2009) Training-induced changes in structural and mechanical properties of the patellar tendon are related to muscle hypertrophy but not to strength gains. *J Appl Physiol* 107:523-530
- Thepaut-Mathieu C, Van Hoecke J, Maton B (1988) Myoelectrical and mechanical changes linked to length specificity during isometric training. *J Appl Physiol* 64:1500-1505
- Tsaopoulos DE, Baltzopoulos V, Maganaris CN (2006) Human patellar tendon moment arm length: Measurement considerations and clinical implications for joint loading assessment. *Clin Biomech (Bristol, Avon)* 21:657-667

Weir JP, Housh TJ, Weir LL (1994) Electromyographic evaluation of joint angle specificity and cross-training after isometric training. *J Appl Physiol* 77:197-201

Weiss LW FA, Wood LE, Relyea GE, Melton C (2000) Comparative effects of deep versus shallow squat and leg-press training on vertical jumping ability and related factors. *J Strength Cond Res* 14:241-247

Wickiewicz TL, Roy RR, Powell PL, Perrine JJ, Edgerton VR (1984) Muscle architecture and force-velocity relationships in humans. *J Appl Physiol* 57:435-443

Woo SL, Gomez MA, Amiel D, Ritter MA, Gelberman RH, Akeson WH (1981) The effects of exercise on the biomechanical and biochemical properties of swine digital flexor tendons. *J Biomech Eng* 103:51-56

Abbreviations

Coefficient of variation	CV
Counter Movement jump	CJ
Cross-sectional area	CSA
Deep squat	DS
Dual energy x-ray absorption	DEXA
Lean Body Mass	LBM
Magnetic resonance imaging	MRI
Pearson correlation coefficient	r
Procollagen type 1 N-propeptide	PINP
Repetition maximum	RM
Series elastic component	SEC
Shallow squat	SS
Squat jump	SJ
Standard deviation	SD
Standard error	SE
Stretch shortening cycle	SSC

Tables

Table 1. Pretest characteristics of subjects in the SS group and in the DS group (means \pm SD).

	Shallow squat group (n=9)	Deep squat group (n=8)
Age (years)	23 \pm 3	25 \pm 6
Weight (kg)	80 \pm 15	79 \pm 6
Height (cm)	178 \pm 6	181 \pm 5
Peak torque (Nm) (isometric at 105°)	241 \pm 66	242 \pm 29
Jump height (cm) (squat jump)	33.9 \pm 3.6	32.8 \pm 3.3
Muscle CSA (cm ²) (front thigh)	95.6 \pm 14.1	95.2 \pm 7.3
Tendon CSA (mm ²) (middle part)	162 \pm 9	166 \pm 12

Table 2. Periodization and progression of strength training

Week	Monday	Wednesday	Friday
1	Familiarization	Familiarization	Familiarization
2	Pretesting	Pretesting	Pretesting
3	3x10 RM	3x8 (submax)*	4x5 RM
4	3x10 RM	3x10 (submax)	4x5 RM
5	3x10 RM	3x8 (submax)	4x5 RM
6	3x10 RM	3x10 (submax)	4x5 RM
7	3x10 RM	3x8 (submax)	4x5 RM
8	3x10 RM	3x10 (submax)	4x5 RM
9	3x6 RM	3x8 (submax)	5x3 RM
10	3x6 RM	3x10 (submax)	5x3 RM
11	3x6 RM	3x8 (submax)	5x3 RM
12	3x6 RM	3x10 (submax)	5x3 RM
13	3x6 RM	3x8 (submax)	5x3 RM
14	3x6 RM	3x10 (submax)	5x3 RM
15	Posttesting	Posttesting	Posttesting

* 8 reps with a 12-13 RM load

Table 3. Muscle architecture before and after training in the SS and DS group. Pre-post values given as means \pm SD; % change \pm SEM.

	Shallow squat group		Deep squat group	
	Pre	Post	Pre	Post
Muscle thickness (cm)	2,47 \pm 0,37	2,54 \pm 0,33	2,51 \pm 0,26	2,60 \pm 0,32
Fasicle angle (°)	18,5 \pm 3,0	22,6 \pm 3,7*	18,6 \pm 2,8	21,7 \pm 2,0*

* P < 0.05

Table 4. Correlation of selected outcomes at pre- and posttest.

Outcome	Outcome	Pretest	Posttest Δ
Isometric strength at 105°	Leg LBM	r = 0.79*	r = 0.29
Isometric strength at 75°	Leg LBM	r = 0.75*	r = -0.03
1 RM deep squat	Front thigh muscle CSA	r = 0.66*	r = 0.53*
1 RM shallow squat	Front thigh muscle CSA	r = 0.77*	r = -0.22
Mean patellar tendon CSA	Leg LBM	r = 0.56*	r = 0.11
Mid patellar tendon CSA	Isometric strength at 105°	r = 0.60*	r = 0.29
SJ height	Isometric strength at 105°	r = 0.54*	r = 0.55*
SJ height-CMJ height	Pennation angle	r = 0.66*	
Isometric strength at 105°	Pennation angle	r = 0.53*	r = 0.06

* P < 0.05

Figure Captions

Fig. 1 Illustration of the deepest position in the SS (left) and DS exercise (right) The external moment arms indicated are estimated from an average subject with regard to lifting technique and height (180 cm) The ground reaction forces represent a body mass of 80 kg and an external load of 200 kg in the SS, and 100 kg in the DS

Fig. 2 One repetition maximum (1 RM) in the DS and SS exercise measured pre- and post intervention * = significant change from pretest, # = significant difference between groups from pre to posttest

Fig. 3 Change in isometric knee-extension peak torque measured at knee angles of 40°, 75° and 105° (0° is full extension) * = significant change from pretest, # = significant difference between groups from pre- to posttest

Fig. 4 Change in front thigh muscle CSA (upper panel) and back thigh (lower panel) * = significant change from pretest (section 9 was the most proximal)

Fig. 5 Leg lean body mass (LBM) pre- and post intervention* = significant change from pretest

Fig. 6 Change in jump height in SJ and CMJ * = significant change from pretest, # = significant difference between groups from pre- to posttest

Illustrations

Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.




Figure 6.




Figure 7.

