

Dalene, K. E., Anderssen, S. A., Ekelund, U., Thorén, A.-K. H., Hansen, B. H., Kolle, E. (2016). Permanent play facility provision is associated with children's time spent sedentary and in light physical activity during school hours: A cross-sectional study. *Preventive Medicine Reports*, 4, s. 429-434.

Dette er siste tekst-versjon av artikkelen, og den kan inneholde små forskjeller fra forlagets pdf-versjon. Forlagets pdf-versjon finner du på www.sciencedirect.com: <http://dx.doi.org/10.1016/j.pmedr.2016.08.011>

This is the final text version of the article, and it may contain minor differences from the journal's pdf version. The original publication is available at www.sciencedirect.com: <http://dx.doi.org/10.1016/j.pmedr.2016.08.011>

Permanent play facility provision is associated with children's time spent sedentary and in light physical activity during school hours: a cross-sectional study

Knut Eirik Dalene*^a, Sigmund A. Anderssen^a, Ulf Ekelund^a, Anne-Karine Halvorsen Thorén^b, Bjørge H. Hansen^a, Elin Kolle^a.

Address: ^a Department of Sports Medicine, the Norwegian School of Sport Sciences, P.O. Box 4014 Ullevål Stadion, N-0806 Oslo, Norway and ^b Department of Landscape Architecture and Spatial Planning, the Norwegian University of Life Sciences, P.O. Box 5003 NMBU, N-1432 Ås.

Email: Knut Eirik Dalene* – k.e.dalene@nih.no; Sigmund A. Anderssen – sigmund.anderssen@nih.no; Ulf Ekelund – ulf.ekelund@nih.no; Anne-Karine Halvorsen Thorén - kine.thoren@nmbu.no; Bjørge H. Hansen – b.h.hansen@nih.no; Elin Kolle – elin.kolle@nih.no

* Corresponding author

List of abbreviations¹

Word counts:

- *Abstract: 227 words*
- *Main text: 3496 words*

¹ PA: Physical Activity; PE: Physical Education; CPM: Counts per Minute; LPA: Light Physical Activity; MVPA: Moderate-to-Vigorous Physical Activity; SES: Socioeconomic Status; CI: Confidence Interval; BMI: Body Mass Index; SOPA: School's Outdoor Play area(s); PPF: Permanent Play Facilities

Abstract

Objective: To study the associations between: 1) number of permanent outdoor play facilities per pupil and 2) the size of the outdoor play area per pupil with sedentary time and physical activity (PA) during school hours in six-, nine-, and 15-year olds. We conducted a cross-sectional study of nationally representative samples of Norwegian six- (n=1071), nine- (n=1421) and 15-year-olds (n=1106) in 2011 (the Physical Activity among Norwegian Children Study). The participation rate was 56.4%, 73.1% and 57.8% for six-, nine- and 15-year olds, respectively. We assessed PA objectively for seven consecutive days using accelerometers, the size of a school's outdoor play area (SOPA) using an online map service and the permanent play facility (PPF) provision using a standardized form during school site visits. We successfully measured SOPA and PPF in 99 schools, from which 3040 participants provided valid accelerometer data. We used generalized least-squares random-effects models with robust variance estimation to assess associations. Our results indicate that better provision of permanent play facilities may reduce sedentary time and increase time spent in light PA among six-year-olds. Permanent play facility provision was not associated with sedentary time or PA among nine- and 15-year-olds. Associations found between outdoor play area size, physical activity and sedentary time were negligible. Future research should investigate what types of permanent play facilities may be associated with physical activity in both children and adolescents.

Key words: PHYSICAL ACTIVITY, CHILD, ADOLESCENT, SCHOOL, RECESS.

1 Introduction

Since almost all children spend a large proportion of their awake time in school, this arena provides a unique setting for physical activity (PA) promotion. During and adjacent to the school day, children may have several opportunities to be physically active, e.g. through active travel, physical education (PE) and recess. Intervention studies aimed at promoting PA in all these settings have shown promising results (Lonsdale et al. 2013;Larouche et al. 2014;Ickes et al. 2013). However, because it is already compulsory in most schools and does not compete with academic interests (Ickes et al. 2013), recess might be a particularly attractive arena for PA promotion. Children also seem to be more physically active in school free play than during PE lessons (Sleap and Warburton 1996), and more physically active outdoors compared with indoors (Gray et al. 2015). Unstructured free play during recess has

been shown to contribute 5-40% of recommended daily PA (Ridgers et al. 2006), indicating that some schools might have a large PA promoting potential through simple, low-cost strategies.

Designing outdoor play areas that stimulate as many pupils as possible to be physically active is, however, a multifaceted process. For example, studies indicate that girls and boys use different areas of their school's outdoor play area (SOPA) when being physically active (Fjørtoft et al. 2009; Anthamatten et al. 2014), that PA levels are higher in areas with a naturalistic feel (Fjørtoft 2004) and that colorful playground markings can increase recess PA (Blaes et al. 2013). Both the size of SOPA and the availability of permanent play facilities (PPF) are basic components of a schoolyard design, and studies indicate that both factors may be important to stimulate PA (D'Haese et al. 2013; Escalante et al. 2012; Nielsen et al. 2010). However, previous research is limited by the use of subjective measures of PA and small sample sizes (Haug et al. 2010; Ridgers et al. 2010b). Furthermore, studies investigating the association between the size and PPF content of SOPA with time spent sedentary among children and adolescents are limited. Even though debated, studies have indicated that sedentary time might pose a negative effect on cardiovascular risk factors already at a young age (Healy and Owen 2010). Therefore, further research is necessary to identify the importance of the size and PPF content of SOPA for both PA and sedentary time.

Therefore, the aim of this study was to assess the associations between: 1) number of permanent play facilities and 2) the size of the outdoor play area with objectively measured sedentary time and physical activity during school hours in a representative sample of pupils from Norwegian schools.

2 Materials and methods

2.1 Participants

The participants in this cross-sectional study, the Physical Activity among Norwegian Children Study, were nationally representative samples of six-, nine- and 15-year-olds. Statistics Norway randomly selected the cohort using cluster sampling, with school as the primary unit. When a school agreed to participate, we invited all pupils in first, fourth or tenth grade to participate. In total, 5,757 pupils from 107 schools were invited. We obtained written informed consent from 3,598 participants and their primary guardians, yielding a participation rate of 56.4%, 73.1% and 57.8% for six-, nine- and 15-year-olds, respectively. The Regional

Committee for Medical Research Ethics and the Norwegian Social Science Data Services reviewed and approved the study. We conducted the study according to the Helsinki declaration.

2.2 Anthropometrics

We measured weight and height to the nearest 0.1 kg (Seca 877, SECA GmbH, Hamburg, Germany) and 0.1 cm (wall-mounted measuring tape), respectively, while the participants wore light clothing and no shoes. Body mass index (BMI) was calculated as kg/m^2 .

2.3 Physical activity

We measured PA using ActiGraph accelerometers (models GT1M and GT3X+; ActiGraph, LLC, Pensacola, Florida, USA). Children's free-living PA measured with ActiGraph accelerometers has previously been shown to correlate moderately well with activity energy expenditure measured by doubly labeled water ($r = 0.66$, $p < 0.001$) (Ekelund et al. 2001). The participants were fitted with the accelerometers on their right hip during school visits, and instructed to wear the monitor during all waking hours for seven consecutive days, except during showering and bathing. Using the Actilife software (ActiGraph, LLC, Pensacola, Florida, USA), we initialized the accelerometers to sample vertical accelerations (30 Hz), and to start recording at 06:00 on the day after the monitors were attached in order to eliminate reactivity-bias (Dossegger et al. 2014). We used KineSoft (KineSoft Software, Rothesay, New Brunswick, Canada) to analyze the accelerometer files.

An epoch length of 10 seconds was used, which has been deemed suitable for children (McClain et al. 2008). We defined non-wear as intervals ≥ 20 consecutive minutes with no activity recordings, and wear time by subtracting non-wear from school hours. In Norway, school normally starts between 8:00 and 9:00 and ends between 13:00 and 14:45, depending on school and grade. To ensure that we only included school hours, we defined schooldays as 9:00-13:00 for six- and nine-year-olds and 9:00-14:00 for 15-year-olds. These periods include morning-, lunch- and afternoon recess for all grades. We excluded all schooldays with ≥ 60 minutes of non-wear and included participants if they had accumulated ≥ 2 valid schooldays of accelerometer data. We collected all data from March to December in 2011 (no measures in July due to summer holidays). Measurements were evenly distributed across the school year, with the exception of August and December during which only 82 and 95 pupils were measured, respectively.

We used counts·min⁻¹ (CPM) as a measure of overall school PA. We calculated CPM by dividing the total number of school day counts by the total number of school day wear minutes. To investigate time spent sedentary, in PA of light intensity (LPA) and of moderate-to-vigorous intensity (MVPA), we used cut-points of <100 CPM (1-1.5 METs), 100-1999 CPM (1.6-2.9 METs) and ≥2000 CPM (≥3 METs), respectively (Andersen et al. 2006).

2.4 Play facilities/Area size

During school visits, the research team registered the number of PPF using a standardized form. Subsequently, we calculated the number of PPF per pupil. To measure the size of SOPA we used a polygon measurement tool and updated electronic maps from the Norwegian Mapping Authority (finn.no 2011). We calculated SOPA by subtracting areas of buildings, car parks and other areas with car traffic from the school's total outdoor area, and then calculated the SOPA per pupil. Others have used similar methods (Pagels et al. 2014; Ridgers et al. 2010a; Nilsen 2014).

Through interviews with teachers, we received information on recess period organization potentially influencing the availability of space and play facilities (e.g. access to areas outside school property and sectioning of SOPA during recess).

2.5 Socioeconomic status

We used the highest education level of the participant's parents (data from Statistics Norway) as a proxy for socioeconomic status (SES) and computed four SES groups: low (primary school, lower secondary school, vocational high school), middle low (secondary school/high school), middle high (undergraduate degree) and high (graduate degree).

2.6 Sample size calculations

We based the sample size calculations on the ability to detect subgroup differences in CPM. With respect to this, 516 individuals in each age and sex group allowed us to detect subgroup differences of 7% using a two-tailed test ($1 - \beta = 0.80$; two-tailed $\alpha = 0.05$). Because of cluster sampling, we incorporated a design effect of 1.1, yielding a final target sample size of 567 individuals in each age and sex group.

2.7 Statistical analysis

We performed all statistical analysis using Stata 13.1 (StataCorp. 2013. Stata Statistical Software: TX: StataCorp LP.). We used independent samples t-test to investigate sex differences, and one-way ANOVA with Bonferroni corrections to assess differences between the three age groups. For our main analyzes, we ran all the models separately for the different age groups. To account for cluster sampling, we used GLS-re models with robust variance estimation. Initially, we entered the interaction terms *sex*number of PPF* and *sex*play area size*. The interaction terms were not statistically significant. Consequently, we did not stratify the main analyses by sex but rather included sex as a covariate.

We adjusted all analyses for accelerometer wear time (except analyses of CPM), measurement month, sex, and SES, and the dummy variables "access to areas outside school property", "sectioning of play areas", "recess at different time points for different classes" and "allowed to spend recess indoors". We also adjusted for number of PPF in analyses with the size of SOPA as the independent variable.

3 Results

Of the 3,598 participants, 3,040 from 99 schools met the inclusion criteria. Because of construction work, we did not get valid measurements in three schools (n=212). The remainder of the excluded participants did not provide valid PA measurements (n=346). Table 1 displays descriptive characteristics of the study sample. Participants meeting the inclusion criteria were similar to those who did not in terms of age and BMI. However, a higher proportion of the excluded six- and nine-year-old participants were categorized in the two lowest SES categories. In general, there were only small differences in BMI and SES between boys and girls within the age groups.

3.1 Physical activity

The participants provided 4.2 ± 0.9 valid schooldays of PA measurements (mean \pm SD). Table 2 displays the participants' school day PA and sedentary time. School day PA and sedentary time were significantly different between all the age groups ($p < 0.001$) and significantly different between girls and boys within the age groups ($p < 0.001$). For the six-, nine- and 15-year-olds, the mean \pm SD proportion of weekday time spent sedentary accumulated during school hours was $31 \pm 5\%$, $31 \pm 5\%$ and $38 \pm 7\%$, respectively. The mean \pm SD proportion of weekday MVPA accumulated during school hours was $36 \pm 8\%$, $35 \pm 10\%$ and $36 \pm 15\%$, respectively.

3.2 Permanent play facilities

We registered more than 50 unique PPF across the participating schools. Swings (94.5%), climbing frames (87.9%), soccer goals (85.5%) and sand boxes (79.5%) were the most common permanent play facilities in primary schools. The most common permanent play facilities in lower secondary schools were basketball hoops (85.3%), soccer goals (79.0%) and beach volleyball nets (33.5%). The absolute number of PPF and the number of PPF per pupil in primary schools were significantly higher than in lower secondary schools ($p < 0.001$) (Table 3).

The participants' overall PA and time spent in MVPA were not associated with the number of PPF per pupil. Among six-year-olds, however, there was a significant negative association between the number of PPF per pupil and time spent sedentary and a significant positive association between the number of PPF per pupil and time spent in LPA. These associations translate to daily changes in time spent sedentary and in LPA of -3.8 and 2.2 minutes, respectively, if the number of PPF per pupil increased from 0.1 to 0.2 (Table 4).

3.3 Outdoor play area size

The size of SOPA per pupil in primary schools was significantly larger than in lower secondary schools ($p < 0.001$) (Table 3). For the six- and nine-year-olds, we did not find an association between the size of SOPA per pupil and overall PA, LPA, MVPA or time spent sedentary. Among the 15-year-olds, we found the size of SOPA per pupil to be positively associated with LPA and negatively associated with MVPA (Table 5). These associations translates to an increase in LPA of 0.9 min/d and a decrease in MVPA of 0.4 min/d if the size of SOPA increased by 10m² per pupil.

4 Discussion

The results from the present study suggest a weak association between outdoor PPF availability, time spent sedentary and time spent in LPA among six-year-olds. An increase in the number of PPF from 0.1 to 0.2 per pupil, which equates to a doubling of PPF in an average Norwegian primary school, was associated with 3.1% less sedentary time and 2.5% more time spent in LPA during school hours. However, since the influence of PPF on sedentary time and PA is mainly restricted to recess, these weak, although statistically significant, associations may not be negligible. Primary and lower secondary schools in

Norway provide approximately 60 min/day of recess (10-15 minutes of morning recess, 30-40 minutes of lunch recess and 10-15 minutes of afternoon recess). Although speculative, if the observed associations were in fact restricted to recess, they would translate to ~6.3% less sedentary time and ~3.6% more LPA. However, we did not observe any associations between sedentary time and PA with PPF in nine-year-olds. This may be explained by differences between age groups in time spent outdoors during school hours. In the Norwegian school system part of the taught classes in the first grade are outdoors classes, possibly contributing differences in observed associations.

Studies investigating the isolated association between PPF provision and objectively measured sedentary time in children are limited. Results from Ridgers et al. (2010) support our finding that PPF provision is negatively associated with sedentary time during school hours (Ridgers et al. 2010a). In their study, children without access to fixed equipment during recess engaged in 8.2% more sedentary activity than children provided with fixed equipment.

We did not find PPF provision to be associated with overall PA or MVPA in children. This is supported by two studies conducted in Australia, where no association between equipment availability (other than balls) and MVPA (Zask et al. 2001) or energy expenditure (Harten et al. 2008) was observed. In contrast, Ridgers et al. (2010) suggested a positive association between PPF provision and MVPA (Ridgers et al. 2010a). The latter is also supported by three other cross-sectional studies that used accelerometers to assess PA (Nielsen et al. 2010;Nielsen et al. 2012;Taylor et al. 2011). However, the strength of the associations reported in these studies varied considerably (Nielsen et al. 2010;Taylor et al. 2011). Consequently, studies differ in their conclusions with regard to the actual importance of PPF provision for children's PA during school hours.

Contradictory results may reflect actual differences in the everyday life of children in different study populations, e.g. due to different school policies regarding PA. However, in three of the studies reporting an association, only PPF that had previously been observed to be used for play and/or sports activities during break-time were counted. In addition, PPF that facilitated active play for several small groups of children at the same time were counted as more than one item (Nielsen et al. 2010;Nielsen et al. 2012;Taylor et al. 2011). This might have given a more detailed and realistic picture on PPF accessibility than in the present study, where we counted all individual play structures as one item. It is therefore possible that the associations between PPF provision, sedentary time and LPA found in the present study are

underestimated. One could argue that a doubling of the sheer number of PPF (from ~22 to ~44 in an average Norwegian primary school) is neither realistic nor practical when we consider the relatively modest associations observed in the present study. However, investing in PPF that promote PA for many children at the same time might both be realistic and practical. Further research is therefore needed to identify what sort of PPF increases PA-levels the most.

Although a few experimental studies have investigated the isolated effect of altering PPF availability (Ickes et al. 2013; Parrish et al. 2013; van Sluijs et al. 2007; Ridgers et al. 2010b), we are only aware of one such study with a long term follow-up. In this study, Ridgers et al. (2007) investigated the effect of redesigning the playground environment in elementary schools on MVPA and vigorous PA (VPA) during recess (Ridgers et al. 2007). Results demonstrated significant intervention effects after 6 weeks and 6 months (Ridgers et al. 2007), but at 12 months, the only significant intervention effect that remained was higher VPA during lunch recess (Ridgers et al. 2010b). This might indicate a novelty effect of the intervention and, furthermore, that regular changes in the outdoor playing environment might be necessary in future interventions to increase PA in the long term.

Using questionnaires to assess PA, Haug et al. (2010) investigated the association between characteristics of the outdoor school environment and PA in a nationally representative sample of Norwegian 13-15 year olds. They found that adolescents with access to the maximum number of play facilities had almost three times higher odds of being physically active during recess than adolescents attending schools not providing play facilities. Although comparability is limited because of the different methods used to assess both PA and play facilities, this is in contrast to our findings. We are not aware of studies that have investigated the association between objectively measured PA or sedentary time and PPF availability in adolescents.

Few cross-sectional studies have investigated the association between the size of SOPA and objectively measured PA in children, and the results are equivocal. Two studies conducted by Nielsen et al. (Nielsen et al. 2010; Nielsen et al. 2012) support our findings of no association. On the other hand, five studies report positive associations between the size of SOPA and MVPA during recess (Sallis et al. 2001; Harten et al. 2008; Ridgers et al. 2010a; D'Haese et al. 2013; Escalante et al. 2012). Furthermore, interventional studies indicate a positive effect of increasing the size of SOPA per pupil on PA during recess (Loucaides et al. 2009; D'Haese et

al. 2013;Harten et al. 2008). Although small sample sizes and short follow-up limit the generalizability of these studies, they contrast with our findings.

One possible reason for the differing results might be due to the actual size of SOPA. In the present study, and in the studies by Nielsen et al., the size of SOPA per pupil was much larger than in the other studies. In PANCS2, only four of the 60 participating primary schools provided $<15\text{m}^2$ of outdoor play space per pupil, while none of the 18 schools in the study by Nielsen et al. (Nielsen et al. 2012) conducted on Danish children provided $<77\text{m}^2$ per pupil. In comparison, none of the 11 participating schools in the two studies by Ridgers et al. (Ridgers et al. 2010a) and D'Haese et al. (D'Haese et al. 2013) provided $>16.9\text{m}^2$ per pupil. It is therefore likely that smaller outdoor play areas might inhibit the PA level of children, but that most Norwegian schools provide children with sufficient outdoor play space to be physically active. Explorative analyses of the third of schools ($n=16$) providing the least play space per pupil in the present study ($4\text{-}40\text{m}^2$) did however indicate a positive association between play space and MVPA among nine-year-olds (data not shown). Further research on a larger sample of schools with smaller outdoor play areas (e.g. $<40\text{m}^2$) could be useful for developing general recommendations on the minimum outdoor play space per pupil that should be provided with regard to PA.

Discrepancy in results between studies could also be due to differences in methods used to measure PA. In our study, and in the studies by Nielsen et al. (Nielsen et al. 2010;Nielsen et al. 2012), PA was measured objectively and continuously for several days. The other studies measured PA levels during recess only (D'Haese et al. 2013;Escalante et al. 2012;Harten et al. 2008;Ridgers et al. 2010a;Sallis et al. 2001). Isolating the PA measurements to recess could be more sensitive and therefore enable the detection of smaller differences in PA. However, if children are aware of the PA-monitoring, either as consequence of being observed (Ridgers et al. 2010a;Sallis et al. 2001) or being equipped with a PA monitor just before recess (D'Haese et al. 2013;Escalante et al. 2012;Harten et al. 2008), they might alter their normal recess behavior (Dossegger et al. 2014). Thus, a Hawthorne effect can't be excluded (McCambridge et al. 2014).

Although the size of SOPA was negatively associated with MVPA and positively associated with LPA among 15-year-olds, these associations were weak and likely not clinically meaningful. When we also take in to consideration that only five of the 44 included lower

secondary schools provided $\leq 20\text{m}^2$ of outdoor play area per pupil, we could expect that the size of SOPA does not seem to be a limiting factor for PA among the 15-year-olds.

4.1 Study limitations and strengths

A major strength of the present study is the large, nationally representative sample of children and adolescents. Another strength is the objective and continuous measure of PA over multiple days. Because of known difficulties with accurately recalling details about PA, especially among children (Sallis and Saelens 2000), objective measurement with accelerometers is considered the best option in large scale studies (Westerterp 2009). Lastly, the high number of participants from a large number of schools allowed us to include several covariates in the statistical models.

This study also has some important limitations. First, this is a cross-sectional study, and we can therefore not make inferences about cause and effect. Second, because several schools did not provide us with class schedules, we were not able to use isolated recess PA/sedentary time in the analyses, or to control for PE. However, we used analyses that partly account for nesting effects within schools, and we do not have any indications that recess or PE durations were not randomly distributed between schools. Third, we did not consider the use and quality of the PPF. Therefore, it is unknown how many pupils actually used the different play facilities, or how much PA they could potentially generate. Fourth, because we used vertical accelerations of the hip to assess PA, it is likely that the intensity (energy expenditure) of PA involving substantial upper-body movements, such as climbing, was underestimated (Lee and Shiroma 2013). Lastly, we do acknowledge that landscape features, such as areas with a naturalistic feel and areas with different surfaces, may influence the PA level of children (Anthamatten et al. 2014; Fjørtoft 2004). Because of the risk of overfitting the regression models, we chose not to adjust for this. Additional explorative analyses using soft surface area, asphalt area or treetop-covered area as the dependent variable in the model did not change the observed results.

5 Conclusions

Our results indicate that increasing the sheer number of PPF in SOPA may be beneficial to reduce sedentary time and increase time spent in LPA among six-year-olds, but not among nine- and 15-year-olds. In order to recommend cost-effective changes to SOPA, there is a need to identify what types of PPF that increase PA-levels the most, especially in adolescents.

The size of SOPA did not seem to be a limiting factor for PA in the present study. This may be explained by the large outdoor areas generally observed in Norwegian schools.

6 Conflicts of interest statement

The authors declare that there are no conflicts of interest.

7 Acknowledgements

Financial support was received from the Norwegian Directorate of Health and the Norwegian School of Sport Sciences. The authors thank all the test personnel for their work during the data collection and Professor Ingar M. Holme for statistical guidance.

8 References

Andersen, L.B., Harro, M., Sardinha, L.B., Froberg, K., Ekelund, U., Brage, S., and Anderssen, S.A., 2006 Physical activity and clustered cardiovascular risk in children: a cross-sectional study (The European Youth Heart Study). *Lancet* 368, 299-304.

Anthamatten, P., Brink, L., Kingston, B., Kutchman, E., Lampe, S., and Nigg, C., 2014 An Assessment of Schoolyard Features and Behavior Patterns in Children's Utilization and Physical Activity. *Journal of Physical Activity & Health* 11, 564-573.

Blaes, A., Ridgers, N.D., Aucouturier, J., Van, P.E., Berthoin, S., and Baquet, G., 2013 Effects of a playground marking intervention on school recess physical activity in French children. *Prev.Med* 57, 580-584.

D'Haese, S., Van, D.D., De, B., I, and Cardon, G., 2013 Effectiveness and feasibility of lowering playground density during recess to promote physical activity and decrease sedentary time at primary school. *Bmc Public Health* 13, 1154.

Dossegger, A., Ruch, N., Jimmy, G., Braun-Fahrlander, C., Mader, U., Hanggi, J., Hofmann, H., Puder, J.J., Kriemler, S., and Bringolf-Isler, B., 2014 Reactivity to accelerometer measurement of children and adolescents. *Med Sci Sports Exerc* 46, 1140-1146.

Ekelund, U., Sjostrom, M., Yngve, A., Poortvliet, E., Nilsson, A., Froberg, K., Wedderkopp, N., and Westerterp, K., 2001 Physical activity assessed by activity monitor and doubly labeled water in children. *Medicine and Science in Sports and Exercise* 33, 275-281.

Escalante, Y., Backx K, Saavadra JM, Garcia-Hermoso A, and Dominguez AM, 2012 Play Area and Physical Activity in Recess in Primary School. *Kinesiology* 44, 123-129.

finn.no, 2011 <http://kart.finn.no/>.

Fjørtoft, I., 2004 Landscape as Playscape: The Effects of Natural Environments on children's Play and Motor Development. *Children, Youth and Environments* 14(2), 2004.

Fjørtoft, I., Kristoffersen, B.r., and Sageie, J., 2009 Children in schoolyards: Tracking movement patterns and physical activity in schoolyards using global positioning system and heart rate monitoring. *Landscape and Urban Planning* 93, 210-217.

Gray, C., Gibbons, R., Larouche, R., Sandseter, E.B., Bienenstock, A., Brussoni, M., Chabot, G., Herrington, S., Janssen, I., Pickett, W., Power, M., Stanger, N., Sampson, M., and Tremblay, M.S., 2015 What Is the Relationship between Outdoor Time and Physical Activity, Sedentary Behaviour, and Physical Fitness in Children? A Systematic Review. *Int J Environ.Res Public Health* 12, 6455-6474.

Harten, N., Olds, T., and Dollman, J., 2008 The effects of gender, motor skills and play area on the free play activities of 8-11 year old school children. *Health Place* 14, 386-393.

Haug, E., Torsheim, T., Sallis, J.F., and Samdal, O., 2010 The characteristics of the outdoor school environment associated with physical activity. *Health Education Research* 25, 248-256.

Healy, G.N. and Owen, N., 2010 Sedentary behaviour and biomarkers of cardiometabolic health risk in adolescents: an emerging scientific and public health issue. *Rev.Esp.Cardiol.* 63, 261-264.

Ickes, M.J., Erwin, H., and Beighle, A., 2013 Systematic review of recess interventions to increase physical activity. *J Phys Act Health* 10, 910-926.

Larouche, R., Saunders, T.J., Faulkner, G., Colley, R., and Tremblay, M., 2014 Associations between active school transport and physical activity, body composition, and cardiovascular fitness: a systematic review of 68 studies. *J Phys Act Health* 11, 206-227.

Lee, I.M. and Shiroma, E.J., 2013 Using accelerometers to measure physical activity in large-scale epidemiological studies: issues and challenges. *British Journal of Sports Medicine*.

Lonsdale, C., Rosenkranz, R.R., Peralta, L.R., Bennie, A., Fahey, P., and Lubans, D.R., 2013 A systematic review and meta-analysis of interventions designed to increase moderate-to-vigorous physical activity in school physical education lessons. *Preventive Medicine* 56, 152-161.

Loucaides, C.A., Jago, R., and Charalambous, I., 2009 Promoting physical activity during school break times: piloting a simple, low cost intervention. *Prev.Med* 48, 332-334.

McCambridge, J., Witton, J., and Elbourne, D.R., 2014 Systematic review of the Hawthorne effect: New concepts are needed to study research participation effects(). *Journal of Clinical Epidemiology* 67, 267-277.

McClain, J.J., Abraham, T.L., Brusseau, T.A., Jr., and Tudor-Locke, C., 2008 Epoch length and accelerometer outputs in children: comparison to direct observation. *Med Sci Sports Exerc* 40, 2080-2087.

Nielsen, G., Bugge, A., Hermansen, B., Svensson, J., and Andersen, L.B., 2012 School Playground Facilities as a Determinant of Children's Daily Activity: A Cross-Sectional Study of Danish Primary School Children. *Journal of Physical Activity & Health* 9, 104-114.

Nielsen, G., Taylor, R., Williams, S., and Mann, J., 2010 Permanent play facilities in school playgrounds as a determinant of children's activity. *J Phys Act Health* 7, 490-496.

Nilsen, A., 2014 Available outdoor space and competing needs in public kindergartens in Oslo . *FORMakademisk* 7, 1-13.

Pagels, P., Raustorp, A., De Leon, A.P., Martensson, F., Kylin, M., and Boldemann, C., 2014 A repeated measurement study investigating the impact of school outdoor environment upon physical activity across ages and seasons in Swedish second, fifth and eighth graders. *Bmc Public Health* 14, 803.

Parrish, A.M., Okely, A.D., Stanley, R.M., and Ridgers, N.D., 2013 The Effect of School Recess Interventions on Physical Activity: A Systematic Review. *Sports Medicine* 43, 287-299.

Ridgers, N.D., Fairclough, S.J., and Stratton, G., 2010a Variables associated with children's physical activity levels during recess: the A-CLASS project. *The International Journal of Behavioral Nutrition and Physical Activity* 7, 74.

Ridgers, N.D., Stratton, G., and Fairclough, S.J., 2006 Physical activity levels of children during school playtime. *Sports Med* 36, 359-371.

Ridgers, N.D., Fairclough, S.J., and Stratton, G., 2010b Twelve-Month Effects of a Playground Intervention on Children's Morning and Lunchtime Recess Physical Activity Levels. *Journal of Physical Activity & Health* 7, 167-175.

Ridgers, N.D., Stratton, G., Fairclough, S.J., and Twisk, J.W.R., 2007 Long-term effects of a playground markings and physical structures on children's recess physical activity levels. *Preventive Medicine* 44, 393-397.

Sallis, J.F., Conway, T.L., Prochaska, J.J., McKenzie, T.L., Marshall, S.J., and Brown, M., 2001 The association of school environments with youth physical activity. *Am.J.Public Health* 91, 618-620.

Sallis, J.F. and Saelens, B.E., 2000 Assessment of physical activity by self-report: status, limitations, and future directions. *Res Q.Exerc.Sport* 71, S1-14.

Sleap, M. and Warburton, P., 1996 Physical activity levels of 5-11-year-old children in England: cumulative evidence from three direct observation studies. *Int J Sports Med* 17, 248-253.

Taylor, R.W., Farmer, V.L., Cameron, S.L., Meredith-Jones, K., Williams, S.M., and Mann, J.I., 2011 School playgrounds and physical activity policies as predictors of school and home time activity. *The International Journal of Behavioral Nutrition and Physical Activity* 8, 38.

van Sluijs, E.M.F., McMinn, A.M., and Griffin, S.J., 2007 Effectiveness of interventions to promote physical activity in children and adolescents: systematic review of controlled trials. *BMJ* 335, 703.

Westerterp, K.R., 2009 Assessment of physical activity: a critical appraisal. *Eur.J.Appl.Physiol* 105, 823-828.

Zask, A., van, B.E., Barnett, L., Brooks, L.O., and Dietrich, U.C., 2001 Active school playgrounds-myth or reality? Results of the "move it groove it" project. *Prev.Med* 33, 402-408.

Table 1: Characteristics of the study sample in the Physical Activity among Norwegian Children Study (2011) by age and sex (n=3040).

	6-year-olds		9-year-olds		15-year-olds	
	Girls	Boys	Girls	Boys	Girls	Boys
n	489	479	663	625	393	391
Age (yrs.) ^a	6.6 (0.4)	6.6 (0.4)	9.6 (0.4)	9.6 (0.4)	15.1 (0.6)	15.1 (0.6)
Height (cm) ^a	121.0 (5.5) ^c	122.2 (5.8)	138.0 (6.5)	138.7 (6.8)	164.7 (6.4) ^c	172.9 (8.0)
Weight (kg) ^{ab}	23.8 (4.2)	24.0 (3.8)	33.7 (6.8)	33.9 (6.9)	57.1 (9.4) ^c	62.0 (12.0)
BMI (kg/m ²) ^a	16.2 (1.9)	16.0 (1.6)	17.6 (2.7)	17.5 (2.7)	21.1 (3.1)	20.7 (3.3)
Parents' educ. level (%)						
Low	7.5	6.8	10.8	9.0	10.9	12.7
Middle low	30.8	31.7	31.1	36.3	33.1	35.4
Middle high	45.8	45.7	43.0	40.7	39.1	37.7
High	15.8	15.9	15.0	14.0	16.9	14.3

BMI = body mass index; educ. = education.

^a Mean (standard deviation).

^b The weight was corrected by -0.3 kg for all participants to account for clothes.

^c Significantly different from boys within age group (all p-values ≤ 0.001).

Table 2: Mean (SD) physical activity and minutes of time spent sedentary among Norwegian children and adolescents during school hours^a

	6-year-olds		9-year-olds		15-year-olds	
	Girls	Boys	Girls	Boys	Girls	Boys
n	489	479	663	625	393	391
Overall PA (CPM) ^b	765±211 ^{c, e}	845±227 ^c	607±183 ^{d, e}	750±206 ^d	358±138 ^e	475±173
Sedentary (min/d) ^b	127±18 ^{c, e}	118±20 ^c	150±23 ^{d, e}	137±23 ^d	237±36 ^e	214±29
LPA (min/d) ^b	85±14 ^{c, e}	88±13 ^c	71±14 ^{d, e}	76±15 ^d	56±16 ^e	69±18
MVPA (min/d) ^b	31±9 ^{c, e}	37±11 ^c	26±9 ^{d, e}	34±11 ^d	20±10 ^e	26±11

PA=physical activity; CPM=counts per minute; LPA=light physical activity; MVPA=moderate-to-vigorous physical activity.

^a 9 AM to 1 PM for six- and nine-year-olds, 9 AM to 2PM for 15-year-olds

^b Mean ± standard deviation.

^c Significantly different from nine- and 15-year-olds (p<0.001).

^d Significantly different from 15-year-olds (p<0.001).

^e Significantly different from boys in the same age group (p≤0.045).

Table 3: Permanent play facility provision and the size school's outdoor play area in schools participating in the Physical Activity among Norwegian Children Study in 2011 (n=99)

	6-year-olds	9-year-olds	15-year-olds
Permanent play facilities ^a	22.2±7.5	21.7±7.8	10.9±7.1 ^b
Per pupil ^a	0.095±0.055	0.093±0.058	0.037±0.033 ^b
SOPA (m ²) ^a	15249±7958	15128±8018	14428±7279
Per pupil (m ²) ^a	65.6±45.2	62.9±43.0	49.9±35.7 ^b

SOPA = school's outdoor play area

^a Mean ± standard deviation

^b Significantly different from 6- and 9-year-olds (p<0.001)

Table 4: Associations between permanent play facility provision, physical activity and sedentary time among Norwegian children and adolescents in 2011 (n=2588)^a

	Age	n pupils (n schools)	B ^{b, c}	95% CI
Overall PA (CPM)	6	837 (55)	19.91	-26.09, 65.90
Sedentary (minutes)	6	837 (55)	-3.78*	-7.28, -0.28
LPA (minutes)	6	837 (55)	2.16**	0.53, 3.79
MVPA (minutes)	6	837 (55)	1.67	-0.55, 3.89
Overall PA (CPM)	9	1126 (55)	7.35	-36.65, 51.35
Sedentary (minutes)	9	1126 (55)	-1.92	-6.23, 2.39
LPA (minutes)	9	1126 (55)	1.93	-0.36, 4.22
MVPA (minutes)	9	1126 (55)	0.04	-2.52, 2.59
Overall PA (CPM)	15	625 (36)	-25.08	-94.74, 44.57
Sedentary (minutes)	15	625 (36)	0.17	-7.29, 7.64
LPA (minutes)	15	625 (55)	0.77	-4.08, 5.61
MVPA (minutes)	15	625 (36)	-0.90	-5.95, 4.14

PA=physical activity; CPM=counts per minute; LPA=light physical activity; MVPA=moderate-to-vigorous physical activity.

^a Data on one or more of the covariates in the statistical models were missing for 452 of the 3040 participants that met the inclusion criteria, therefore the results from the analyses are based on a total of 2588 participants.

^b Beta values represent daily change associated with increasing the number of permanent play facilities per pupil by 0.1.

^c Analyses adjusted for: accelerometer wear time (except analyses of CPM); measurement month; socioeconomic status; the dummy variables "access to areas outside school property during recess", "sectioning of the play area during recess", "recess at different time points for different classes" and "allowed to spend recess indoors".

* p=0.034

** p=0.009

Table 5: Associations between the size of school's outdoor play area, physical activity and sedentary time among Norwegian children and adolescents in 2011 (n=2588)^a

	Age	n pupils (n schools)	B ^{b, c}	95% CI
Overall PA (cpm)	6	837 (55)	3.43	-3.80, 10.67
Sedentary (minutes)	6	837 (55)	-0.01	-0.54, 0.52
LPA (minutes)	6	837 (55)	-0.02	-0.38, 0.33
MVPA (minutes)	6	837 (55)	0.03	-0.24, 0.29
Overall PA (minutes)	9	1126 (55)	2.40	-4.07, 8.88
Sedentary (minutes)	9	1126 (55)	0.08	-0.75, 0.59
LPA (minutes)	9	1126 (55)	0.06	-0.36, 0.47
MVPA (minutes)	9	1126 (55)	0.03	-0.32, 0.39
Overall PA (cpm)	15	625 (36)	-2.0	-3.39, 3.40
Sedentary (minutes)	15	625 (36)	-0.47	-1.26, 0.32
LPA (minutes)	15	625 (36)	0.86*	0.21, 1.50
MVPA (minutes)	15	625 (36)	-0.41**	-0.77, -0.05

PA=physical activity; CPM=counts per minute; LPA=light physical activity; MVPA=moderate-to-vigorous physical activity.

^a Data on one or more of the covariates in the statistical models were missing for 452 of the 3040 participants that met the inclusion criteria, therefore the results from the analyses are based on a total of 2588 participants.

^b Beta values represent daily change associated with increasing outdoor play area size by 10m².

^c Analysis adjusted for: accelerometer wear time (except analyses of CPM); measurement month; socioeconomic status; number of permanent play facilities; the dummy variables "access to areas outside school property during recess", "sectioning of the play area during recess", "recess at different time points for different classes" and "allowed to spend recess indoors".

* p=0.009

** p=0.027.