



## Original research

# Towards more valid simulations of slopestyle and big air jumps: Aerodynamics during in-run and flight phase

Fabian Wolfspurger<sup>a,\*</sup>, Frédéric Meyer<sup>b,2</sup>, Matthias Gilgien<sup>c,d,3</sup>

<sup>a</sup> Winter Sports and Climate Group, WSL-Institute of Snow and Avalanche Research SLF, Switzerland

<sup>b</sup> Department of Sport Science, University of Lausanne, Switzerland

<sup>c</sup> Department of Physical Performance, Norwegian School of Sport Sciences, Norway

<sup>d</sup> Center for Alpine Sports Biomechanics, St. Moritz Health and Innovation Foundation, Switzerland

## ARTICLE INFO

## Article history:

Received 30 November 2019

Received in revised form 19 March 2021

Accepted 11 May 2021

Available online 18 May 2021

## Keywords:

Accident prevention

Skiing

Snowboarding

Aerodynamics

Apparel

Body posture

## ABSTRACT

**Objectives:** This study aimed to investigate air drag and lift during the in-run and flight phase of ski and snowboard slopestyle and big air, to allow more valid modeling of jumps and hence reduce injury risk.

**Design:** We present an experimental, multiple single athlete study based on wind tunnel measurements of 4 skiers and 3 snowboarders.

**Methods:** Measurements were carried out in a closed loop wind tunnel, measuring airflow speed and 3D forces acting on the athletes. Athletes performed trials in typical postures at 35, 60 and 85 km/h wearing slim-, regular- and wide fit apparel. Drag and lift area ( $c_D A$ ;  $c_L A$ ) were calculated and analyzed using linear and multiple regression to describe their dependencies on posture, apparel and speed.

**Results:**  $c_D A$  values were higher than earlier assumed and ranged from 0.3 to 0.95 m<sup>2</sup> for skiers and from 0.35 to 0.55 m<sup>2</sup> for snowboarders, primarily dominated by posture, and followed by apparel.  $c_L A$  ranged from −0.1 to 0.45 m<sup>2</sup> for skiers and from 0.04 to 0.17 m<sup>2</sup> for snowboarders. To facilitate more valid jump modeling posture- and apparel-dependent formulations for air drag coefficients were provided and the consequences of sport specific differences on modeling were highlighted.

**Conclusions:** Applying the air drag coefficients and relationships determined in this study will help to improve validity of jump modeling in big air and slopestyle. The variability in aerodynamic forces in slopestyle and big air is caused by differences between sports, posture and apparel.

© 2021 The Authors. Published by Elsevier Ltd on behalf of Sports Medicine Australia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Practical implications

- The presented data and models will help to improve jump design and thus prevent injuries.
- Skiers' capacity to compensate for low speed with changes in body posture and drag area is much larger than for snowboarders.
- Course builders should take care of the snowboarders' limited ability to compensate for low in-run speed in jump design.
- Athletes and coaches should be aware, that body posture and apparel can strongly influence take-off speeds.

## 1. Introduction

Ski and snowboard (SNB) slopestyle and big air (BA) are young but rapidly developing winter sports with high injury rates.<sup>1–5</sup> Jumps are considered a key injury risk factor.<sup>6–8</sup> Jump modeling was suggested as a method to help course builders to validate and improve jump design and enhance jump safety.<sup>9–18</sup> The principles of jump kinematics and kinetics are well established. Angle and speed of the center of mass trajectory at take-off, air drag and lift during the flight phase, and the shape of the landing are the initial determinants of jump length, height and impact. Take-off speed is regulated by the shape of the in-run, which determines the energy to accelerate the athlete by gravity, and the braking forces of snow friction and air drag.<sup>19</sup> Finally, athletes have a last chance to modify their trajectory by absorbing or pushing on his leg ('pop'), just before take-off.<sup>20</sup> In competitive alpine skiing, the relative contributions to energy dissipation from air drag and snow friction have been determined. The distribution is 23% for air drag and 77% for snow friction in giant slalom, and reaches 51% for air drag and 49% for snow friction in downhill.<sup>21</sup> Such information is entirely lacking for ski and snowboard slopestyle and BA, resulting in

\* Corresponding author.

E-mail addresses: [wolfspurger@slf.ch](mailto:wolfspurger@slf.ch) (F. Wolfspurger), [matthias.gilgien@nih.no](mailto:matthias.gilgien@nih.no) (M. Gilgien).

<sup>1</sup> WSL-Institut für Schnee- und Lawinenforschung SLF, Switzerland.

<sup>2</sup> University of Lausanne, Quartier UNIL-Centre, Switzerland.


<sup>3</sup> Norwegian School of Sport Sciences, Department of Physical Performance, Norway.

considerable uncertainties when modeling the take-off speed. Hence, to date, park designers have to rely on their experience when building in-runs for jumps. To improve the validity of jump modeling, realistic values for snow friction and air drag need to be established. This study aimed to establish realistic air drag and lift force values for BA and slopestyle ski and snowboard, to improve the validity of computer simulations of jumps in these sports. Air drag and lift are dependent on air properties, ambient wind, athlete speed, and the athlete's body posture and apparel.<sup>22–27</sup> Apparel and body posture are the factors athletes can influence to regulate take-off speed during the in-run. To date, air drag values have to be estimated from values available for other winter sports such as alpine skiing, where clothing and body posture and hence air drag are very different.<sup>28,29</sup> Therefore, the aim of this study was to measure typical mean values and the range of air drag and lift for elite level athletes in ski and snowboard, slopestyle and BA, as a function of the athletes' posture, apparel, speed and anthropometrics, using wind tunnel testing for postures typical for both the in-run and the airborne phase.

## 2. Methods

*Wind tunnel measurements* were carried out in an atmospheric closed loop wind tunnel with a closed test section of  $7.5 \times 11$  m, at RUAG Aerospace Center at Emmen, Switzerland.<sup>30</sup> Four male athletes (37, 34, 32, 37 years; 90.2, 74.2, 90.3, 71.6 kg; 182, 184, 191, 175 cm) participated in the study and were labeled as athletes A to D. Their anthropometry was quantified by their body surface ( $A_b$ ), calculated from body height and weight according to Boyd et al.<sup>31</sup>

To investigate drag and lift forces during ski and snowboard *slopestyle in-runs*, four *skiers* (A to D) and three *snowboarders* (A, C and D) performed two trials each with three typical in-run postures at 35, 60 and 85 km/h, wearing regular fit snow sports apparel. Three *typical postures* (Fig. 1) were chosen to capture the whole range of realistic in-run postures in ski and snowboard. For skiing, the postures were as follows. 1) In the low posture the athlete had fully flexed knees with the arms wrapped around the thighs. 2) In the mid posture athletes held their knees with straight arms and an extended upper body. 3) In



**Fig. 1.** a–b) Skier B in low (left), mid and extended (right) in-run posture wearing regular (a) and wide-fit (b) apparel. c–d) Snowboarder D in mid (left), extended and extended rotated (right) posture wearing slim (c) and wide-fit (d) apparel. e) Airborne postures measuring drag area at vertical inflow hitting the snowboard/ski from below. f) Left: Snowboarder D wearing regular apparel performing a mute grab with inflow direction from behind (rotated 90°). Right: Skier B performing a mute grab with inflow direction from behind (rotated 180°).

the high posture athletes stood upright with slight flexion of the knees and the hips, with arms held straight along the upper body. For snowboarding, we tested 1) a typical riding posture, characterized by considerable knee flexion, slight elbow flexion and upper arm abduction; 2) an upright posture with the trunk held parallel to the transverse axis; and 3) an upright posture with the trunk rotated 50° to 80° in the riding direction. For both upright postures, the arms were held crossed on the back. The postures were further quantified by the extension height  $h_{ext}$ , defined as the vertical distance between the ski boot sole and the antenna mounted on the athletes' helmet (taken from side view). We found that  $h_{ext}$  was an inadequate descriptor of typical postures in snowboarding. We therefore introduced an ordinal scaled posture number ( $n_{pos} = 1; 2; 3$  for mid, extended and extended rotated).

To investigate the influence of *different apparel* on the aerodynamic forces, skiers B and C and snowboarder D performed additional trials wearing slim-fit and wide-fit apparel. To provide apparel fitting categories that had practical relevance and were easy to apply, we introduced an ordinal scaled apparel fit number ( $n_{fit}$ ) as follows: extra slim (1, race suit); slim (2, regular jeans and sweatshirt); regular (3, regular ski/SNB apparel); wide (4, oversized ski/SNB apparel); and extra wide (5, double oversized ski/SNB apparel). The difference between extra-large sized apparel and the athlete's regular size was used to distinguish between wide (skier B, D) and extra-wide fits (snowboarder D). The regular sizes were deduced from the athletes' chest and waist girth (A to D: 1.05/0.95 m, 0.95/0.815 m, 1.04/0.98 m, 0.955/0.815 m), based on the European norm 13,402–2:2002 for "Size designation of clothes" (A and C size large; B and D size medium).

Additional trials at 60 km/h were performed to investigate lift and drag in an *airborne-like position*, holding the skis/snowboard under inflow directions between 0° and 180° (Fig. 1f). To quantify drag forces in the late flight phase, when the vertical airflow component increases, another test trial was made, rotating the athletes' vertical axis into the horizontal direction, so that the airflow hit the skier/snowboarder from below (Fig. 1e).

For each trial drag force  $F_D$ , lift force  $F_L$ , dynamic pressure  $q$ , airflow speed  $v$  and air density  $\rho$ , were measured at 0.5 Hz during 20 s. The sensor accuracy was  $\pm 0.01$  N for the forces and  $\pm 0.01$  Pa for the pressure. Drag area ( $c_D A$ ) and lift area ( $c_L A$ ) were calculated.

$$c_L A = F_L / q \quad \text{with } q = 0.5 \cdot \rho \cdot v^2 \tag{1}$$

$$c_D A = F_D / q \tag{2}$$

The dependencies of  $c_D A$  and  $c_L A$  on posture and speed were analyzed for the regular apparel trials for skiers and snowboarders, using least squares first or second order regression. One-parameter models of the dominant variable were provided as a first approximation of the aerodynamic coefficients. Mean ( $\pm \sigma$ )  $c_D A$  and  $c_L A$  values were calculated for each posture. The means of the different postures were then compared by their relative differences to quantify the impact of posture on  $c_D A$  within the group of snowboarders and skiers. Beyond that, the dependencies of  $c_D A$  on the full set of explanatory variables (posture, apparel and speed) were analyzed separately for those athletes who conducted both posture and apparel tests, using multiple least squares linear regression. The ordered categorical variables  $n_{fit}$  and  $n_{pos}$  were implemented to the regression models using two indicator variables each ( $X_{p1}, X_{p2}; X_{f1}, X_{f2}$ ) to represent the three tested posture and fit categories.<sup>32,33</sup> Mean ( $\pm \sigma$ )  $c_D A$  values were calculated for each posture and athlete, and were compared using relative differences. For all regression models, adjusted coefficients of determination ( $R_{adj}^2$ ) and slopes of regression ( $m$ ) were stated (partly in supplementary materials) if significant ( $p \leq 0.001$ ) to quantify the strength and relevance

of those relationships. The uncertainties of the regression coefficients, as well as model outputs, were given by their 95% confidence intervals ( $\pm CI/2$ ), partly as supplementary material (Table A5).

### 3. Results

For the skiers, change in *body posture* had the largest impact on  $c_D A$ . Posture-induced  $c_D A$  changes deviated from the mid posture by  $-41 \pm 5\%$  for the tuck posture to  $+36 \pm 5\%$  for the extended posture (Fig. 2; Table A1). For snowboarders, changes in posture influenced  $c_D A$  less but were still a main influence factor:  $c_D A$  deviated from the extended posture by  $-6 \pm 2\%$  for the mid posture to  $+15 \pm 8\%$  for the extended rotated posture (Fig. 2b; Table A1). We did not find a significant *influence of speed* on  $c_D A$  for ski and snowboard on the group level. For skiers and snowboarders wearing the most commonly used type of apparel (regular fit) Eqs. (3) and (4) were set up to calculate drag area as a function of body posture ( $h_{ext}$  for ski and  $n_{pos}$  for snowboard):

$$c_D A_{Ski} = -0.137 + 0.491 \cdot h_{ext} \text{ for } 0.89 \text{ m} \leq h_{ext} \leq 2.00 \text{ m}; (R_{adj}^2 = 0.91) \tag{3}$$

$$c_D A_{SNB} = 0.0466 - 0.027 \cdot X_{p1} + 0.070 \cdot X_{p2} \tag{4}$$

$$X_{p1} = \begin{cases} 1 & \text{if } n_{pos} = 1 \\ 0 & \text{if } n_{pos} > 1 \end{cases} \quad X_{p2} = \begin{cases} 1 & \text{if } n_{pos} = 3 \\ 0 & \text{if } n_{pos} < 3 \end{cases} \quad \text{for } n_{pos} = [1; 2; 3]; (R_{adj}^2 = 0.53)$$


During the in-run, athletes commonly use the mid posture. Hence, to model the typical development of an athlete's in-run speed, Eqs. (3) and (4) propose drag areas of  $0.634 \pm 0.035 \text{ m}^2$  for skiers and  $0.439 \pm 0.014 \text{ m}^2$  for snowboarders, using an average mid posture as equation input ( $h_{ext\_mid\_mean} = 1.571 \pm 0.072$ ;  $n_{pos} = 1$ ).

*Aerodynamic lift* was measured to provide also data to model the snow friction force during the in-run revealing mean  $c_L A$ s of  $0.319 \pm 0.038 \text{ m}^2$  for the skiers and  $0.081 \pm 0.008 \text{ m}^2$  for the snowboarders, both in mid posture wearing regular clothes (Table A2). Posture influenced the  $c_L A$  strongly for the skiers ( $R_{adj}^2 = 0.80$ ), but only slightly for snowboarders ( $R_{adj}^2 = 0.37$ ) (see regression models and scatter plots in Table A5 and Fig. A1). For skiers, the  $c_L A$  peaked in mid posture when the airflow hit the inclined upper body.

*Individual deviations* from the generalized posture-dependent drag and lift models can be caused by *anthropometric differences*, deviations caused by *apparel of different fit* or by *individual speed dependencies* of the athletes'  $c_D A$ s (Fig. 3). The contribution of *apparel* to  $c_D A$  was analyzed for two skiers and one snowboarder. In all postures, apparel affected the  $c_D A$  of the tested skiers, but to a lesser extent than posture did (Table A3). The largest effects of apparel, averaged over all speeds, occurred in the mid posture, where  $c_D A$  changed from slim to wide apparel by  $-9\%$  to  $+21\%$  for skier B, and by  $-14\%$  to  $+13\%$  for skier C (Fig. 2a; Table A1). For snowboarder D, apparel changed  $c_D A$  by approximately the same amount as posture, inducing  $c_D A$  changes in the most common mid posture from  $-8\%$  to  $+15\%$  (Fig. 2b; Table A1). Applying individual two-parameter models including *apparel* in addition to the *body posture* (Eqs. (5) to (7)) increased the explanatory power distinctively ( $0.05 \leq \Delta R_{adj}^2 \leq 0.35$ ; Table A3), which confirmed the importance to consider apparel fit when estimating  $c_D A$ s of BA or SS athletes. In contrast, adding speed as third explanatory variable, did not improve the models remarkably ( $\Delta R_{adj}^2 \leq 0.01$ ). Therefore, speed was not retained as regressor of the individual  $c_D A$  models.

$$c_D A_{SkiB} = -0.142 + 0.461 \cdot h_{ext} - 0.043 \cdot X_{f1} + 0.098 \cdot X_{f2} \quad (R_{adj}^2 = 0.96) \tag{5}$$

$$X_{f1} = \begin{cases} 1 & \text{if } n_{fit} = 2 \\ 0 & \text{if } n_{fit} > 2 \end{cases} \quad X_{f2} = \begin{cases} 1 & \text{if } n_{fit} = 5 \\ 0 & \text{if } n_{fit} < 5 \end{cases} \quad n_{fit} = [2; 3; 5] \quad 0.95 \text{ m} \leq h_{ext} \leq 2.00 \text{ m}$$



**Fig. 2.** Measured drag areas and extension heights ( $n = 720$ ) of four skiers wearing regular apparel in low, mid and extended postures at speeds of 35, 60 and 85 km/h. b) Measured drag areas ( $n = 720$ ) of three snowboarders wearing regular apparel in mid ( $n_{pos} = 1$ ), extended ( $n_{pos} = 2$ ) and extended rotated ( $n_{pos} = 3$ ) postures at speeds of 35, 60 and 85 km/h.

$$c_{DA_{SKIC}} = -0.141 + 0.523 \cdot h_{ext} - 0.114 \cdot X_{f1} - 0.012 \cdot X_{f2} \quad (R_{adj}^2 = 0.91)$$

$$X_{f1} = \begin{cases} 1 & \text{if } n_{fit} = 2 \\ 0 & \text{if } n_{fit} > 2 \end{cases} \quad X_{f2} = \begin{cases} 1 & \text{if } n_{fit} = 4 & n_{fit} = [2; 3; 4] \\ 0 & \text{if } n_{fit} < 4 & 0.93 \text{ m} \leq h_{ext} \leq 1.98 \text{ m} \end{cases} \quad (6)$$

$$c_{DA_{SNBo}} = 0.410 + 0.031 \cdot X_{p1} + 0.103 \cdot X_{p2} - 0.028 \cdot X_{f1} + 0.062 \cdot X_{f2} \quad (R_{adj}^2 = 0.94)$$

$$X_{f1} = \begin{cases} 1 & \text{if } n_{fit} = 2 \\ 0 & \text{if } n_{fit} > 2 \end{cases} \quad X_{f2} = \begin{cases} 1 & \text{if } n_{fit} = 5 \\ 0 & \text{if } n_{fit} < 5 \end{cases} \quad n_{fit} = [2; 3; 5]$$

$$X_{p1} = \begin{cases} 1 & \text{if } n_{pos} = 1 \\ 0 & \text{if } n_{pos} > 1 \end{cases} \quad X_{p2} = \begin{cases} 1 & \text{if } n_{pos} = 3 \\ 0 & \text{if } n_{pos} < 3 \end{cases} \quad n_{pos} = [1; 2; 3] \quad (7)$$

The contribution of *anthropometric differences* was relatively small for the skiers with  $c_{DA}$  variations of up to 10% between athlete A and C in mid and extended posture, but considerably for the snowboarders with up to 21% deviation between A and C in mid posture (Table A1).

In contrast to the group level, significant *speed dependencies* were found for individual athletes (Table A4). Among the skiers, negative speed dependencies existed only for the larger athletes A and C in mid and high postures. For skier C, wearing wide fit apparel, a maximal  $c_{DA}$  variation was found from +15% to -13% over the tested speed range (referred to  $c_{DA}$  average over all speeds). This correlation disappeared wearing slim fit apparel. A slightly positive relationship between  $c_{DA}$  and speed was observed in low postures for all skiers, which was only significant for athlete B and D causing maximal  $c_{DA}$  variations from -5% to 7% (B, regular). For all tested snowboarders, a slight positive speed dependency was found for mid and extended postures causing maximal  $c_{DA}$  variations from -6% to 5% (D, regular), which weakened or disappeared when an upright rotated posture was adopted (Fig. 3b; Table A4).


In summary, the  $c_{DA}$  of skiers were dominated primarily by posture (77%  $c_{DA}$  variation), followed by apparel (30%), speed (28%) and individual differences (10%) (Fig. 2a). For snowboarders, posture (21%), apparel (23%), and individual differences (21%) influenced  $c_{DA}$  to a comparable extent, whereas speed (11%) had a smaller impact (Fig. 2b).

Drag areas for typical *airborne postures*, performing a mute grab at inflow directions of zero, 90 and 180° were  $0.379 \pm 0.005 \text{ m}^2$ ,  $0.567 \pm 0.010 \text{ m}^2$ , and  $0.382 \pm 0.005 \text{ m}^2$  for skier B and  $0.325 \pm 0.013 \text{ m}^2$ ,  $0.453 \pm 0.014 \text{ m}^2$ , and  $0.419 \pm 0.009 \text{ m}^2$  for snowboarder D. Corresponding lift areas were  $0.104 \pm 0.011 \text{ m}^2$ ,  $0.134 \pm 0.018 \text{ m}^2$ , and  $0.044 \pm 0.004 \text{ m}^2$  for skier B and  $0.008 \pm 0.019 \text{ m}^2$ ,  $0.057 \pm 0.014 \text{ m}^2$ , and  $0.040 \pm 0.014 \text{ m}^2$  for snowboarder D. The drag areas due to *vertical inflow* were  $0.412 \pm 0.071 \text{ m}^2$  for skier B and  $0.635 \pm 0.004 \text{ m}^2$  for snowboarder D.

#### 4. Discussion

To our knowledge, this study is the first to analyze air drag and lift in snowboard. Until now, modeling of BA and slopestyle jumps were performed using  $c_{DA}$  values from alpine skiing. This study showed that  $c_{DA}$  issued from alpine skiing studies (ranging from 0.25 to 0.5  $\text{m}^2$  for giant slalom,<sup>34</sup> and 0.15  $\text{m}^2$  for downhill discipline<sup>35</sup>) underestimated the true  $c_{DA}$  for BA and slopestyle.

Due to the current lack of any comparable data and the fact that regular apparel is most frequently worn during elite level competitions<sup>36</sup> (observations of 163 athletes at FIS World Cup at Seiser Alm 2018; Fig. A2), we are convinced that the models derived in this study provide a useful contribution to *improve the understanding and modeling of in-run mechanics* in slopestyle and BA. For skiers and snowboarders, wearing regular fit apparel, posture-dependent formulations were deduced for  $c_{DA}$  and  $c_{LA}$ . The speed dependency of  $c_{DA}$  was largely significant in the individual models, but not consistent in direction, and had smaller effect than apparel and posture. For skiers in mid and extended posture the  $c_{DA}$  was mostly negatively related with speed, which corresponds to findings on alpine skiers,<sup>37</sup> whereas for snowboarders all significant relations of  $c_{DA}$  with speed, were positive. This discrepancy might be caused by the different inflow (frontal vs. lateral) affecting the Reynolds Number range or slight changes of posture at higher speeds.<sup>37</sup> As speed effects were small compared to the other factors influencing  $c_{DA}$  in the in-run, these were neglected in our model.<sup>15,19</sup> We suggest that course builders use the posture-dependent models (Eqs. (3), (4), A1 and A2) to simulate an average in-run, and the individual, apparel and posture-dependent data (Fig. 3, Tables A1 & A2) as estimates for extreme aerodynamic coefficients. Course builders need to build jumps that allow for a certain range of speed at take-off, with a critical lower limit of take-off speed. However, this lower limit is not sharp, since it changes with external conditions and user groups. Therefore, a certain range is needed to compensate for headwind, snow conditions and differences in mass between sexes and snow friction between snowboards and skis. If in-run speed is at the low end, athletes have some capacity to compensate for this by pushing off at the start and reducing air drag by choosing a low posture during the in-run. This study shows that the capacity to compensate for limited speed in the in-run is substantially smaller for snowboarders than for skiers as the range of  $c_{DA}$  for realistic postures is about tripled for skiers compared to snowboarders and minimum  $c_{DA}$ s of skiers are approximately 0.1  $\text{m}^2$  smaller than of snowboarders. The reason for this difference may be that snowboarders' balance setting allows them to manipulate  $h_{ext}$  only in a small range compared to skiers. In addition, squatting does not lead to overlapping of body segments, and consequently does not



**Fig. 3.** a) Mean  $c_{DPA}$  values ( $\pm 2\sigma$ ) of one (wide/slim fit) or two trials (regular fit) for skiers A to D for low (red), mid (blue) and extended posture (black) with regular (circle), slim (bar) and wide fit (triangle) apparel. b) Mean  $c_{DPA}$  values ( $\pm 2\sigma$ ) of one (wide/slim fit) or two trials (regular fit) for snowboarders A, C and D for mid (red), extended (blue) and extended rotated (black) posture with regular (circle), slim (bar) and wide fit (triangle) apparel. Note the differently scaled y-axes for skiers and snowboarders.

affect the frontal area as it does in skiing. However, snowboarders should be aware that upper body orientation has a strong influence on the  $c_{DPA}$ , and can be used to reduce air drag when speed is critical. As

more and more high-level competitions are held using the same course for males and females, ski and snowboard course construction should pay particular attention to snowboarders' limited ability to compensate

for head winds and elevated snow friction using alterations in body posture. However, this study also shows that athletes can substantially reduce air drag with their choice of apparel, if take-off speed is critical: Wearing wide fit apparel compared to slim fit increased  $c_{DA}$  as much as changing from the mid to the extended posture for skiers and changing to the extended rotated posture for snowboarders (Fig. 2).

For modeling of the flight phase, this study contributed the first values for drag and lift. For compact posture holding a grab, we recommend using rounded averages over all tested inflow directions for  $c_{DA}$  and  $c_{LA}$  of 0.44 m<sup>2</sup> and 0.09 m<sup>2</sup> for skiers, and 0.40 m<sup>2</sup> and 0.04 m<sup>2</sup> for snowboarders. Although the results show distinct differences depending on the inflow direction, we consider averaging from 0° to 180° is reasonable, as athletes mostly aim to rotate full numbers of semi twists.

The limitations of the test design included the small number of athletes, the lack of small and female athletes as well as a rather rough quantification of the apparel fit. The number and choice of athletes was constrained by the financial means for wind tunnel testing and the fact, that the athletes tested in the wind tunnel were included in a second study on snow friction, where the individual air drag values from wind tunnel testing were applied to distinguish air drag and snow friction for these athletes. Although the tested athletes did not represent the overall population of slopestyle and BA athletes, the test efficiency was kept as high as possible within the study's financial limits.

## 5. Conclusion

This study presented for the first time a data set that describes the aerodynamic coefficients for slopestyle and BA athletes. The presented models allow the calculation of aerodynamic coefficients as a function of the main influencing factors posture and apparel. The models and data allow the estimation of typical air drag values including the upper and lower limits and differences between snowboarders and skiers, and can hence be applied to improve the validity of jump simulations.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jsams.2021.05.005>.

## Funding information

The authors thank the International Olympic Committee (IOC) and the Mid-Sweden University for their financial contribution to the study.

## Declaration of interest statement

The authors have no conflict of interest to report.

## Confirmation of ethical compliance

The study was approved by the ethics committee at the Norwegian School of Sport Sciences, and the Norwegian Centre for Research Data, and conducted according to the Declaration of Helsinki. All subjects gave their written informed consent prior to participation.

## References

1. Soligard T, Steffen K, Palmer-Green D et al. Sports injuries and illnesses in the Sochi 2014 Olympic Winter Games. *Br J Sport Med* 2015;49(7):441-447. doi:10.1136/bjsports-2014-094538.
2. Torjussen J, Bahr R. Injuries among elite snowboarders (FIS Snowboard World Cup). *Br J Sports Med* 2006;40(3):230-234.
3. Florenes TW, Heir S, Nordsletten L et al. Injuries among World Cup freestyle skiers. *Br J Sports Med* 2010;44(11):803-808.
4. Florenes TW, Nordsletten L, Heir S et al. Injuries among World Cup ski and snowboard athletes. *Scand J Med Sci Sports* 2012;22(1):58-66.
5. Soligard T, Palmer D, Steffen K et al. Sports injur. Published online and illness incidence in t018 Olympic Winter Games: a prospective study of 2914 athletes from 92 countries. *Br J Sports Med* 2019. Published online. doi:10.1136/bjsports-2018-100236.
6. Russell K. *The relationship between injuries and terrain park feature use among snowboarders in Alberta (thesis)*, Calgary, Canada, Univ Calgary, 2011. Published online.
7. Henrie M, Petron D, Pepper M et al. Comparison of ski and snowboard injuries that occur inside versus outside terrain parks: 2001: board# 130 June 3 9:00 AM–10:30 AM. *Med Sci Sport Exerc* 2010;42(5):473-474.
8. Dohjima T, Sumi Y, Ohno T et al. The dangers of snowboarding: a 9-year prospective comparison of snowboarding and skiing injuries. *Acta Orthop* 2001;72(6):657-660.
9. Scher I, Shealy J, Stepan L et al. *Terrain park jump design: would limiting equivalent fall height reduce spine injuries?*, *Skiing Trauma and Safety*, ASTM International, 2015.
10. Shealy JE, Scher I, Stepan L et al. Jumper kinematics on terrain park jumps: relationship between takeoff speed and distance traveled. *J ASTM Int* 2010;17(10):1.
11. McNeil JA, McNeil JB. Dynamical analysis of winter terrain park jumps. *Sport Eng* 2009;11(3 LB-1028):159-164.
12. Levy D, Hubbard M, McNeil JA et al. A design rationale for safer terrain park jumps that limit equivalent fall height. *Sport Eng* 2015:1-13. Published online.
13. Moore JK, Hubbard M. skijumpdesign: a ski jump design tool for specified equivalent fall height. *J Open Source Softw* 2018;3(28):818. doi:10.21105/joss.00818.
14. Hubbard M, McNeil J, Petrone N et al. *Impact performance of standard tabletop and constant equivalent fall height snow park jumps*, *Skiing Trauma and Safety*, ASTM International, 2015.
15. McNeil JA, Hubbard M, Swedberg A. Designing tomorrows snow park jump. *Sport Eng* 2012;15(1):1-20.
16. Böhm H, Senner V. *Safety in big jumps: Relationship between landing shape and impact energy determined by computer simulation*, *Skiing Trauma and Safety*, ASTM STP, 2008. p. 1510.
17. Hubbard M, Swedberg A. Design of Terrain Park jump landing surfaces for constant equivalent fall height is robust to uncontrollable factors. *Ski Trauma Saf* 2012;19:75-94.
18. Shealy JE, Stone F. Tabletop jumping: engineering analysis of trajectory and landing impact. *J ASTM Int* 2008;5(6):1.
19. Schindelwig K, Platzter HP, Mössner M et al. Safety recommendations of winter terrain park jumps into airbags. *J Sci Med Sport* 2019;22. doi:10.1016/j.jsams.2018.09.229.
20. McNeil J. Modelling the 'Pop' in winter terrain park jumps. *Ski Trauma Saf* 2012;19:1553.
21. Gilgien M, Kröll J, Spörri J et al. Application of dGNSS in alpine ski racing: basis for evaluating physical demands and safety. *Front Physiol* 2018;9(MAR):145. doi:10.3389/fphys.2018.00145.
22. Alam F, Chowdhury H, Moria H. A review on aerodynamics and hydrodynamics in sports. *Energy Procedia* 2019. doi:10.1016/j.egypro.2019.02.158.
23. Watanabe K, Ohtsuki T. Postural changes and aerodynamic forces in alpine skiing. *Ergonomics* 1977. Published online. doi:10.1080/001401377080online.
24. Kaps P, Nachbauer W, Mössner M. *Determination of kinetic friction and drag area in Alpine skiing*, *Skiing Trauma and Safety*, Vol. Tenth, 2009. doi:10.1520/stp37926s.
25. Nørstrud H. *Basic aerodynamics*, CISM International Centre for Mechanical Sciences, Courses and Lectures, 2008. doi:10.1007/978-3-211-89297-8\_1.
26. Ainegren M, Jonsson P. Drag area. Published online, frontal area and drag cross-country skiing techniques. *Proceedings* 2018. Published online. doi:10.3390/proceedings2060313.
27. Brownlie L, Larose G, D'Auteuil A et al. Factors affecting the aerodynamic drag of alpine skiers. *Procedia Eng* 2010. ;2(2):2375-2380. <http://www.sciencedirect.com/science/article/pii/S187705810002560>.
28. Barelle C, Tavernier M. Experimental creation of a model for the aerodynamic braking coefficient in Alpine skiing. *Arch Physiol Biochem* 2000;108(1-2):138.
29. Elfmark O, Bardal LM. An empi. Published online, model of aerodynamic skiing. *Proceedings* 2018. Published online. doi:10.3390/proceedings2060310.
30. Steiling D, Hauser A. *Large wind tunnel Emmen LWTE facility description*, Ruag Schweiz AG, 2018.
31. Boyd E. The growth of the surface area of the human body. *J R Stat Soc* 1937;100(1):111. doi:10.2307/2980290.
32. Tutz G. *Regression for categorical data*, 2011. doi:10.1017/CBO9780511842061.002.
33. Montgomery DC. *Introduction to linear regression analysis*, 5th ed., Wiley, 2012.
34. Meyer F, Le Pelley D, Borrani F. Aerodynamic drag modeling of Alpine skiers performing giant slalom turns. *Med Sci Sport Exerc* 2011;44(6):1109-1115.
35. Barelle C, Ruby A, Tavernier M. Experimental model of the aerodynamic drag coefficient in alpine skiing. *J Appl Biomech* 2004;20(2):167-176.
36. Wolfsperger F, Meyer F, Guillaume SGM. *Towards more valid simulations of slopestyle and big air jumps*, In: Lenka Stepan P, Irving Scher P, Rick Greenwald PD, eds. *Book of Abstracts of the 23rd International Congress on Snow Sport Trauma and Safety*, ISSS, 2019. p. 92.
37. Elfmark O, Reid R, Bardal LM. Blockage correction and Reynolds number dependency of an Alpine skier: a comparison between two closed-section wind tunnels. *Proceedings* 2020;49(1). doi:10.3390/proceedings2020049019.