

Silva, L., Castro, M. A., Marta, S., Almeida, J., Vaz, J., Pezarat-Correia, P., Cabri, J. M. H. (2014). Cross-cultural adaptation and validation of the Portuguese Survey of musculoskeletal conditions, playing characteristics and warm-up patterns of golfers. *Journal of Back and Musculoskeletal Rehabilitation*, under utgivelse. doi:10.3233/BMR-140582

---

Dette er siste tekst-versjon av artikkelen, og den kan inneholde små forskjeller fra forlagets pdf-versjon. Forlagets pdf-versjon finner du på [www.iospress.nl](http://www.iospress.nl):  
<http://dx.doi.org/10.3233/BMR-140582>

---

This is the final text version of the article, and it may contain minor differences from the journal's pdf version. The original publication is available at [www.iospress.nl](http://www.iospress.nl): <http://dx.doi.org/10.3233/BMR-140582>

---

## TITLE PAGE

# Cross-cultural adaptation and validation of the Portuguese Survey of musculoskeletal conditions, playing characteristics and warm-up patterns of golfers

**Silva, Luís**<sup>1</sup>; Faculdade de Motricidade Humana, Universidade de Lisboa, Estrada da Costa, 1499-002 Cruz Quebrada – Dafundo, Portugal

**Castro, Maria António**; Coimbra College of Health Technology, Polytechnic Institute of Coimbra, Rua 5 de Outubro - SM Bispo, Apartado 7006, 3046-854 Coimbra

**Marta, Sérgio**; Faculdade de Motricidade Humana, Universidade de Lisboa, Estrada da Costa, 1499-002 Cruz Quebrada – Dafundo, Portugal

**Almeida, José**; Faculdade de Motricidade Humana, Universidade de Lisboa, Estrada da Costa, 1499-002 Cruz Quebrada – Dafundo, Portugal

**Vaz, João**; Faculdade de Motricidade Humana, Universidade de Lisboa, Estrada da Costa, 1499-002 Cruz Quebrada – Dafundo, Portugal

**Pezarat-Correia, Pedro**; Faculdade de Motricidade Humana, Universidade de Lisboa, Estrada da Costa, 1499-002 Cruz Quebrada – Dafundo, Portugal

**Cabri, Jan**, Norwegian School for Sport Sciences, Dept. Physical Performance, P.o. box 4014, Ullevål Stadion, 0806 Oslo, Norway

<sup>1</sup> Corresponding author

Mobile phone: +351 927121682; email: [luisbsilva@hotmail.com](mailto:luisbsilva@hotmail.com); Institutional fax: (+351) 21 415 12 48.

## ABSTRACT

**BACKGROUND:** The University of Western Ontario Questionnaire for Musculoskeletal Conditions in Senior Golfers (MSK Golfers) was developed in Canada because of a lack of knowledge concerning musculoskeletal conditions directly related to golf play and warm-up, although the high injury incidence in golf practice. This lack of epidemiological measures also exists for the Portuguese golf population.

**OBJECTIVE:** The purpose of this study was to translate and cross-culturally adapt the MSK Golfers questionnaire into Portuguese and to test its construct validity and reproducibility.

**METHODS:** The MSK Golfers was translated from English to Portuguese and tested for psychometric properties. Sixty-one golfers, aged between 14 and 70 years and with at least 1 year of practice in golf, were recruited. The validity of the MSK Golfers was assessed by evaluating data quality (missing, floor and ceiling effects). Reproducibility analysis included intra-class correlation coefficients (ICC) (2,1) and Cohen's Kappa coefficient.

**RESULTS:** The ICC values for continuous items ranged from 0.634 to 0.998 with the exception of one item on golf activity. Kappa statistics for the categorical items ranged between 0.714 and 1.00.

**CONCLUSIONS:** The Portuguese version of the MSK Golfers, including playing characteristics and warm-up patterns of golfers, showed a high reliability for a golfing population with an age range of 14 to 70 years.

**Keywords:** Questionnaire, Sport Medicine, Musculoskeletal Injuries, Golf

1

2

## 1. INTRODUCTION

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

Golf is practiced by people of various ages and skill levels, who undertake widely varying amounts of practice, and have objectives including competition, leisure and tourism. Although golf is considered a sport with a moderate aerobic component, during the swing, the body segments and club form an energy chain to transfer to the ball the necessary velocity and trajectory that maximize distance and accuracy [1,2]. Usually, the variables used to measure swing outcome are ball displacement, shot accuracy, club head velocity, and club face angle, with performance being better when golfers focus on these swing outcomes rather than a given body movement [3]. To achieve these objectives, the golfer applies extreme muscle strength [4] at maximum ranges of motion [5], and with high-velocity joint movement [6,7]. The repeated execution of the swing during a round of golf can help to explain the high incidence of golf-related injuries. Lindsay et al [8] reported that more than 70% of golf players experienced injuries severe enough to result in them playing at an unsatisfactory skill level over a short period of time. Age-related golf practice, rates of participation, declines in strength, flexibility and coordination, and the efficacy of conditioning programs for improving performance and preventing injuries are major areas of research related to senior golfers [8,9]. Playing golf provides moderate intensity exercise for seniors, but musculoskeletal injuries can result from unsafe participation and/or due to preexisting musculoskeletal disorders. Cann et al [9] considered four specific concerns for the senior golfer: injury rehabilitation coordinated by therapists; warm up routines; club-fitting/coaching on proper technique; and pre-season conditioning programs. However, there is a lack of studies separating the effects of swing modification from physical rehabilitation, and on the effectiveness of changes in swing leading to a decrease in complaints such as low back pain [10].

26           The main complaint reported is low back pain, followed by elbow and wrist  
27 problems [6,11,12] but golf-related practice risks and health benefits have not been fully  
28 established and the available literature related to golf injuries is descriptive and  
29 controversial [12]. There is a need for tools that help the understanding of how exercise  
30 habits and injury incidence may be related with respect to the mechanisms of golf  
31 injuries. For instance, investigation is needed on the role of specific physical activities,  
32 skills and overuse (such as repeated swing trials at driving ranges) in the cause,  
33 prevention and rehabilitation of acute and chronic disorders [13]. Studies are also  
34 needed on changes in muscle activation, such as in the erector spinae, in golfers with  
35 low back pain [14]. The available literature shows the need for an instrument that can  
36 categorize the golfing population, particularly senior players [8,15]. Studies on temporal  
37 parameters in golfers with and without low back pain have used the Rolling Morris  
38 questionnaire [14,16], but there is a lack of instruments to satisfactorily group golfers,  
39 particularly in terms of relating low back pain to golf practice.

40           A survey of musculoskeletal conditions, playing characteristics and warm-up  
41 patterns in golfers was developed by Fox et al. [15] in response to the lack of information  
42 concerning musculoskeletal conditions affecting senior golfers. The questionnaire is a  
43 short participant-reported form, with 37 continuous (eg. golf handicap, minutes of warm-  
44 up) and categorical variables (eg. presence or absence of any injuries in the past 3 years  
45 related to golf activities, which caused the player to stop or modify their golf practice).  
46 The survey takes about 10 minutes to complete. Palmer et al. [17] applied this survey to  
47 100 golf club members, reporting five groups of musculoskeletal (MSK) conditions:  
48 36.8% of those surveyed had rheumatologic conditions; 18.4% had tendinitis; 18.4%  
49 reported muscle strains; 10.5% had vertebral/disk pathologies, and 15.8% noted other  
50 conditions. Half of the respondents reported having suffered from MSK conditions, and  
51 a large proportion of golfers indicated they spent just few minutes on warm-up and  
52 stretching exercises before performing the swing.

53 To be able to characterize Portuguese golf players and their usual golfing  
54 behavior, this survey of musculoskeletal conditions, playing characteristics and warm-up  
55 patterns of golfers (MSK Golfers) was translated and cross-culturally adapted for the  
56 Portuguese population. The Portuguese version was then tested for psychometric  
57 properties in terms of validity and reproducibility in Portuguese-speaking golfers.

58

59

## 2. METHODS

60

### 2.1. Study Design

61

62

63

64

The study was carried out using a two-step procedure: firstly, the MSK Golfers  
was translated and cross-culturally adapted; and secondly, the Portuguese version of  
MSK Golfers was tested for psychometric properties, reliability and validity, in a cross-  
sectional design with a 2-week follow-up for test-retest evaluation.

65

66

### 2.2. Translation and cross-cultural adaptation

67

68

69

70

71

72

Translation and cross-cultural adaptation of the MSK Golfers was performed  
according to international guidelines [18,19] under license from the original author. The  
English MSK Golfers was translated into Portuguese independently by two Portuguese  
native translators (one a doctor with experience in sports medicine and one a  
professional translator). The translations were discussed in a consensus panel to  
produce a preliminary version before it was translated back into English.

73

74

75

76

77

Two translators and native speakers of English, who were blinded to the original  
MSK Golfers, performed the back-translation. A second panel, consisting of a  
physiotherapist, a golf coach and the researchers in our group, reviewed all translations.  
The committee discussed discrepancies until consensus on a pre-final version was  
achieved. The goal of the pre-final Portuguese MSK Golfers was that it should be as

78 concise and easy to understand as possible. The pre-final Portuguese version of MSK  
79 Golfers was then tested on six golfers. None of the golfers had difficulties in  
80 understanding the meaning of items or responses. A suggestion to adapt the questions  
81 according to the Portuguese golf season was made which was not considered by the  
82 expert panel. A third expert consensus panel completed the final version of MSK Golfers.  
83 The English and Portuguese versions are presented in additional file (Appendix).

84 In total, 61 golfers were included in the study. Eligible participants were golfers  
85 with at least 1 year of practice in golf, aged 14 and over, who were able to speak, read  
86 and write in Portuguese. The exclusion criteria were having less than 1 year's golfing  
87 experience and being under the age of 14. Baseline characteristics for the test-retest  
88 subgroup are presented in Table 1. All participants received written and oral information  
89 about the study. Signed informed consent was obtained from all participants. The study  
90 was approved by the Ethical Committee of the Faculty of Human Kinetics - University of  
91 Lisbon (FMH – UL) in 2011.

92

93 Insert table 1 here.

94

### 95 **2.3. Procedures and Measures**

96 Participants consented to participate by filling in the MSK Golfers test and  
97 undergoing a retest, preferably with a two-week interval.

98 The MSK Golfers comprises 37 items focusing on golfers' habits. The original  
99 MSK Golfers was evaluated in senior golfers (over 50 years of age) and was found to be  
100 a reliable and valid measure of golfers' playing conditions [15], involving 6 questions  
101 related to demographics, 7 questions about the player's general history of illness and  
102 injury, 3 questions about the respondent's golf game, 2 questions about their golf swing,

103 6 questions about their golf course, 4 questions about the amount they played, 6  
104 questions about warm-up and exercises and 3 questions about golf injuries (see  
105 Appendix).

106

#### 107 **2.4. Statistical Analysis**

108 The validation procedures were similar to the original, in order to facilitate  
109 reproducibility and comparison. Agreement between answers to the test and re-test  
110 questionnaires was assessed, using a two-week interval. This period was chosen to  
111 avoid biases in answers due to factors such as fatigue, learning, or memory effects, and  
112 to provide close enough periods to avoid genuine changes in the variables that were  
113 being measured. Descriptive statistics for quantitative variables such as “*handicap*” or  
114 “*On average how many yards do you hit your drive*”, and proportions of transformed  
115 ordinal variables such as “*Typically, how often are you aware of low back pain after*  
116 *playing 18 holes?*” were presented as mean  $\pm$  standard deviation. Nominal qualitative  
117 variables such as “*Do you routinely perform any of your golf stretches away from the*  
118 *course/practice range?*” (Portuguese version) were presented as the respective  
119 frequencies.

120 The validity of the Portuguese MSK Golfers was assessed by evaluating data  
121 quality (missing, floor and ceiling effects). Test-retest reliability was analyzed using  
122 Cohen’s Kappa statistic for items with nominal scales and using Intraclass Correlation  
123 Coefficients with two-way random single measures (2,1) (ICC) for items with quantitative  
124 and transformed ordinal scales [20-22]. All data were processed in SPSS 19.0 [23].  
125 Pearson’s *r* was also calculated for items with categorical scales, to provide a reference  
126 for assessing ICC.

127



### 3. RESULTS

128

129           The survey of musculoskeletal conditions, playing characteristics and warm-up  
130 patterns of golfers showed acceptable psychometric properties in terms of  
131 comprehensibility, consistency (agreement), construct validity and reproducibility when  
132 applied to golfers with an age range of 14–70 years.

133           The reliability results of the Past Medical History items showed a high agreement  
134 between the two occasions (Table 2), despite the item about knowing any  
135 Contraindications to Activity showing a low Kappa value (0.659); overall reliability was  
136 acceptable. The Kappa values for Heart Condition, Chest Pain with Activity and chest  
137 Pain at Rest are not shown because there was perfect agreement.

138

139           Insert table 2 here.

140

141           The items that measured the playing level of the subject (Table 3) showed a high  
142 level of agreement between the two occasions, with the worst value being the ICC for  
143 the “On average, how many yards do you hit your drive” showing an ICC of 0.837. The  
144 variable “Are your golf clubs customized to fit your golf swing” (fitting) showed a  
145 difference between the two occasions of 3.3%, but with a high ICC of 0.997.

146           Of the Transportation on Golf Course Items (Table 4), only the “*On average, how*  
147 *often do you push your clubs around the course on a cart?*” item had an ICC lower than  
148 0.9 (ICC=0.705). All the others items had ICC values higher than 0.9.

149

150           Insert table 3 and table 4 here.

151

152 All Reliability of Frequency of Golf Activity Items (Table 5) showed an ICC higher  
153 than 0.7, with the exception of the Class Sessions Off Season item (ICC=0.456), being  
154 the lowest value found in the entire questionnaire. Reliability of Frequency of Golf Activity  
155 items achieved a maximum ICC value of 0.920. For these items it was noted that the  
156 lower handicap players tended not to respond because they undertake daily practice.

157

158 Insert table 5

159

160 Interestingly, despite the time used for warm-up (Table 6) ranging up to 9  
161 minutes, and this item showing an ICC of 0.793, some golfers had no specific warm-up  
162 routine habits. The duration of the warm-up used for stretching was 3.95 and 3.64  
163 minutes for the first and second occasions, respectively (ICC=0.832). The highest  
164 concordance found for the reliability of warm-up and exercise habits items was in the  
165 question "*Do you routinely participate in a cardiovascular conditioning program apart*  
166 *from golfing?*" showing a kappa of 0.964, but the other nominal variables gave kappa  
167 values higher than 0.7.

168

169 Insert table 6

170

171 Table 7 indicates the reliability levels for the injury items. The new added question  
172 "*How much time have you stopped*" achieved an ICC of 0.672, and golfers with injuries  
173 reported having stopped for 25 and 27 days for occasions 1 and 2, respectively. The  
174 question that related low back pain to sports practice also yielded a high ICC of 0.905.  
175 Both the general injury incidence and musculoskeletal injury incidence were higher than  
176 20%, showing a high kappa value.

177

178           Insert table 7

179

180

#### 4. DISCUSSION

181           In this article the process of cross cultural adaptation of the MSK golfer to  
182 Portuguese was described as well as its validity and reliability.

183           The survey of musculoskeletal conditions, playing characteristics and warm-up  
184 patterns of golfers for the Portuguese population showed a high reliability. This study  
185 took into account the recommendations stated by Portney & Watkins [21] for ICC values.  
186 A good reliability for ICC is considered to be  $\geq 0.75$ , however, the authors stated that for  
187 most clinical measurements, reliability should exceed 0.90 to ensure reasonable validity.  
188 In this study, of the variables measured, 44.5% exceeded ICC = 0.9 and 72.4% were  
189 above ICC = 0.7. Also, in the original version, the named "skill" items were highly reliable,  
190 with ICC's above 0.90, and "Transportation on the Course" variables ranged between  
191 0.692 and 0.921. The Portuguese validation showed "skill" ICC's higher than 0.9 and  
192 close to 1.0 (Table 3). Only the drive range had an ICC = 0.837. For the Transportation  
193 on Golf Course Items, the Portuguese version also showed ICC's greater than 0.9, with  
194 the exception of "Proportion of Rounds Push" item (ICC = 0.705).

195           The Portuguese version includes a question to complement the survey section  
196 on golf-related injuries. Question 35 states "*Have you suffered any injuries in the past 3*  
197 *years while playing or practicing golf, which caused you to stop or modify your game?*"  
198 When the respondent answers "yes," in the Portuguese version he/she is referred to the  
199 question "*How much time have you stopped*". The unit measure of this question is "*days*".  
200 Original authors felt this upgrade very useful.

201 The reliability of Past Medical History items was similar to the original version with  
202 the Kappa showing a high agreement, but the first three items (heart condition, chest  
203 pain with activity, and Chest Pain at Rest) in the present study showed a frequency of  
204 100% for the answer “no” on both occasions. The Kappa values for these variables are  
205 not presented because they are determined by cross-tabulating ratings for two coders,  
206 and the agreement expected by chance is determined by the marginal frequencies of  
207 each rate [20]. In the original version more than 39.5% of respondents answered  
208 “Orthopaedic Problem Aggravated by Activity”, while the present study had 16.7% of  
209 respondents in this category. This group of items was based on the Physical Activity  
210 Readiness (PAR-Q) form from Public Health Agency of Canada and the Canadian  
211 Society for Exercise Physiology [24], validated for people with ages ranging from 15 to  
212 69 years. In the present study one respondent was 14 years old, which presents a  
213 limitation. Fox et al [15] noted similar age-related limitations with the Par-Q (British  
214 Columbia Ministry of Health, 1978 cited [15]), because their version was validated for  
215 individuals aged 18 to 69 and they had participants aged 70 years and older. They  
216 reported the use of Par-Q only as an accepted measure of general health, not as a  
217 screening device and not critical in these study types.

218 Comparing Portuguese and Canadian versions of the survey with respect to  
219 “Warm-up and Exercise Habits”, Portuguese golfers appeared to spend more time  
220 warming-up prior to playing, with  $9.15 \pm 5.82$  minutes on the first occasion and  $9.02 \pm 4.66$   
221 minutes on the second occasion, while the Canadian golfers’ mean values on occasions  
222 one and two were  $4.47 \pm 0.59$  and  $4.07 \pm 0.54$  minutes, respectively. The ICC’s for both  
223 versions were similar, ICC = 0.793 in the Portuguese version and ICC = 0.750 for the  
224 Canadians. The amount of this period which was devoted to stretching showed for  
225 occasions one and two:  $3.95 \pm 3.61$  and  $3.64 \pm 3.98$  for the Portuguese participants (ICC  
226 = 0.832) and  $2.9 \pm 0.49$  and  $2.7 \pm 0.50$  minutes for Canadians, respectively (ICC = 0.791).  
227 Parmer et al. [17] considered senior’ golf warm-up activities as “surprisingly limited.”

228 They found answers indicating that some spent less than 1 minute, both in warm-up  
229 (16.6%) and stretching (36%), and that over 75% of all participants spent less than five  
230 minutes. Golfers with MSK conditions spent more time warming-up than those who had  
231 not had that experience.

232 The imbalance in the relationship between the samples in terms of sex ratios is  
233 related to the actual Portuguese golf population, and also applies to specific injuries. In  
234 the literature, there is a higher percentage of male golfers [25,26], but shoulder, knee  
235 and elbow injuries associated to overuse were found to be similar in both sexes [11,27].  
236 The findings on the most likely area of injury in female golfers were elbow and lower  
237 back, related to overuse, poor conditioning, and incorrect swing mechanisms [28].

238 In this study, the injury incidence was low compared with the data reported by  
239 Lindsay et al [8] and Fox et al [15]. However, both validations showed a relative absence  
240 of self-identified musculoskeletal injuries linked to golf participation. These differences  
241 could be associated with the inclusion of young athletes in the present study. Compared  
242 with the original MSK Golfers test, the Portuguese version covered a wider age range.  
243 Of the participants, 6.6% of the participants were aged 14 to 18 years, 45.9% were  
244 between 18 and 40 years of age, 31.1% were between 40 and 60, and the other 16.4%  
245 were over 60 years of age. The differences found between genders reflect the  
246 Portuguese reality in golf. Fox et al [15] used participants aged 50 years and older, and  
247 one-half of their respondents stated a pre-existing musculoskeletal condition affecting  
248 golf practice. Cabri et al. [12] reported that practicing golf for long periods could be the  
249 main reason for the prevalence of injuries in players aged between 50 and 65 years. This  
250 population has more probability of injury due to age-related changes such as decreasing  
251 strength, flexibility, coordination and balance. Also, many senior golfers already have  
252 preexisting MSK conditions [17]. Recreational golfers are not exposed to the repetitive  
253 stresses experienced by professional players, who spend significant time practicing at  
254 the driving range, the putting green, and playing the course [15,28]. Recreational golfers

255 usually have more options for varying the type of swing, method of transport and total  
256 duration of play [17]. Controversially, injury rates in amateur golfers may vary between  
257 mean values of 1.19–1.31 per year, while professionals have a higher prevalence with  
258 ratios near 2 injuries per year due to the time spent playing [27], rather than technical  
259 aspects or warm-up habits [28,29].

260 The health benefits of golf are related to the amount of physical activity during  
261 the game, including the method of transportation around the course. The respective  
262 items in the Portuguese version of the test showed a higher reliability than the original.  
263 In the Canadian version, the proportion using power carts was approximately 71%, unlike  
264 the 17% reported by Palmer et al. [17] and the 20% in the present study, according to  
265 which the respondents carry their clubs for approximately 43% of the rounds. Similar,  
266 Palmer et al. [17] reported that two-thirds of the respondents had a preference for walking  
267 the golf course at least 50% of the time. Careless driving and improper use of power  
268 carts were associated with traumatic injuries [30,31].

269 The question about the incidence of low back pain after 18 holes showed a mean  
270 value of about 18% of the time. This result is lower than expected, since low back pain  
271 is the major complaint among golfers [32-35] and is the main cause for a reduction in  
272 playing and skill level decrease in elite golfers [32]. Nevertheless, it is necessary take  
273 into account the specificity of the question when it asks “after 18 holes”, in other words,  
274 after a full round of the golf course. Besides the question specificity, Palmer et al. [17]  
275 also found higher incidence of low back pain, with 42% of the respondents having  
276 complaints.

277

278

## 5. CONCLUSION

279 The survey of musculoskeletal conditions, playing characteristics and warm-up  
280 patterns of golfers for the Portuguese population showed high reliability. The question

281 about golf practice time lost due to injury, which was added in the Portuguese version,  
282 also showed good reliability. In the Portuguese version the age range was larger than in  
283 the original version, but the reliability remained high. This survey demonstrates an  
284 excellent applicability and reliability and can be regarded as valid and useful for the  
285 characterization of the Portuguese golf population.

286

## 287 **Acknowledgments**

288 The project “Neuromuscular activity in the golf swing with implications for the  
289 practice and in the prevention of overuse injuries” was supported by the Portuguese  
290 Foundation for Science and Technology (FCT) fund (PTDC/DES/105176/2008).

291 The authors would like to thank to David Lindsay and Filomena Carnide for their  
292 comments on the manuscript. The authors would like to thank to Judy McDonald for  
293 English review.

294

295

296

297

298

299

300

301

302

## REFERENCES

303

- 304 [1] Farrally MR, Cochran AJ, Crews DJ, Hurdzan MJ, Price RJ, Snow JT, et al. Golf  
305 science research at the beginning of the twenty-first century. *J Sports Sci.* 2003;  
306 21(9): 753-765.
- 307 [2] Hume PA, Keogh J, Reid D. The role of biomechanics in maximising distance and  
308 accuracy of golf shots. *Sports Med.* 2005; 35(5):429-49.
- 309 [3] Keogh JW, Hume PA. Evidence for biomechanics and motor learning research  
310 improving golf performance. *Sports Biomech.* 2012; 11(2):288-309.
- 311 [4] Hosea TM. and Gatt CJ. Back pain in golf. *Clin Sports Med.* 1996; 15(1): 37-53.
- 312 [5] Hosea TM, Gatt CJ, Galli KM, Langrana NA, Zawadasky JP. Biomechanical analysis  
313 of the golfer's back. In: Cochrane AJ, editor. *Science and Golf.* London: Chapman  
314 & Hall; 1990. pp. 43-48.
- 315 [6] McCarroll M, Retting AC, Shelbourne K. Injuries in the amateur golfer. *Phys*  
316 *Sportsmed.* 1990; 18: 122-126.
- 317 [7] Zheng N, Barrentine SW, Fleisig GS, Andrews JR. Kinematic Analysis of Swing in  
318 Pro and Amateur Golfers. *Int J Sports Med.* 2008; 29(6): 487-493.
- 319 [8] Lindsay DM, Horton JF, Vandervoort AA. A Review of Injury Characteristics, Aging  
320 Factors and Prevention Programmes for the Older Golfer. *Sports Med.* 2000;  
321 30(2): 89-103.
- 322 [9] Cann AP, Vandervoort AA, Lindsay DM. Optimizing the benefits versus risks of golf  
323 participation by older people. *J Geriatr Phys Ther.* 2005; 28(3): 85-92.
- 324 [10] Gluck GS, Bendo JA, Spivak JM. The lumbar spine and low back pain in golf: a  
325 literature review of swing biomechanics and injury prevention. *Spine J,* 2008;  
326 8(5), 778-788.



- 327 [11] Gosheger G, Liem D, Ludwig K, Greshake O, Winkelmann W. Injuries and overuse  
328 syndromes in golf. *Am J Sports Med.* 2003; 31(3): 438-443.
- 329 [12] Cabri J, Sousa J, Kots M, Barreiros J. Golf-related injuries: a systematic review. *Eur*  
330 *J Sport Sci.* 2009; 9(6):353–366.
- 331 [13] Suckel A. [Sports medicine analysis of golf "swing" and lesions occurring during golf  
332 practice]. *Sportverletz Sportschaden.* 2002; 16(1):31-35
- 333 [14] Cole MH, Grimshaw PN. Electromyography of the trunk and abdominal muscles in  
334 golfers with and without low back pain. *J Sci Med Sport.* 2008; 11(2): 174-181.
- 335 [15] Fox E, Lindsay DE, Vandervoort AA. Musculoskeletal Injury Questionnaire for Senior  
336 Golfers. In Thain, E, editor. *Science and Golf IV. Proceedings of the World*  
337 *Scientific Congress of Golf.* St. Andrews, Scotland, July 22-26, 2002. pp. 88-99.  
338 London: Routledge.
- 339 [16] Horton JF, Lindsay DM, Macintosh BR. Abdominal muscle activation of elite male  
340 golfers with chronic low back pain. *Med Sci Sports Exerc.* 2001; 33(10): 1647-  
341 1654.
- 342 [17] Palmer JL, Young SD, Fox E, Lindsay DM., Vandervoort, A. A. Senior Recreational  
343 Golfers: a survey of musculoskeletal conditions, playing characteristics, and  
344 warm-up patterns. *Physiother Can.* 2003; 55(2): 79-85.
- 345 [18] Beaton DE, Bombardier C, Guillemin F, Ferraz MB. Guidelines for the Process of  
346 Cross-Cultural Adaptation of Self-Report Measures. *Spine.* 2000; 25:3186-3191.
- 347 [19] Guillemin F, Bombardier C, Beaton D. Cross-cultural adaptation of health-related  
348 quality of life measures: literature review and proposed guidelines. *J Clin*  
349 *Epidemiol.* 1993; 46:1417-1432.

- 350 [20] Hallgren, KA. Computing Inter-Rater Reliability for Observational Data: An Overview  
351 and Tutorial. *Tutor Quant Methods Psychol.* 2012; 8(1): 23-34.
- 352 [21] Portney L, Watkins M. *Foundations of Clinical Research: Applications to Practice.*  
353 3rd ed. Upper Saddle River, New Jersey: Pearson Prentice Hall; 2009.
- 354 [22] Shrout PE, Fleiss JL. Intraclass Correlations: uses in assessing rater reliability.  
355 *Psychol Bull.* 1979; 86(2):420-428.
- 356 [23] McGraw KO, Wong SP. Forming inferences about some intraclass correlation  
357 coefficients. *Psychol Methods.* 1996; 1(1):30-46.
- 358 [24] ACSM. *Acsm's Guidelines for Exercise Testing and Prescription.* 8th ed. Baltimore:  
359 Lippinkott Williams & Wilkins; 2010.
- 360 [25] Bechler J, Jobe F, Pink M, Perry J, Ruwe P. Electromyographic analysis of the hip  
361 and knee during the golf swing. *Clin J Sport Med.* 1995; 5(3):162-166.
- 362 [26] Bulbulian R, Ball K, Seaman D. The short golf backswing: effects on performance  
363 and spinal health implications. *J Manipulative Physiol Ther.* 2001; 24(9):569-575.
- 364 [27] Theriault G, Lachance P. Golf injuries. An overview. *Sports Med.* 1998; 26(1): 43-  
365 57.
- 366 [28] McCarroll JR. The frequency of golf injuries. *Clin Sports Med,* 1996; 15(1): 1-7.
- 367 [29] Metz JP. Managing Golf Injuries. Technique and equipment changes to aid  
368 treatment. *Phys Sportsmed.* 1999; 27(7): 41-56.
- 369 [30] Rahimi SY, Singh H, Yeh DJ, Shaver EG, Flannery AM, Lee MR. Golf associated  
370 head injury in the pediatric population: a common sports injury. *J Neurosurg.*  
371 2005; 102(2 Suppl): 163-166.

372 [31] Watson DS, Mehan TJ, Smith GA, McKenzie LB. Golf cart-related injuries in the  
373 U.S. Am J Prev Med. 2008; 35(1): 55-59.

374 [32] Finch C, Sherman C, James T, Farrally M, Cochran A. The epidemiology of golf  
375 injuries in Victoria, Australia: evidence from sports medicine clinics and  
376 emergency department presentations. In: M. Farrally & A. Cochran, editors.  
377 Science and Golf III: Proceedings of the World Scientific Congress of Golf.  
378 Champaign, IL.: Human Kinetics; 1999. pp. 73-82.

379 [33] McHardy A, Pollard H, Luo K. Golf injuries: a review of the literature. Sports Med.  
380 2006; 36(2):171-87.

381 [34] Sugaya H, Tsuchiya A, Moriya H, Morgan DA, Banks SA. Low back injury in elite  
382 and professional golfers: An epidemiologic and radiographic study. In: M. Farrally  
383 & A. Cochran, editors. Science and Golf III: Proceedings of the World Scientific  
384 Congress of Golf. Champaign, IL.: Human Kinetics; 1999. pp. 83-91.

385 [35] Vad VB, Bhat AL, Basrai D, Gebeh A, Aspergren DD, Andrews JR. Low back pain  
386 in professional golfers: the role of associated hip and low back range-of motion  
387 deficits. Am J Sports Med. 2004; 32(2): 494-497.

388

389

390

391

## TABLES

**Table 1 – Subject Characteristics**

	<b>N</b>	<b>Age (years)</b>	<b>Height (m)</b>	<b>Mass (kg)</b>	<b>Experience (years)</b>
<b>Male</b>	56	42.4±14.76	1.75±0.1	78.6±10.6	11.1±6.1
<b>Female</b>	5	34.6±19.9	1.66±0.1	65.6±7.6	8.2±4.7
<b>Total</b>	61	41.8±15.2	1.74±0.1	77.6±11.0	10.9±6.0

**Table 2 – Reliability of Past Medical History Items**

	Frequency (occ1)		Frequency (occ2)		<i>Kappa</i>
	Yes	No	Yes	No	
	(%)	(%)	(%)	(%)	
<b>Heart Condition</b>	-	100	-	100	-
<b>Chest Pain with Activity</b>	-	100	-	100	-
<b>Chest Pain at Rest</b>	-	100	-	100	-
<b>Loss of Balance or Loss of Consciousness</b>	5.1	94.9	10	90.0	0.733
<b>Orthopaedic Problem Aggravated by Activity</b>	16.7	83.3	16.4	83.6	0.880
<b>Ingesting Blood Pressure or Heart Medication</b>	13.1	86.9	11.5	88.5	0.924
<b>Known Contraindication to Activity</b>	3.3	96.7	1.6	98.4	0.659

**Table 3 – Reliability of Subject Skill Items**

	Mean±SD		ICC	r
	(occ1)	(occ2)		
<b>Years Played</b>	10.93±6.08	10.89±6.13	0.998	0.998
<b>Driver Range</b>	197.46±28.89	196.39±31.09	0.837	0.838
<b>7-Iron Range</b>	119.34±21.46	120.45±21.42	0.970	0.971
<b>Handicap</b>	17.54±9.25	17.62±9.24	0.997	0.997

  

	Frequency (%)		Frequency (%)		Kappa
	Yes	No	Yes	No	
<b>Fitting</b>	36.1	63.9	32.8	67.2	0.855
<b>Handed</b>	-	100	-	100	-
<b>Swing Side</b>	98.4	1.6	98.4	1.6	1.0

**Table 4 – Reliability of Transportation on Golf Course Items by proportion of rounds**

	<b>Mean±SD</b> <b>(occ1)</b>	<b>Mean±SD</b> <b>(occ2)</b>	<b>ICC</b>	<b>r</b>
<b>Using Power Cart</b>	20.08±28.93	20.25±28.92	0.992	0.992
<b>Carrying Clubs</b>	42.13±40.32	44.18±40.77	0.943	0.944
<b>Pulling Clubs</b>	29.84±32.54	31.48±32.79	0.952	0.952
<b>Pushing Clubs</b>	8.55±18.98	10.74±22.41	0.705	0.714
<b>Using Electric Cart</b>	12.46±29.41	11.15±27.51	0.946	0.948

**Table 5 – Reliability of Frequency of Golf Activity Items**

	<b>Mean±SD (occ1)</b>	<b>Mean±SD (occ2)</b>	<b>ICC</b>	<b>r</b>
<b>Rounds Played</b>				
<b>Early Season</b>	5.60±4.76	5.95±5.49	0.914	0.924
<b>Mid-Season</b>	7.33±5.86	8.20±7.26	0.881	0.907
<b>Late Season</b>	5.64±4.53	6.55±6.18	0.773	0.819
<b>Off Season</b>	6.29±6.67	6.31±6.79	0.863	0.862
<b>Practice Sessions</b>				
<b>Early Season</b>	4.71±5.36	4.67±4.7	0.888	0.894
<b>Mid-Season</b>	5.42±8.42	5.16±5.42	0.796	0.872
<b>Late Season</b>	4.98±5.52	4.70±4.84	0.917	0.925
<b>Off Season</b>	5.14±6.83	4.95±4.81	0.761	0.805
<b>Putting Freq.</b>				
<b>Early Season</b>	3.55±4.91	3.40±3.67	0.901	0.939
<b>Mid-Season</b>	4.09±8.12	3.84±4.58	0.717	0.834
<b>Late Season</b>	3.67±5.02	3.48±4.09	0.920	0.939
<b>Off Season</b>	4.60±7.60	3.76±4.33	0.775	0.906
<b>Class Sessions</b>				
<b>Early Season</b>	1.51±2.89	1.24±1.38	0.634	0.816
<b>Mid-Season</b>	1.52±3.62	1.48±2.59	0.903	0.952
<b>Late Season</b>	1.55±2.85	1.28±1.57	0.708	0.837
<b>Off Season</b>	1.76±4.50	1.20±1.34	0.456	0.836



**Table 6 – Reliability of Warm-up and Exercise Habits**

	<b>Mean±SD (occ1)</b>	<b>Mean±SD (occ2)</b>	<b>ICC</b>	<b>r</b>	
<b>Warm-up</b>	9.15±5.82	9.02±4.66	0.793	0.810	
<b>Stretching</b>	3.95±3.61	3.64±3.98	0.832	0.849	
	<b>Frequency (%)</b>		<b>Frequency (%)</b>		<b>Kappa</b>
	<b>Yes</b>	<b>No</b>	<b>Yes</b>	<b>No</b>	
<b>Stretching During Round</b>	24.6	75.4	27.9	72.1	0.746
<b>Stretching Away from Course</b>	39.3	60.7	39.3	60.7	0.794
<b>Strengthening Program</b>	48.3	51.7	52.5	47.5	0.797
<b>Endurance Program</b>	61.7	38.3	61.0	39.0	0.964

**Table 7 – Reliability of Injury Items**

	Mean±SD (occ1)	Mean±SD (occ2)	ICC	r	
<b>Stopped Playing because of Injury</b>	24.55±9.34	27.27±13.30	0.672	0.714	
<b>Low Back Pain after 18 holes</b>	17.54±20.81	18.64±24.91	0.905	0.919	
	Frequency (%)		Frequency (%)		<i>Kappa</i>
	Yes	No	Yes	No	
<b>Injury incidence</b>	21.7	78.3	25.4	74.6	0.906
<b>Musculoskeletal Injury</b>	23.3	76.7	23.3	76.7	0.814

## APPENDIX

### Original Version

#### UNIVERSITY OF WESTERN ONTARIO

#### Musculoskeletal Conditions in Senior Golfers Questionnaire

#### First, we would like to ask you a few questions about yourself:

1. What is your date of birth? \_\_\_\_\_  
(day) (month) (year)
2. What is your height? \_\_\_\_\_
3. What is your weight? \_\_\_\_\_
4. Are you male or female? (circle the number of your answer)
  1. male
  2. female
5. How many years have you been playing golf? \_\_\_\_\_
6. Are you right or left handed? (circle)
  1. right
  2. left

#### Now we would like to ask you about your history of illness and injuries:

7. Has your doctor ever said that you have a heart condition and that you should only do physical activity recommended by a doctor? (circle)
  1. yes
  2. no
8. Do you feel pain in your chest when you do physical activity? (circle)
  1. yes
  2. no
9. In the past month, have you had chest pain when you were not doing physical activity? (circle)
  1. yes
  2. no
10. Do you ever lose your balance because of dizziness or do you ever lose consciousness? (circle)
  1. yes
  2. no
11. Do you have a bone or joint problem that could be made worse by changes in physical activity?  
(circle)
  1. yes
  2. no
12. Is your doctor currently prescribing drugs (for example water pills) for your blood pressure or heart condition? (circle)
  1. yes
  2. no
13. Do you know of any other reason why you should not do physical activity? (circle)
  1. yes
  2. no

If yes, please specify:

#### Next, we would like to ask a few questions about your golf game:

14. On average, how many yards do you hit your driver? \_\_\_\_\_
15. On average, how many yards do you hit your 7 iron? \_\_\_\_\_
16. What is your golf handicap? \_\_\_\_\_  
(Please give an approximate handicap if you do not have an official one)

**Now we would like to ask you about your golf swing:**

17. Do you swing your golf club left or right? (circle)
- 1. left
  - 2. right
18. Are your golf clubs customized to fit your golf swing? (circle)
- 1. yes
  - 2. no

**We would like to ask you how you get around the golf course:**

19. On average, how often do you use a power cart around the course? (circle)
- 0% of the time ...15% ..... 30% ..... 50% ..... 65% ..... 80% ..... 100% of the time**
20. On average, how often do you carry your clubs around the course? (circle)
- 0% of the time ...15% ..... 30% ..... 50% ..... 65% ..... 80% ..... 100% of the time**
21. On which side of your body do you carry your clubs? (circle)
- 1. left
  - 2. right
  - 3. alternate
  - 4. double strap

22. On average, how often do you pull your clubs around the course on a cart? (circle)
- 0% of the time ...15% ..... 30% ..... 50% ..... 65% ..... 80% ..... 100% of the time**
23. On average, how often do you push your clubs around the course on a cart? (circle)
- 0% of the time ...15% ..... 30% ..... 50% ..... 65% ..... 80% ..... 100% of the time**
24. On average, how often do you use an electronic cart to carry your clubs? (circle)
- 0% of the time ...15% ..... 30% ..... 50% ..... 65% ..... 80% ..... 100% of the time**

**We would like to know how much golf you play.**

25. On average, how many rounds of golf do you play *in a single month* during the following times:
- Early Season (April - May) \_\_\_\_\_
- Mid Season (June - August) \_\_\_\_\_
- Late Season (September - October) \_\_\_\_\_
- Off Season (November - March) \_\_\_\_\_
26. On average, how many times *in a single month* do you go to the practice range during the following times: Early Season (April - May) \_\_\_\_\_
- Mid Season (June - August) \_\_\_\_\_
- Late Season (September - October) \_\_\_\_\_
- Off Season (November - March) \_\_\_\_\_
27. On average, how many times *in a single month* do you practice putting?
- Early Season (April - May) \_\_\_\_\_
- Mid Season (June - August) \_\_\_\_\_
- Late Season (September - October) \_\_\_\_\_
- Off Season (November - March) \_\_\_\_\_
28. On average, how many times *in a single month* do you take lessons from a golf professional?
- Early Season (April - May) \_\_\_\_\_
- Mid Season (June - August) \_\_\_\_\_
- Late Season (September - October) \_\_\_\_\_
- Off Season (November - March) \_\_\_\_\_

**We would like to ask you about your warm-up.**

29. On average, how much time do you spend warming up prior to playing or practicing? \_\_\_\_\_ minutes
30. How much of this warm up time is spent stretching? \_\_\_\_\_ minutes

31. Once you've started a round, do you routinely perform any golf stretches while out on the course? (circle)

- 1. yes
- 2. no

**Now, just a few questions about other exercises you might do.**

32. Do you routinely perform any of your golf stretches away from the course/practice range? (circle)

- 1. yes
- 2. no

33. Do you routinely do any strengthening exercises? (circle)

- 1. yes
- 2. no

34. Do you routinely participate in a cardiovascular conditioning program apart from golfing? (circle)

- 1. yes
- 2. no

**Finally, we would like to ask about your golfing injuries.**

35. Have you suffered **ANY** injuries in the past 3 years while playing or practicing golf, which caused you to stop or modify your game for at least 2 weeks? (circle)

- 1. yes
- 2. no

if yes, please tell us which part(s) and side of your body was hurt

---

and the medical name or diagnosis of each injury, if you know it (e.g., tennis elbow, low back strain)

---

36. Typically, how often are you aware of low back pain after golfing 18 holes? (circle)

**0% of the time ...15% ..... 30% ..... 50% ..... 65% ..... 80%..... 100% of the time**

37. Have you suffered **ANY** muscle or joint conditions in the past 3 years which affected your golf game? (circle)

- 1. yes
- 2. no

if yes, please tell us which part(s) and side of your body was hurt  
and the medical name or diagnosis of each condition, if you know it

---

***Thank you very much for completing this questionnaire***

## Portuguese version

### Universidade Técnica de Lisboa Faculdade de Motricidade Humana

#### Questionário sobre condições músculo-esqueléticas em jogadores de golfe adultos (Fox, Lindsay & Vandervoort, 2002)

Em primeiro lugar, gostaríamos de lhe fazer algumas perguntas sobre si:

1. Qual é a sua data de nascimento? \_\_\_\_\_  
(dia) (mês) (ano)
2. Qual é a sua altura? \_\_\_\_\_ cm
3. Qual é o seu peso? \_\_\_\_\_ kg
4. Sexo masculino ou feminino? (deve escolher a opção adequada assinalando-a com um círculo no número correspondente à sua resposta)
  1. Masculino
  2. Feminino
5. Há quantos anos joga golfe? \_\_\_\_\_ anos/ meses (**riscar o que não interessa**)
6. É destro ou esquerdino? (Círculo)
  1. Destro
  2. Esquerdino

#### Gostaríamos agora de o questionar acerca do seu histórico de doenças e lesões:

7. O seu médico alguma vez lhe disse que tinha um problema cardíaco e que só deveria fazer actividade física recomendada por um médico? (Círculo)
  1. Sim
  2. Não
8. Sente dor no peito quando faz exercício físico? (círculo)
  1. Sim
  2. Não
9. No último mês, alguma vez teve dor no peito quando não estava a fazer actividade física? (Círculo)
  1. Sim
  2. Não
10. Alguma vez sentiu tonturas chegando a perder o equilíbrio ou alguma vez desmaiou?
  1. Sim
  2. Não
11. Tem algum problema ósseo ou articular que possa ser agravado pela prática de actividade física? (Círculo)
  1. Sim
  2. Não
12. Toma medicamentos (por exemplo, diuréticos) para a tensão arterial ou para problemas cardíacos? (Círculo)
  1. Sim
  2. Não
13. Conhece algum outro motivo que o possa impedir de fazer exercício físico? (círculo)
  1. Sim
  2. Não

Em caso afirmativo, especifique, por favor: \_\_\_\_\_

**De seguida, gostaríamos de fazer algumas perguntas sobre o seu jogo/prática de golfe:**

14. Em média, quantos metros bate com o seu driver? \_\_\_\_\_

15. Em média, quantos metros bate com o seu ferro 7? \_\_\_\_\_

16. Qual é o seu handicap? \_\_\_\_\_

(Refira o seu handicap aproximado, caso não tenha um oficial)

**Gostaríamos agora de o questionar sobre o seu swing:**

17. O seu swing é destro ou esquerdino? (Círculo)

1. Esquerdino

2. Destro

18. Os seus tacos de golfe estão adaptados (fitting) ao seu swing? (Círculo)

1. Sim

2. Não

**Gostaríamos de perguntar como se desloca no campo de golfe:**

19. Em média, com que frequência usa o *buggie* durante o percurso de jogo? (Círculo)

**0% do tempo** ..... 15% ..... 30% ..... 50% ..... 65% ..... 80% ..... **100% do tempo**

20. Em média, com que frequência transporta os seus tacos durante o percurso de jogo? (Círculo)

**0% do tempo** ..... 15% ..... 30% ..... 50% ..... 65% ..... 80% ..... **100% do tempo**

21. De que lado do corpo transporta os seus tacos? (Círculo)

1. Esquerda

2. Direito

3. Alternado

4. Duas alças

22. Em média, com que frequência puxa os seus tacos num trolley durante o percurso de jogo? (Círculo)

**0% do tempo** ..... 15% ..... 30% ..... 50% ..... 65% ..... 80% ..... **100% do tempo**

23. Em média, com que frequência empurra os seus tacos num trolley durante o percurso de jogo? (Círculo)

**0% do tempo** ..... 15% ..... 30% ..... 50% ..... 65% ..... 80% ..... **100% do tempo**

24. Em média, com que frequência usa um trolley eléctrico para transportar os seus tacos? (Círculo)

**0% do tempo** ..... 15% ..... 30% ..... 50% ..... 65% ..... 80% ..... **100% do tempo**

**Gostaríamos de saber qual a sua regularidade na prática de golfe.**

25. Em média, quantas vezes joga num só mês durante as seguintes épocas:

Época inicial (Abril-Maio) \_\_\_\_\_

Época média (Junho-Agosto) \_\_\_\_\_

Época final (Setembro-Outubro) \_\_\_\_\_

Fora de época (Novembro-Março) \_\_\_\_\_

26. Em média, quantas vezes num só mês vai para o *driving range* treinar durante as seguintes épocas:

Época inicial (Abril-Maio) \_\_\_\_\_

Época média (Junho-Agosto) \_\_\_\_\_

Época final (Setembro-Outubro) \_\_\_\_\_

Fora de época (Novembro-Março) \_\_\_\_\_

27. Em média, quantas vezes num só mês treina o putting?

Época inicial (Abril-Maio) \_\_\_\_\_

Época média (Junho-Agosto) \_\_\_\_\_

Época final (Setembro-Outubro) \_\_\_\_\_

Fora de época (Novembro-Março) \_\_\_\_\_

28. Em média, quantas vezes num só mês tem aulas com um profissional de golfe?

Época inicial (Abril-Maio) \_\_\_\_\_  
Época média (Junho-Agosto) \_\_\_\_\_  
Época final (Setembro-Outubro) \_\_\_\_\_  
Fora de época (Novembro-Março) \_\_\_\_\_

**Gostaríamos de o questionar sobre o seu aquecimento.**

29. Em média, quanto tempo dedica ao seu aquecimento antes de jogar ou praticar? \_\_\_\_\_ minutos
30. Quanto tempo do aquecimento dedica a alongamentos? \_\_\_\_\_ minutos
31. Uma vez iniciada uma volta, costuma realizar alguns alongamentos específicos para o golfe? (círculo)
1. Sim
  2. Não

**Agora, apenas algumas perguntas sobre outros exercícios que possa fazer.**

32. Por rotina, executa alguns dos seus alongamentos específicos para o golfe quando não está a praticar golfe? (círculo)
1. Sim
  2. Não
33. Costuma realizar exercícios de força? (círculo)
1. Sim
  2. Não
34. Costuma realizar treino cardiovascular, para além do golfe? (círculo)
1. Sim
  2. Não

**Finalmente, gostaríamos de lhe perguntar sobre as lesões sofridas durante a prática de golfe.**

35. Sofreu ALGUMA lesão nos últimos 3 anos, durante o jogo ou prática de golfe, que o tenha levado a parar ou modificar o seu jogo? (círculo)
1. Sim \_\_\_\_\_ (refira o tempo de paragem) \_\_\_\_\_ (dias)
  2. Não

em caso afirmativo, diga-nos qual/quais a(s) parte(s) e lado do seu corpo ficou lesionada

\_\_\_\_\_

e a designação clínica ou diagnóstico de cada lesão, se souber (ex., "cotovelo de tenista", desconforto lombar)

\_\_\_\_\_

36. Por norma, com que frequência sente dor lombar depois de fazer 18 buracos? (círculo)

0% do tempo ... 15% ..... 30% ..... 50% ..... 65% ..... 80%..... 100% do tempo

37. Sofreu ALGUM problema muscular ou articular nos últimos 3 anos que tenha afectado a sua prática de golfe? (círculo)
1. Sim
  2. Não

em caso afirmativo, diga-nos qual/quais a(s) parte(s) e lado do seu corpo ficou afectado(s)

\_\_\_\_\_

e a designação clínica ou diagnóstico de cada problema, se souber.

\_\_\_\_\_

**Muito obrigado por ter respondido a este questionário**