

Steffen, K., Andersen, T. E., Krosshaug, T., van Mechelen, W., Myklebust, G., Verhagen, E. ALM, Bahr, R. (2010). ECSS Position Statement 2009: Prevention of acute sports injuries. *European Journal of Sport Science*, 10, 223-236.

Dette er siste tekst-versjon av artikkelen, og den kan inneholde ubetydelige forskjeller fra forlagets pdf-versjon. Forlagets pdf-versjon finner du på www.informaworld.com: <http://dx.doi.org/10.1080/1746139090358517>

This is the final text version of the article, and it may contain insignificant differences from the journal's pdf version. The original publication is available at www.informaworld.com: <http://dx.doi.org/10.1080/1746139090358517>

ECSS Position Statement 2009: Prevention of acute sports injuries

K. Steffen¹, T.E. Andersen¹, T. Krosshaug¹, W. van Mechelen², G. Myklebust¹, E. Verhagen², R. Bahr¹

¹Oslo Sports Trauma Research Center, Department of Sports Medicine, Norwegian School of Sport Sciences, Oslo, Norway

²Department of Public and Occupational Health, EMGO Institute, VU University Medical Center, Amsterdam, Netherlands

Running title: Prevention of acute sports injuries

Key words: athletic injury, prevention, risk factors, injury mechanisms, safety equipment, exercise programs, health policy

Correspondence to:

Kathrin Steffen, Oslo Sports Trauma Research Center, Department of Sports Medicine, Norwegian School of Sport Sciences, PO Box 4014 Ullevål Stadion, 0806 Oslo, Norway.

Fax: +47 23 26 23 07

e-mail: kathrin.steffen@nih.no

Abstract

To maximize the health benefits of sports and exercise and minimize the direct and indirect costs associated with injuries, developing and adopting injury prevention strategies is an important goal. The aim of this ECSS consensus paper on injury prevention is to review current evidence on injury prevention methods and training programs aimed at reducing the most common or severe types of acute injuries. The target audience is everyone involved in protecting the health of the athlete, i.e. coaches, referees, medical staff, sports governing bodies, as well as athletes themselves. Effective sports injury prevention requires successful implementation of efficacious interventions. This paper reviews the main mechanisms and risk factors for acute injuries to the head, shoulder, elbow, hand/wrist, groin, thigh, knee and ankle, as well as the evidence supporting various strategies to prevent them. Approaches that have been proven successful include: 1) using equipment designed to reduce injury risk, 2) adopting the rules of play, and 3) specific exercise programs developed to reduce injury risk. Sports organizations should adopt available injury prevention strategies as part of their policies.

Introduction

The physical activity guidelines of the American College of Sports Medicine recommend adults to achieve 20–30 min of vigorous exercise at least five days a week for optimum functional capacity and health (Haskell et al., 2007). Numerous health benefits of physical activity have been well documented, resulting in public health support for regular physical activity and exercise. Regular physical activity reduces the risk of premature mortality in general, and of coronary heart disease, hypertension, colon cancer, obesity, and diabetes mellitus in particular (Haskell et al., 2007; LaMonte et al., 2005).

The question is whether these health benefits outweigh the risk of potential injury and long-term disability associated with sports participation, especially at the elite level (Engebretsen & Bahr, 2009). A Finnish study on the incidence of chronic disease and life expectancy of former elite level athletes, revealed that the rate of hospitalization for heart disease, respiratory disease and cancer was lower for former athletes than for a control group of non-athletes. However, former athletes were more likely to have been hospitalized for musculoskeletal disorders (Kujala et al., 1996), and a follow-up study showed that former team sport athletes had an increased risk of knee osteoarthritis (Kettunen et al., 2001). Thus, although evidence suggests that sports participation is beneficial from a public health perspective, injuries also have significant negative side effects on both the short and long term.

Some injury types are of particular concern, either because they can be severe, such as head and knee injuries, or common, such as ankle sprain or hamstring strain injuries. These represent a considerable problem for the athlete, the team, and, given the popularity of sports, for the society at large. Thus, to maximize the health benefits of sports and exercise and to minimize the direct

and indirect costs associated with injury, developing methods to prevent sports injuries is a significant goal. However, it should be kept in mind that the risk of injury will to a certain extent always remain when participating in sports.

In order to develop effective injury prevention strategies it is critical to know the causes of injuries. As described by Meeuwisse et al. (2007), one important goal is to map the intrinsic and extrinsic factors that put an athlete at risk of injury. In addition, a precise description of the inciting event, the injury mechanism, is necessary (Bahr & Krosshaug, 2005).

The aim of this ECSS consensus paper on injury prevention is to review current evidence on injury prevention strategies and training programs aimed at reducing the most common or severe types of acute injuries. To better understand the rationale of effective injury prevention measures, each section also briefly describes the most important injury characteristics, risk factors, and mechanisms. General recommendations on the implementation of injury prevention strategies are presented in the final section of this paper.

Methods

Individuals with extensive experience in prevention research related to each of the body regions to be covered were identified. Each of them was asked to write a section outlining the current status regarding the characteristics, risk factors and mechanisms of injury to a specific region/injury type, as well as methods to prevent the injury in question. We did not require them to do a formal search of the literature, although most did to supplement their own literature database. Their draft was circulated among the group members until a final version was reached, which was acceptable to all members.

Head

Injury characteristics: Head and neck injuries are common across many sports. Horse riding, ice hockey, skiing/snowboarding, soccer and other football codes (e.g. American and Australian football) are sports where head injuries can result from a fall or from direct contact with sports equipment or other athletes, either by chance or through poor individual skills or rule violations. Recent reviews (Ackery et al., 2007; Hagel, 2005) clearly show that a head injury is the most frequent reason for hospital admission and the most common cause of death among skiers and snowboarders. Fortunately, serious head injuries are rare. The majority (90%) of head injuries are minor, defined as mild concussions (Straume-Næsheim et al., 2005). Concussions typically result in rapid but short-lived impairment of neurological function that resolves spontaneously.

Although most athletes with head injuries recover uneventfully following a single concussive episode, repetitive mild head trauma may cause cognitive impairment (Straume-Næsheim et al., 2005; Kirkendall et al., 2001). However, computerized neuropsychological testing did not reveal any impairment from heading exposure or previous concussions in a cohort of professional soccer players (Straume-Næsheim et al., 2005).

Risk factors: Several risk factors for concussive injuries have been suggested. There is agreement in the skiing and snowboarding literature that beginners have an increased injury risk in general and a higher risk for head injuries in particular (Ackery et al., 2007; Hagel, 2005). In snowboarding, for instance, beginners easier tend to lose their balance, which can result in a backwards fall with an impact to the back of the head. Snow parks with half pipes, rails and other slope modifications are popular and may contribute to an increased head injury risk as individual ability is often overestimated (Ackery et al., 2007; Hagel, 2005). The risk factors for catastrophic head and neck injuries are not known.

Mechanisms: Head injuries in sports arise mainly due to impacts either with an opposing player, the ground/surface, or equipment. In soccer, a blow of the elbow or arm to the opponents head, as well as a head-to-head contact are the main mechanisms for concussions (Fuller et al., 2005; Andersen et al., 2004a). In skiing/snowboarding, concussions mainly result from a fall or landing with a surface head crash, while even more severe head injuries may follow from collisions with other skiers or objects, e.g. trees, rocks, or lift towers (Hagel, 2005).

Prevention: Two main approaches have been used to prevent head injuries; using a helmet and rule modifications.

Helmets or padded headgear are used in many high-energy and collision sports to prevent head injury. A meta-analysis, based on five well-conducted case-control studies on pedal cycle helmets in transport and recreation, have confirmed that helmets provide a 63-88% reduction in the risk of head, brain and severe brain injury for all ages of bicyclists (Thompson et al., 2000). In recreational skiing and snowboarding, there is solid evidence showing that using a helmet is associated with a 22-60% reduction in head injury rates (Sulheim et al., 2006; Hagel, 2005; Macnab et al., 2002). Thus, it is strongly recommended that recreational skiers and snowboarders wear helmets. There are no data from competitive cycling, skiing or snowboarding, but helmets are compulsory in the competition regulations for these and several other sports. Protective equestrian helmets are also widely recommended. Interestingly, since helmets were made compulsory for professional jockeys in 1993-94, no significant changes in concussion rates have been observed (McCrorry & Turner, 2005). However, research on padded headgear (soft shell helmets) indicates that they do not reduce the incidence of a concussion or serious head injuries in Rugby Union football (McIntosh et al., 2009), and similarly, data from soccer (Andersen et al.,

2004a) and Australian rules football (McIntosh & McCrory, 2005) suggest that the currently available head gears are unlikely to reduce concussion incidences.

Rule modifications represent one strategy to reduce the risk of collisions and head impacts in sports, albeit direct scientific evidence is lacking. Studies from elite soccer show that approximately 40% of all concussions can be attributed to elbow and arm impacts in heading duels (Fuller et al., 2005; Andersen et al., 2004a). As a result, FIFA introduced a stricter rule interpretation prior to the 2006 World Cup making contact to an opponent's head with the elbow/upper arm during heading duels a 'red card' offence. In ice hockey, injury prevention programs are becoming effective in reducing the number of injuries, especially head, cervical spine and spinal cord injuries when banning checking from behind (McIntosh & McCrory, 2005).

Shoulder

Injury characteristics: Acute shoulder injuries are typical in sports which involve powerful and intentional body contact, like certain football codes, or ice hockey, but they are also common in sports which are characterized by high-energy falls on the shoulder, like in skiing, snowboarding, ice skating, and cycling.

The severity of an acute shoulder injury depends on the direction of the forces and the anatomic structure affected. Most acute shoulder injuries are minor, resulting in soft tissue contusions around the shoulder. More severe trauma may result in a shoulder dislocation, clavicle fracture, a acromioclavicular and sternoclavicular joint dislocation or a fracture of the upper humerus (Krogsgaard et al., 2009).

Risk factors: There is no evidence on risk factors associated with acute shoulder injuries.

Mechanisms: An acute shoulder injury occurs either directly by falls or blows on the shoulder, or indirectly caused by transmission of forces through the arm.

Prevention: There is no evidence in the literature for methods to protect an athlete from acute shoulder injuries. However, any measures to reduce the risk of falls in sports should reduce the risk of shoulder injury. Shoulder pads can theoretically spread the forces from an impact over a larger area of the body and absorb energy, just like a helmet. In American football, shoulder pads protect the shoulder by dissipating the forces. There are, however, no studies that have proven the effect of shoulder pads on shoulder injury risk reduction. In addition, hard shoulder pads have the disadvantage that they can hurt other players, particularly if they are used to butt others (Krogsgaard et al., 2009). Proper falling techniques can be taught to potentially lessen the direct impact placed across the shoulder joint by learning to roll when landing from a fall, rather than landing directly on the shoulder. Head-first and diveback sliding techniques in baseball and softball can result in upper extremity injuries. However, changing techniques, e.g. banning sliding or forbidding head-first/diveback sliding is impractical and unsatisfactory to players, and instructional courses regarding injury prevention have proved ineffective since players did not attend (Janda, 2003). Even if direct evidence from team handball is lacking, the adoption of strict penalties for tackling the attacker's arm from behind or holding the player's arm when shooting, has been assumed to prevent acute lesions to the shoulder joint. To sum up, there is not much evidence on either risk factors or mechanisms for acute shoulder injuries, and, hence, either for effective preventive strategies.

Elbow

Injury characteristics: Acute elbow injuries are most common in 1) contact and collision sports, e.g. American football, rugby, and martial arts, 2) sports that place an athlete at elevated heights,

e.g. high jump, ski jumping, and gymnastics, and 3) high-speed sports, e.g. alpine skiing, speed skating and cycling. Also sports such as weightlifting, boxing, shot put and javelin place acute and heavy loads across the elbow joint during training and competition and expose the elbow to a potential injury risk. Athletes who fall on an extended elbow with the hand in full supination can suffer a number of injury types, including bone bruises, ligament strains, muscle contusions and skin abrasions, and in the worst cases fractures and dislocations, even with injury to neurovascular structures (Hutchinson & Andrews, 2009).

Risk factors: There is no evidence on risk factors associated with acute elbow injuries.

Mechanisms: An acute elbow injury may either occur by direct impact onto the elbow, e.g. by a fall or a collision, or by indirect impact resulting in torques across the elbow, as by a fall on an outstretched hand; the classical injury description. When the elbow is hyperextended and when moments and forces are excessive, the collateral ligaments will rupture and damage the anterior capsule. If the forces continue, the elbow will dislocate. In judo for instance, an athlete is allowed to place the opponent's elbow in an elbow lock, a position of hyperextension. Indeed, if the athlete on the receiving end of this maneuver does not "tap-out" to concede defeat, continued force may lead to an elbow dislocation (Hutchinson & Andrews, 2009).

Prevention: There is little research on injury prevention methods, although it has been suggested that elbow injuries may be prevented through protective padding and education of athletes on how to fall properly (Hutchinson & Andrews, 2009). Minimizing risk exposures and energy transfer during falls could be accomplished via rule enforcement, more severe penalties for violations, or rule changes. An example that has been shown to be effective was limiting the height of cheerleader towers which, in turn, reduced the potential energy and risk of injury from falls (Boden et al., 2003). In judo, banning the ability of a competitor to hyperextend the elbow

and subsequently dislocate it until his opponent taps out (gives up) would likely reduce the risk of elbow injuries in judo sport (Hutchinson & Andrews, 2009). Unfortunately, similar to traumatic shoulder injuries, there is not much evidence on how to prevent acute shoulder injuries. However, as for acute shoulder injuries, many acute elbow injuries are simply accidents occurring during the usual performance of a high-risk sport.

Hand/Wrist

Injury characteristics: Sprains of finger and wrist joints as well as fractures of the wrist and fingers are typical for ball sports and other sports involving falling activities/accidents. Wrist fractures are common in snowboard sports (Hagel, 2005; Langran and Selvaraj, 2004). Wrist injuries account for 8% of injuries among elite snowboarders (Torjussen & Bahr, 2006; Torjussen & Bahr, 2005), and up to 32% among recreational boarders (Langran & Selvaraj, 2004; Machold et al., 2002). In less serious cases, falls result in sprains of the wrist joint (Hagel, 2005). ‘Skier’s thumb’ is a tear of the ulnar collateral ligament of the first metacarpophalangeal joint, leading to instability of the joint (Mogan & Davis, 1982), while volleyball players can tear the radial collateral ligament of the same joint when playing defense (‘volleyball thumb’) (Bahr & Reeser, 2003). Most other finger injuries in e.g. volleyball and team handball are mild ligament injuries with which the athlete continues to train or compete, with or without exposed fingers taped.

Risk factors: Finger sprains take time to heal fully, and a history of a previous finger sprain exposes this finger to new injuries. Novice snowboarders are 2-3 times more likely to injure their wrist than experienced boarders (Goulet et al., 2009; Langran & Selvaraj, 2004; Machold et al., 2002). The popularity of snow parks with half pipes, rails and jumps, contribute to an increased risk of wrist sprains and fractures as a result of poor or overestimated physical and technical ability (Hagel, 2005).

Mechanisms: Complicated fractures of the wrist or forearm may occur as a result of direct falls or of unsuccessful jumps, typically seen in skiers and snowboarders. When balance is lost, snowboarders, unlike skiers, cannot ‘step out’ a leg in order to recover the lost balance, as both feet are firmly attached to the board. The instinctive protective reaction is to reach out to break the landing, thus placing the wrist and fingers at risk for injury (Machold et al., 2002). The typical mechanism for skier’s thumb is when a skier falls with the ski pole left in the hand forcing the thumb into adduction and extension (Browne et al., 1976). The mechanism of finger sprains in volleyball and team handball are similar; hyperextension when the ball unexpectedly hits the finger during defense or blocking.

Prevention: There is agreement in the literature that most of the wrist fractures in snowboarding can be prevented by use of wrist protectors, at least among beginners (Russell et al., 2007; Rønning et al., 2001). Wrist protectors are, beside helmet use, one of the simplest strategies to prevent injuries in snowboarding. For every 50 snowboarders who wear a wrist guard, one wrist injury will be averted (Russell et al., 2007). However, the optimal type of wrist guard to most effectively prevent wrist injuries is not defined. There is no conclusive evidence whether or not wrist guards increase the risk of shoulder, shoulder girdle, or elbow injuries (Russell et al., 2007; Hagel, 2005). For skiers, strapless poles have not shown to decrease the incidence of skier's thumb. However, if skiers are trained to discard the pole during a fall, injury risk might be reduced (Fricker & Hintermann, 1995).

One obvious way to try and reduce the risk of injuries to the upper limb in general and the wrist joint in particular, is proper falling techniques and avoiding to land on an outstretched hand.

Finger sprains among team sport players may be prevented by prophylactic taping of the exposed fingers, especially fingers with previous ligamentous injuries.

Groin

Injury characteristics: The pelvis is the center of load transfer from the upper extremities and the torso to the lower extremities and visa versa, and a number of muscle groups interact to stabilize the region and facilitate energy transfer around the pelvic and hip region, including the adductors, iliopsoas and abdominal and erector spinae muscles. Groin injuries are among the top six most cited injuries in e.g. soccer, rugby, American football, ice hockey, speed skating, swimming and athletics accounting for >10% of all injuries, and 20-40% of all muscle strain injuries (Maffey & Emery, 2007). A strain to the groin muscles may be acute but often becomes chronic in nature. Suffering from persistent groin injuries results in extensive rehabilitation time and longstanding pain (Jansen et al., 2008; Macintyre et al., 2006); as an example, 40% of groin injuries in soccer result in more than one week time loss, and 10% in more than one month (Maffey & Emery, 2007).

Risk factors: There is clear evidence that a previous strain of the groin muscles on the same side is a strong predictor for a recurrent injury (Steffen et al., 2008a; Hägglund et al., 2006; Emery et al., 2005). This may be due to scar tissue formation in the muscle or tendon or inadequately rehabilitated strength or flexibility (Steffen et al., 2008a; Emery et al., 2005; Árnason et al., 2004). Other intrinsic risk factors believed to be involved in groin injuries are reduced adductor strength and flexibility of the hip abductors (Maffey & Emery, 2007; Verrall et al., 2005; Árnason et al., 2004). Decreased adductor muscle strength and an imbalanced adductor-to-abductor muscle strength ratio have been shown to predict groin strain injuries. Ice hockey players were 17 times more likely to sustain an adductor muscle strain if adductor strength was less than 80% of abductor strength (Tyler et al., 2001). Reduced flexibility of the abductor and adductor muscles has been suggested as a risk factor for groin strain injuries (Verrall et al., 2005;

Árnason et al., 2004; Ekstrand & Gillquist, 1983). However, the evidence is conflicting (Maffey & Emery, 2007), as it is for age and sports exposure as risk factors for groin injuries (Maffey & Emery, 2007; Árnason et al., 2004).

Mechanisms: As groin injuries often show a transition into an overuse condition, the exact moment of injury and the injury mechanisms are hard to establish. Common for athletes developing groin problems is that they are engaged in sports involving kicking, rapid changes of direction, accelerations and decelerations (Macintyre et al., 2006). The adductor muscles may acutely be strained during an eccentric contraction, e.g. in a forced abduction, most likely when the limb is in abduction. This could be a sudden resistance caused by an opponent's foot in an attempt to reach a ball or a sliding tackle in soccer, or the eccentric force of the adductor muscles attempting to decelerate the leg during a stride in ice hockey. The adductor muscles can also be strained in a forceful concentric contraction, as during a kick after a ball in the air. The iliopsoas muscle can be acutely strained if a forceful hip flexion is interrupted suddenly, e.g. if tackled or hitting the ground when kicking, but also during an eccentric contraction (Meeuwisse et al., 2007; Maffey & Emery, 2007; Árnason et al., 2004).

Prevention: A previous groin injury is the best established risk factor, emphasizing the need for adequate rehabilitation of the primary groin injury before returning to sports (Steffen et al., 2008a; Hägglund et al., 2006; Árnason et al., 2004). It is not known whether stretching of adductor/abductor and iliopsoas muscles can prevent groin injuries. However, a functional flexibility of these muscles and range of motion of the hip joint may be important (Árnason et al., 2004).

As many groin injuries occur during eccentric loading, reduced eccentric strength of the adductor muscles has been suggested as a possible risk factor. However, studies on different athlete

populations (ice hockey, soccer) do not provide conclusive evidence for general strengthening exercises as a prevention method (Maffey & Emery, 2007). Nevertheless, an intervention study including specific strength training of the adductor muscles (static, concentric and eccentric), as well as core stability exercises for the pelvic muscles showed a highly significant effect in the treatment of adductor-related groin injuries (Hölmich et al., 1999). However, this program has not been evaluated as a primary prevention method, although such exercises are commonly included in general prevention programs for lower extremity injuries.

Thigh

Injury characteristics: Common thigh injuries are contusions to the anterior and lateral thigh and strains to the quadriceps or hamstring muscles. Hamstring strains is the most common injury type in many sports, particularly in the football codes, which are characterized by sudden accelerations and decelerations, often followed by changes of direction, and eccentric muscle activity during sprinting and kicking (Askling et al., 2008; Hägglund et al., 2006; Mjølunes et al., 2004; Árnason et al., 2004). Recent studies suggest that the proportion of hamstring strains has increased in soccer during the past decade, as the sport has become more physically demanding with more frequent and higher intensity runs (Hägglund et al., 2006; Árnason et al., 2004). Sports characterized by slow stretching beyond the usual movement of the muscles, in particular hip flexion (e.g. dancing), also have high rates of hamstring injuries (Askling et al., 2008).

Risk factors: The two factors most consistently associated with a hamstring strain injury are a history of a previous injury and age (older players with higher risk) (Árnason et al., 2004). Some studies indicate that decreased hamstring strength (concentric and eccentric), low hamstring-to-quadriceps strength ratio and a side-to-side difference in hamstring strength could be risk factors for hamstring strains (Murphy et al., 2003). Poor flexibility of the hamstring muscles has also

been suggested as a risk factor for muscular strains (Murphy et al., 2003). However, there is a paucity of evidence, and the methodology has been poor, thereby not allowing definitive conclusions to be drawn.

Mechanisms: Most hamstring strains occur during maximal sprinting activities in the late swing phase, when the hamstring muscles are loaded eccentrically to decelerate the forward movement of the leg, or at foot-strike, in the transition from eccentric to concentric muscle action (Askling et al., 2008; Árnason et al., 2008; Proske et al., 2004). It has therefore been postulated that eccentric overload could cause tearing in the muscle tendon unit (Maffey & Emery, 2007; Árnason et al., 2004; Garrett, 1990).

Prevention: Based on the injury mechanism, it has been suggested that eccentric strength is important to prevent hamstring strains. There is good evidence that eccentric strength training of the hamstring muscles increases hamstring muscle torque more effectively than concentric hamstrings training (Mjølunes et al., 2004; Askling et al., 2003) and reduces the incidence of hamstring strains in elite male soccer players (Mjølunes et al., 2004; Árnason et al., 2004; Askling et al., 2003). The Nordic Hamstring lower exercise is a partner exercise where the athlete attempts to resist a forward-falling motion of the upper body using the hamstring muscles to maximize loading in the eccentric phase. In a study on Norwegian and Icelandic male soccer players, this exercise reduced the risk of hamstring strains by 65% (Árnason et al., 2008). There is little or no evidence in the literature to support that flexibility training alone can prevent hamstring strains in soccer or other sports. One explanation may be that most hamstring strains occur during maximal sprinting, when the hamstring muscles are not stretched toward their maximum range (Askling et al., 2008; Árnason et al., 2008; Verrall et al., 2005). Eccentric hamstring training using the Nordic Hamstring exercise have been included in several general

lower extremity injury prevention programs targeting young athletes (Steffen et al., 2008b; Soligard et al., 2008; Gilchrist et al., 2008; Mandelbaum et al., 2005; Olsen et al., 2005).

Knee

Injury characteristics: Typical injuries to the knee are ligament tears of the collateral or cruciate ligaments; these may be associated with meniscal tears, or cartilage and bone damage of varying severity (Renström et al., 2008; Griffin et al., 2006). Anterior cruciate ligament (ACL) injuries are common and serious, and being associated with an increased risk for early osteoarthritis. The majority of ACL-injured athletes will develop osteoarthritis within 15-20 years, regardless of treatment (Myklebust & Bahr, 2005; von Porat et al., 2004; Lohmander et al., 2004). Athletes from pivoting sports such as soccer and other football codes, team handball, and basketball are at risk for sustaining such injuries (Hootman et al., 2007; Myklebust et al., 2003). Alpine skiing is another activity associated with high risk for knee and ACL injuries (Pujol et al., 2007).

Risk factors: As for most other injury types, recent studies have suggested that a history of knee injury is a risk factor for a subsequent knee (Steffen et al., 2008a; Hägglund et al., 2006) or ACL injury (Waldén et al., 2006; Faude et al., 2006). Female athletes are reported to have a 3-6 times higher incidence of non-contact ACL injuries than their male counterparts (Hewett et al., 2006a). However, the reasons for the gender gap are not fully elucidated. Various researchers have suggested differences in anatomy, as well as hormonal and neuromuscular function as potential reasons for the gender gap (for reviews see e.g. Shimokochi & Shultz, 2008; Renström et al., 2008; Griffin et al., 2006). However, there are only a few well-designed prospective studies available, and most of them have assessed only one risk factor in isolation. ACL injuries can most likely not be attributed to one single factor and large scale studies are needed to assess how different risk factors act in combination to put the female athlete at risk.

Thus, there is a paucity of evidence linking all these potential risk factors to non-contact ACL injuries, and a great deal of controversy exists on the importance of the different intrinsic risk factors. From the available evidence, it appears that the predictive values of static anatomic properties such as knee alignment, Q-angle, knee geometry, and leg length is limited (for reviews see e.g. Shimokochi & Shultz, 2008; Renström et al., 2008; Griffin et al., 2006). Differences in mechanical properties of the ACL may be another link to the association between ACL tensile properties and injury risk (Chandrashekar et al., 2006).

There appears to be a consensus emerging from the literature that the likelihood of incurring an ACL injury does not remain constant during the menstrual cycle, with a significantly greater risk during the preovulatory phase than during the postovulatory phase (Hewett et al., 2007).

Compared to anatomical and hormonal risk factors, neuromuscular risk factors are more important to examine in sufficient depth as these are easier modifiable by prevention training. It has been shown that females exhibit greater quadriceps dominance, and have less muscle stiffness and muscle strength. However, none of these factors have been assessed in prospective ACL injury risk factor studies and consequently, it is not possible to determine the relative importance of these factors. So far, there is only one study documenting a link between dynamic knee valgus motion and ACL injury risk (Hewett et al., 2005). However, the limited sample size (only 9 ACL ruptures) means that the results must be interpreted with caution. Deficits in neuromuscular control of the trunk has also been shown to be associated with knee injury risk (Zazulak et al., 2007a).

Investigations from Australian rules football (Orchard, 2002) and team handball (Olsen et al., 2004) suggest that a high shoe-surface friction is connected to an increased ACL injury risk.

Mechanisms: ACL injuries are commonly non-contact in nature and can occur during plant and cut maneuvers or during landings. Although the majority of ACL injuries are non-contact by definition, the movement patterns often involve perturbation by an opponent, e.g. body contact prior to the injury (Krosshaug et al., 2007a). However, even if these few details are known, we still lack vital knowledge about the injury mechanisms. A complete biomechanical description should quantify whole body and knee kinematics, loading directions, magnitudes and rates (Krosshaug et al. 2005).

The biomechanical mechanisms for non-contact ACL injuries are widely debated. The two most discussed theories involve the quadriceps drawer mechanism and valgus loading. Proponents of the first theory argue that ACL injuries result from loading in the sagittal plane only, primarily as a result of anterior shear forces caused by forceful quadriceps contraction when landing or cutting (Yu & Garrett, 2007). This mechanism was demonstrated in a cadaver study (DeMorat et al., 2004), but was criticized for not taking ground reaction forces into consideration. Others argue that knee abduction must be involved in the injury mechanism (Quatman & Hewett, 2009). This theory is partly based on video analyses of ACL-injured athletes where valgus or valgus collapse are frequently reported (Krosshaug et al., 2007a; Krosshaug et al., 2007b; Olsen et al., 2004). The combination of external rotation and valgus can possibly result in femoral notch impingement (Fung & Zhang, 2003).

Also anterior tibial translation induced by a high tibiofemoral compression and tibial plateau inclination has been proposed as possible loading mechanisms (Meyer & Haut, 2008). Internal rotation of the tibia on a relatively extended knee is suggested as an ACL injury mechanism by cadaver studies (Markolf et al., 1995), by athlete interview studies (Myklebust et al., 1998), and in video analysis (Olsen et al., 2004). However, in order to make conclusive statements on the

mechanisms of non-contact ACL injuries, new, sophisticated methods for video analysis (Krosshaug et al., 2007a) combined with other research approaches such as clinical studies and cadaver simulation or mathematical simulation studies are needed (Krosshaug et al., 2005).

The mechanisms of knee injuries in alpine skiing and snowboarding are likely different from those in team sports and at least among recreational skiers. The two main mechanisms, according to Ettliger and co-workers (1995) are the so-called “phantom foot”, i.e. internal rotation in deep flexion due to canting of the inside edge of the tail of the ski, and the ‘boot-induced anterior drawer mechanism’.

Prevention: A growing number of injury prevention programs have been developed to reduce the risk of ligamentous knee injuries in general, and ACL injuries in particular (Renström et al., 2008; Hewett et al., 2006b). They are generally based on the assumption that modifying the dynamic biomechanical risk factors can prevent injuries. Successful prevention programs attempt to alter the dynamic loading of the knee joint through neuromuscular training. Based on the likely injury mechanisms, it is recommended that athletes avoid knee valgus and land with knee flexion to absorb landing forces (Renström et al., 2008; Krosshaug et al., 2007a; Hewett, 2006b).

Training programs that incorporate plyometrics aim to result in safe levels of valgus stress to the knee and alterations in neuromuscular control patterns (Hewett et al., 2006b). As shown in several well-designed intervention studies from soccer and team handball, programs incorporating one or more exercise components to modify dynamic control of the lower limbs have resulted in remarkable reductions in the risk for ACL injuries (Gilchrist et al., 2008; Mandelbaum et al., 2005; Myklebust et al., 2003) or lower extremity injuries (Soligard et al., 2008; Olsen et al., 2005). Balance training alone, and all home-based training without instruction and feedback on proper movement technique (lower limb alignment, two-foot

landings), is probably not as effective as when it is combined with other types of neuromuscular exercise stimuli (Hewett et al., 2006b; Söderman et al., 2000).

Evidence from research on alpine skiers shows that an educational program to increase awareness of and modify behaviour in high-risk situations can prevent ACL injuries (Ettliger et al., 1995).

Ankle

Injury characteristics: An acute lateral ligament injury to the ankle (ankle sprain) represents the dominant injury type in many sports. Most ankle sprains result in an absence from training and competition for not more than a week. Nevertheless, because they are so common in many sports, ankle sprains account for a substantial part of the total injury load. In female youth soccer, for instance, up to 40% of all injuries are ankle sprains (Le Gall et al., 2008; Steffen et al., 2007; Söderman et al., 2001), and in volleyball they account for up to 50% of all reported injuries (Verhagen et al., 2004; Bahr & Bahr, 1997). From a public health perspective, ankle sprains therefore constitute a large part of the medical costs due to sports injuries (Verhagen et al., 2005).

Risk factors: The only well-documented, and probably by far strongest risk factor for an ankle sprain, is a previous sprain sustained in the last 12 months. Studies on various athlete groups found a two- to five-fold increased risk for a lateral ankle ligament injury after suffering a prior ankle injury (Steffen et al., 2008a; Verhagen et al., 2004; Árnason et al., 2004). There is strong evidence that an ankle sprain negatively affects neuromuscular control, believed to be due to trauma to mechanoreceptors of the ankle ligaments causing proprioceptive impairment (Freeman, 1965). As a result, a previously injured ankle is exposed to an increased risk of re-injury (Steffen et al., 2008a; Verhagen et al., 2004; Árnason et al., 2004). There is conflicting evidence about the injury impact of other intrinsic risk factors, such as ankle joint laxity or decreased dorsiflexion of

the ankle (Árnason et al., 2004). Anatomic foot type, when classified as pronated, supinated or neutral, does not appear to be a risk factor for ankle sprains.

Mechanisms: Common mechanisms of ankle sprains are tackling, running or landing on uneven ground or the foot of another player. Two specific injury mechanisms are reported in soccer; 1) player-to-player contact with impact by an opponent on the medial aspect of the leg just before or at foot strike, resulting in a laterally directed force causing the player to land with the ankle in a vulnerable position, and 2) forced plantar flexion where the injured player hits the opponent's foot when attempting to shoot or clear the ball (Andersen et al., 2004b; Giza et al., 2003). In volleyball, ankle sprains typically occur at the net, resulting from contact between the blocker and the opposing attacker across the centerline or when a blocker lands on a teammate's foot when participating in a multi-person block (Verhagen et al., 2004; Bahr & Bahr, 1997).

Prevention: Athletes with a history of ankle sprains represent a group which should be targeted, specifically those with sprain injuries during the preceding 12 months. Ankle proprioception is critical to keep the ankle in a safe 'neutral' position, and this ability is often impaired after injury. External support (bracing, taping) has been shown to reduce the risk of re-injury in athletes with a history of ankle sprains, but not in athletes with no previous history (Surve et al., 1994; Tropp et al., 1985). A cheaper, well investigated and highly effective prophylactic measure is balance training. Such neuromuscular training is thought to improve proprioception by re-establishing and strengthening the protective reflexes of the ankle (Verhagen et al., 2004). Exercises on balance boards or mats have been shown to reduce the risk of ankle sprains by as much as 50% (Emery et al., 2007; Olsen et al., 2005; Verhagen et al., 2004). However, as for external ankle support, the effect appears to be limited only to athletes with a history of previous sprains. It is also important to note that specific technical training to avoid vulnerable situations may be

effective, at least in volleyball (Bahr et al., 1997). In soccer, late tackles resulting in laterally directed blows to the lower leg serve to put the ankle in a vulnerable position when landing or running (Andersen et al., 2004b; Giza et al., 2003). More specific wording of the rules of the game regarding late tackles, in particular two-footed tackles, with stricter penalties, may prevent ankle injuries (Andersen et al., 2004b; Giza et al., 2003).

General conclusions

Reviewing the present literature, varying evidence available on the prevention of acute sports injuries was found. While there appears to be performed quite extensive research to reduce lower extremity injury risk, much less evidence is available on injury risk factors and mechanisms, and consequently on the prevention of acute upper extremity injuries, as injuries to the hand, wrist, elbow, or shoulder. There are also several injury types which have not been included in the paper, partly because they are rare or simply because there is no data available on prevention. For Achilles tendon ruptures, one of these examples, both factors apply. There is to our knowledge no data on prevention, albeit some on risk factors.

In spite of many promising efforts on injury prevention, a better understanding of injury risk factors and mechanisms will in many cases allow us to optimize current injury prevention strategies, as e.g. exercise programs, and consequently result in fewer injuries, a higher lifelong activity in sports, and lower costs for the public health system.

Recommendations for injury prevention

Approaches that have been proven to be successful in preventing injuries include three main strategies: 1) using equipment designed to reduce injury risk, 2) adapting the rules of play, and 3) specific exercise programs developed to reduce injury risk.

Effective sports injury prevention requires successful implementation of efficacious interventions. This, in turn, requires knowledge about the implementation context including how people, their attitudes and safety (or risk) behaviours interact with these interventions (Finch & Donaldson, 2009). In other words, true injury prevention can only be achieved if some form of behavioural change can be invoked in all involved in an athlete's safety and health, e.g. athlete, coach, referee, etc. A number of the acute injuries observed may be prevented if athletes participate in sports within the limits of their personal qualifications, physical condition, and avoid risky situations. Therefore, one of the major goals should be to establish injury prevention habits early in life.

Equipment

The use of helmets has been introduced in skiing, snowboarding, horse riding, ice hockey, cycling and baseball. In some sports their use is strongly recommended. In others helmets are compulsory when competing. In sports in which falls and collisions are part of the game, elbow and shoulder pads are expected to reduce upper extremity injury risk. Similarly, wrist guards are effective to protect from wrist injuries, at least in snowboarding. However, although such external prophylactic measures are proven efficacious, a true injury risk reduction requires the equipment to be used by the athlete.

Fair play and rules of the game

Many of the rules and regulations that govern sports have been put in place to protect the health of the athlete. Coaches and athletes have a responsibility to play by the rules (fair play), and referees are responsible for penalizing behaviour known to cause injuries. Appropriate referee

training emphasizing the safety aspects of the rules is important. Decisions to change rules can be used actively in making the sport safer for the athletes, and such rule changes can follow from systematic analyses of injury mechanisms. Examples include checking from behind in ice hockey, holding of the arm in team handball, or using an active elbow to the opponents' head/neck when heading in soccer.

Exercise/training

The benefits of preventive exercise training on specifically lower limb injury risk have been documented in several trials. There is conclusive evidence that neuromuscular training assists in the reduction of ankle and knee ligament injuries, and that eccentric strength training can prevent muscular strains of the hamstrings and groin.

Comprehensive neuromuscular training programs combining plyometrics, strength, balance, sport-specific technique, and agility exercises can improve biomechanical measures related to lower extremity injury risk (Hewett et al., 2006a; Hewett et al., 2006b; Myer et al., 2004). While training preventive exercises, emphasis should be placed on a proper landing technique, a soft landing on the forefoot, engaging knee and hip flexion upon landing. A two-foot landing is encouraged whenever possible. In cutting and landing manoeuvres, athletes should avoid excessive dynamic valgus of the knee, always aiming for the 'knee-over-toe position'.

Proper rehabilitation of the index injury: The strong association observed between previous injuries, reduced muscle- and joint-specific function, and the risk for recurrent injuries suggests that secondary prevention of re-injuries should be emphasized. It is of major importance to identify players with previous injuries, impaired neuromuscular control and functional deficits in order to provide optimal treatment, and to prevent further injuries. As is true for all injuries,

appropriate rehabilitation programs and time to allow the athlete to become symptom-free before returning to play is necessary (Steffen et al., 2008a; Hägglund et al., 2006; Myklebust & Bahr, 2005). An athlete who is not able to train without pain or other symptoms from a particular body part should be advised to undergo rehabilitation and restrict participation in competitions, as this probably will increase the risk for a new injury. Hägglund et al. (2007) have developed a 10-step rehabilitation program serving as return-to-play guidelines for soccer players, which has been shown to be effective in preventing re-injuries.

Start early and identify athletes at risk: One goal in sports injury prevention is to teach athletes less vulnerable movement patterns, which may be easier to accomplish with young athletes who have not yet established their basic motion patterns. We therefore suggest that programs to improve basic movement skills and strength, awareness, and neuromuscular control of static and dynamic movements should be implemented in physical education curriculae in schools and in youth sports clubs.

Implementation, establish routines and maintenance: How much risk is reduced depends on the compliance with the exercise program (Soligard et al., 2008), and the compliance of coaches and athletes with injury preventive training is a challenge. It may be difficult to motivate coaches and players to follow such exercise programs merely to prevent injuries, unless there is a direct performance benefit as well. However, it should be emphasized that improved performance will result when team members are free of injury and able to train and compete. Moreover, many of the elements included in evidence-based injury preventive exercise programs also result in improved fitness (Myer et al., 2005; Mjølsnes et al., 2004).

Even so, there is a limit to how much time teams and coaches are willing to spend on exercise programs to prevent injury. For youth and adolescent athletes, who typically practice two to four

times a week, it is not realistic to expect them to spend a similar amount of time on injury prevention exercises, even if they were shown to improve performance.

In our opinion, to successfully implement injury prevention exercises in the training program of youth and adolescents on a consistent basis, the injury prevention exercise protocol must be well accepted by coaches and athletes. The duration of the program should not exceed 20 minutes per session, and preferably be designed to replace the ordinary warm-up exercises. Based on previous research, it appears that such prevention programs should include at least 15 training sessions during the first 6-8 weeks of training, preferably in the pre-season period (Steffen et al., 2008b; Soligard et al., 2008; Hewett et al., 2006a; Olsen et al., 2005; Myklebust et al., 2003). Further evidence is needed to determine whether a pre-season examination, e.g. by a drop-jump or single-leg squat test, can be used to identify athletes with higher risk, e.g. poor knee control. To succeed injury prevention habits and training must be maintained throughout the season, which means that increasing difficulty, variation and sport-specific adjustments of the exercises are needed to keep athletes motivated.

Coach education: The coach plays a key role to successfully implement injury prevention methods, as e.g. exercise programs in warm-up routines. To this end, coaches must understand the principles behind successful injury prevention exercise programs and how to deliver them optimally to ensure high compliance. This goal can only be achieved if injury prevention becomes an integrated part of education programs for coaches at all levels. If efficacious interventions are not widely adopted, complied with and sustained as ongoing practice, then it is very unlikely that any significant health impact will be made (Finch & Donaldson, 2009).

Injury prevention policy

The recommendations stated in the previous paragraphs all relate to those preventive strategies directly involved in the well being of the athlete. The uptake of efficacious preventive measures by athletes, coaches, therapists and referees, is likely to have the most direct effect on injury prevention. Nevertheless, an equally important role in the health of an athlete is ascribed to sport governing bodies and their policies. When a sport governing body includes specific injury prevention strategies as part of its policy, clubs and coaches are more likely to comply and adopt these in their regular routine. Negligence of a sporting organization to promote injury prevention as part of their policies can be considered a 'behavioural' injury causing factor.

References

- Ackery, A., Hagel, B. E., Provvidenza, C., & Tator, C. H. (2007). An international review of head and spinal cord injuries in alpine skiing and snowboarding. *Inj Prev, 13*, 368-375.
- Andersen, T. E., Árnason, A., Engebretsen, L., & Bahr, R. (2004a). Mechanisms of head injuries in elite football. *Br J Sports Med, 38*, 690-696.
- Andersen, T. E., Flørenes, T. W., Árnason, A., & Bahr, R. (2004b). Video analysis of the mechanisms for ankle injuries in football. *Am J Sports Med 32*, 69S-79S.
- Árnason, A., Andersen, T. E., Holme, I., Engebretsen, L., & Bahr, R. (2008). Prevention of hamstring strains in elite soccer: an intervention study. *Scand J Med Sci Sports, 18*, 40-48.
- Árnason, A., Sigurdsson, S. B., Gudmundsson, A., Holme, I., Engebretsen, L., & Bahr, R. (2004). Risk factors for injuries in football. *Am J Sports Med 32*, 5S-16S.
- Askling, C. M., Tengvar, M., Saartok, T., & Thorstenson, A. (2008). Proximal hamstring strains of stretching type in different sports: injury situations, clinical and magnetic resonance imaging characteristics, and return to sport. *Am J Sports Med, 36*, 1799-1804.
- Askling, C., Karlsson, J., & Thorstenson, A. (2003). Hamstring injury occurrence in elite soccer players after preseason strength training with eccentric overload. *Scand J Med Sci Sports, 13*, 244-250.
- Bahr, R., & Krosshaug, T. (2005). Understanding injury mechanisms: a key component of preventing injuries in sport. *Br J Sports Med, 39*, 324-329.
- Bahr, R., & Reeser, J. C. (2003). Injuries among world-class professional beach volleyball players. The Federation Internationale de Volleyball beach volleyball injury study. *Am J Sports Med, 31*, 119-125.
- Bahr, R., & Bahr, I. A. (1997). Incidence of acute volleyball injuries: a prospective cohort study of injury mechanisms and risk factors. *Scand J Med Sci Sports, 7*, 166-171.
- Bahr, R., Lian, Ø., & Bahr, I. A. (1997). A twofold reduction in the incidence of acute ankle sprains in volleyball after the introduction of an injury prevention program: a prospective cohort study. *Scand J Med Sci Sports, 7*, 172-177.
- Boden, B. P., Tacchetti, R., & Mueller, F. O. (2003). Catastrophic cheerleading injuries. *Am J Sports Med, 31*, 881-888.
- Browne, E. Z., Dunn, H. K., & Snyder, C. C. (1976). Ski pole thumb injury. *Plast Reconstr Surg, 58*, 17-23.

- Chandrashekar, N., Mansouri, H., Slauterbeck, J. R., & Hashemi, J. (2006). Sex-based differences in the tensile properties of the human anterior cruciate ligament. *J Biomech*, *39*, 2943-2950.
- DeMorat, G., Weinhold, P., Blackburn, T., Chudik, S., & Garrett, T. (2004). Aggressive quadriceps loading can induce noncontact anterior cruciate ligament injury. *Am J Sports Med*, *32*, 477-483.
- Ekstrand, J., & Gillquist, J. (1983). Soccer injuries and their mechanisms: a prospective study. *Med Sci Sports Exerc*, *15*, 267-270.
- Emery, C. A., Rose, M. S., McAllister, J. R., & Meeuwisse, W. H. (2007). A prevention strategy to reduce the incidence of injury in high school basketball: a cluster randomized controlled trial. *Clin J Sport Med*, *17*, 17-24.
- Emery, C. A., Meeuwisse, W. H., & Hartmann, S. E. (2005). Evaluation of risk factors for injury in adolescent soccer: implementation and validation of an injury surveillance system. *Am J Sports Med*, *33*, 1882-1891.
- Engebretsen, L., & Bahr, R. (2009). Why is injury prevention in sports important? In: Bahr, R., & Engebretsen, L. (Eds.), *Sports injury prevention* (pp.1-6). Wiley-Blackwell.
- Ettlinger, C. F., Johnson, R. J., & Shealy, J. E. (1995). A method to help reduce the risk of serious knee sprains incurred in alpine skiing. *Am J Sports Med*, *23*, 531-537.
- Faude, O., Junge, A., Kindermann, W., & Dvorak, J. (2006). Risk factors for injuries in elite female soccer players. *Br J Sports Med*, *40*, 785-790.
- Finch, C. F., & Donaldson, A. (2009). A sports setting matrix for understanding the implementation context for community sport. *Br J Sports Med*, [Epub ahead of print, Feb 6].
- Freeman, M.A. (1965). Instability to the foot after injuries to the lateral ligament of the ankle. *J Bone Joint Surg Br*, *47*, 669-677.
- Fricker, R. & Hintermann, B. (1995). Skier's thumb. Treatment, prevention and recommendations. *Sports Med*, *19*, 73-79.
- Fuller, C. W., Junge, A., & Dvorak, J. (2005). A six year prospective study of the incidence and causes of head and neck injuries in international football. *Br J Sports Med*, *39 Suppl 1*, i3-i9.
- Fung, D.T., & Zhang, L.Q. (2003). Modeling of ACL impingement against the intercondylar notch. *Clin Biomech*, *18*, 933-941.
- Garrett, W. E. (1990). Muscle strain injuries: clinical and basic aspects. *Med Sci Sports Exerc*, *22*, 436-443.
- Gilchrist, J., Mandelbaum, B. R., Melancon, H., Ryan, G. W., Silvers, H. J., Griffin, L. Y., Watanabe, D. S., Dick, R. W., & Dvorak, J. (2008). A randomized controlled trial to prevent

noncontact anterior cruciate ligament injury in female collegiate soccer players. *Am J Sports Med*, 36, 1476-1483.

Giza, E., Fuller, C., Junge, A., & Dvorak, J. (2003). Mechanisms of foot and ankle injuries in soccer. *Am J Sports Med* 31, 550-554.

Goulet, C., Hagel, B. E., Hamel, D., & Legare, G. (2009). Self-reported skill level and injury severity in skiers and snowboarders. *J Sci Med Sport*, [Epub ahead of print, Jan 3].

Griffin, L. Y., Albohm, M. J., Arendt, E. A., Bahr, R., Beynon, B. D., Demaio, M., Dick, R. W., Engebretsen, L., Garrett, W. E., Jr., Hannafin, J. A., Hewett, T. E., Huston, L. J., Ireland, M. L., Johnson, R. J., Lephart, S., Mandelbaum, B. R., Mann, B. J., Marks, P. H., Marshall, S. W., Myklebust, G., Noyes, F. R., Powers, C., Shields, C., Shultz, S. J., Silvers, H., Slauterbeck, J., Taylor, D. C., Teitz, C. C., Wojtys, E. M., & Yu, B. (2006). Understanding and preventing noncontact anterior cruciate ligament injuries: a review of the Hunt Valley II meeting, January 2005. *Am J Sports Med*, 34, 1512-1532.

Hagel, B. (2005). Skiing and snowboarding injuries. *Med Sport Sci*, 48, 74-119.

Hägglund, M., Waldén, M., & Ekstrand, J. (2007). Lower reinjury rate with a coach-controlled rehabilitation program in amateur male soccer: a randomized controlled trial. *Am J Sports Med*, 35, 1433-1442.

Hägglund, M., Waldén, M., & Ekstrand, J. (2006). Previous injury as a risk factor for injury in elite football: a prospective study over two consecutive seasons. *Br J Sports Med*, 40, 767-772.

Haskell, W. L., Lee, I. M., Pate, R. R., Powell, K. E., Blair, S. N., Franklin, B. A., Macera, C. A., Heath, G. W., Thompson, P. D., & Bauman, A. (2007). Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc*, 39, 1423-1434.

Hewett, T.E., Zazulak, B.T., Myer, G.D. (2007). Effects of the menstrual cycle on anterior cruciate ligament injury risk: a systematic review. *Am J Sports Med*, 35, 657-668.

Hewett, T. E., Myer, G. D., & Ford, K. R. (2006a). Anterior cruciate ligament injuries in female athletes: part 1, mechanisms and risk factors. *Am J Sports Med*, 34, 299-311.

Hewett, T. E., Ford, K. R., & Myer, G. D. (2006b). Anterior cruciate ligament injuries in female athletes: part 2, a meta-analysis of neuromuscular interventions aimed at injury prevention. *Am J Sports Med*, 34, 490-498.

Hewett, T. E., Myer, G. D., Ford, K. R., Heidt, R. S., Jr., Colosimo, A. J., McLean, S. G., van den Bogert, A. J., Paterno, M. V., & Succop, P. (2005). Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med*, 33, 492-501.

- Hölmich, P., Uhrskou, P., Ulnits, L., Kanstrup, I. L., Nielsen, M. B., Bjerg, A. M., & Krogsgaard, K. (1999). Effectiveness of active physical training as treatment for long-standing adductor-related groin pain in athletes: randomised trial. *Lancet*, *353*, 439-443.
- Hootman, J. M., Dick, R., & Agel, J. (2007). Epidemiology of collegiate injuries for 15 sports: summary and recommendations for injury prevention initiatives. *J Athl Train*, *42*, 311-319.
- Hutchinson MR, & Andrews JR. (2009). Preventing elbow injuries. In: Bahr, R., & Engebretsen, L. (Eds.), *Sports injury prevention* (pp. 153-174). Wiley-Blackwell.
- Janda, D. H. (2003). The prevention of baseball and softball injuries. *Clin Orthop Relat Res*, 20-28.
- Jansen, J. A., Mens, J. M., Backx, F. J., Kolfschoten, N., & Stam, H. J. (2008). Treatment of longstanding groin pain in athletes: a systematic review. *Scand J Med Sci Sports*, *18*, 263-274.
- Kettunen, J. A., Kujala, U. M., Kaprio, J., Koskenvuo, M., & Sarna, S. (2001). Lower-limb function among former elite male athletes. *Am J Sports Med*, *29*, 2-8.
- Kirkendall, D. T., Jordan, S. E., & Garrett, W. E. (2001). Heading and head injuries in soccer. *Sports Med*, *31*, 369-386.
- Krogsgaard, MR., Safran, M., Rheinländer, P., & Cheung, E. (2009). Preventing shoulder injuries. In: Bahr, & R., Engebretsen, L. (Eds.), *Sports injury prevention* (pp. 134-152). Wiley-Blackwell.
- Krosshaug, T., Andersen, T. E., Olsen, O. E., Myklebust, G., & Bahr, R. (2005). Research approaches to describe the mechanisms of injuries in sport: limitations and possibilities. *Br J Sports Med*, *39*, 330-339.
- Krosshaug, T., Nakamae, A., Boden, B. P., Engebretsen, L., Smith, G., Slauterbeck, J. R., Hewett, T. E., & Bahr, R. (2007a). Mechanisms of anterior cruciate ligament injury in basketball: video analysis of 39 cases. *Am J Sports Med*, *35*, 359-367.
- Krosshaug, T., Slauterbeck, J. R., Engebretsen, L., & Bahr, R. (2007b). Biomechanical analysis of anterior cruciate ligament injury mechanisms: three-dimensional motion reconstruction from video sequences. *Scand J Med Sci Sports*, *17*, 508-519.
- Kujala, U. M., Taimela, S., Viljanen, T., Jutila, H., Viitasalo, J. T., Videman, T., & Battie, M. C. (1996). Physical loading and performance as predictors of back pain in healthy adults. A 5-year prospective study. *Eur J Appl Physiol Occup Physiol*, *73*, 452-458.
- LaMonte, M. J., Blair, S. N., & Church, T. S. (2005). Physical activity and diabetes prevention. *J Appl Physiol*, *99*, 1205-1213.
- Langran, M., & Selvaraj, S. (2004). Increased injury risk among first-day skiers, snowboarders, and skiboarders. *Am J Sports Med*, *32*, 96-103.

- Le Gall, F., Carling, C., & Reilly, T. (2008). Injuries in young elite female soccer players: an 8-season prospective study. *Am J Sports Med*, *36*, 276-284.
- Lohmander, L. S., Östenberg, A., Englund, M., & Roos, H. (2004). High prevalence of knee osteoarthritis, pain, and functional limitations in female soccer players twelve years after anterior cruciate ligament injury. *Arthritis Rheum*, *50*, 3145-3152.
- Machold, W., Kwasny, O., Eisenhardt, P., Kolonja, A., Bauer, E., Lehr, S., Mayr, W., & Fuchs, M. (2002). Reduction of severe wrist injuries in snowboarding by an optimized wrist protection device: a prospective randomized trial. *J Trauma*, *52*, 517-520.
- Macintyre, J., Johson, C., & Schroeder, E. L. (2006). Groin pain in athletes. *Curr Sports Med Rep*, *5*, 293-299.
- Macnab, A. J., Smith, T., Gagnon, F. A., & Macnab, M. (2002). Effect of helmet wear on the incidence of head/face and cervical spine injuries in young skiers and snowboarders. *Inj Prev*, *8*, 324-327.
- Maffey, L., & Emery, C. (2007). What are the risk factors for groin strain injury in sport? A systematic review of the literature. *Sports Med*, *37*, 881-894.
- Mandelbaum, B. R., Silvers, H. J., Watanabe, D. S., Knarr, J. F., Thomas, S. D., Griffin, L. Y., Kirkendall, D. T., & Garrett, W. (2005). Effectiveness of a neuromuscular and proprioceptive training program in preventing anterior cruciate ligament injuries in female athletes: 2-year follow-up. *Am J Sports Med*, *33*, 1003-1010.
- Markolf, K. L., Burchfield, D. M., Shapiro, M. M., Shepard, M. F., Finerman, G. A., & Slauterbeck, J. L. (1995). Combined knee loading states that generate high anterior cruciate ligament forces. *J Orthop Res*, *13*, 930-935.
- McCrory, P., & Turner, M. (2005). Equestrian injuries. *Med Sport Sci*, *48*, 8-17.
- McIntosh, A.S., McCrory, P., Finch, C.F., Best, J.P., Chalmers, D.J., & Wolfe, R. (2009). Does padded headgear prevent head injury in rugby union football? *Med Sci Sports Exerc*, *41*, 306-313.
- McIntosh, A. S., & McCrory, P. (2005). Preventing head and neck injury. *Br J Sports Med*, *39*, 314-318.
- Meeuwisse, W. H., Tyreman, H., Hagel, B., & Emery, C. (2007). A dynamic model of etiology in sport injury: the recursive nature of risk and causation. *Clin J Sport Med*, *17*, 215-219.
- Meyer, E.G., & Haut, R.C. (2008). Anterior cruciate ligament injury induced by internal tibial torsion or tibiofemoral compression. *J Biomechan*, *41*, 3377-3383.
- Mjølsnes, R., Árnason, A., Østhagen, T., Raastad, T., & Bahr, R. (2004). A 10-week randomized trial comparing eccentric vs. concentric hamstrings strength training in well-trained soccer players. *Scand J Med Sci Sports*, *14*, 1-7.

Mogan, J. V., & Davis, P. H. (1982). Upper extremity injuries in skiing. *Clin Sports Med*, 1, 295-308.

Murphy, D. F., Connolly, D. A., & Beynon, B. D. (2003). Risk factors for lower extremity injury: a review of the literature. *Br J Sports Med*, 37, 13-29.

Myer, G. D., Ford, K. R., Palumbo, J. P., & Hewett, T. E. (2005). Neuromuscular training improves performance and lower-extremity biomechanics in female athletes. *J Strength Cond Res*, 19, 51-60.

Myer, G. D., Ford, K. R., & Hewett, T. E. (2004). Methodological approaches and rationale for training to prevent anterior cruciate ligament injuries in female athletes. *Scand J Med Sci Sports*, 14, 275-285.

Myklebust, G., & Bahr, R. (2005). Return to play guidelines after anterior cruciate ligament surgery. *Br J Sports Med*, 39, 127-131.

Myklebust, G., Engebretsen, L., Brækken, I. H., Skjøelberg, A., Olsen, O. E., & Bahr, R. (2003). Prevention of anterior cruciate ligament injuries in female team handball players: a prospective intervention study over three seasons. *Clin J Sport Med*, 13, 71-78.

Myklebust, G., Maehlum, S., Holm, I., & Bahr, R. (1998). A prospective cohort study of anterior cruciate ligament injuries in elite Norwegian team handball. *Scand J Med Sci Sports*, 8, 149-153.

Olsen, O. E., Myklebust, G., Engebretsen, L., Holme, I., & Bahr, R. (2005). Exercises to prevent lower limb injuries in youth sports: cluster randomised controlled trial. *BMJ*, 330, 449-455.

Olsen, O. E., Myklebust, G., Engebretsen, L., & Bahr, R., (2004). Injury mechanisms for anterior cruciate ligament injuries in team handball: a systematic video analysis. *Am J Sports Med*, 32, 1002-1012.

Orchard, J. (2002). Is there a relationship between ground and climatic conditions and injuries in football? *Sports Med*, 32, 419-432.

Proske, U., Morgan, D.L., Brockett, C.L., & Percival, P. (2004). Identifying athletes at risk of hamstring strains and how to protect them. *Clin Exp Pharmacol Physiol*, 31, 546-50.

Pujol, N., Blanchi, M. P., & Chambat, P. (2007). The incidence of anterior cruciate ligament injuries among competitive Alpine skiers: a 25-year investigation. *Am J Sports Med*, 35, 1070-1074.

Quatman, C.E. & Hewett, T.E. (2009). The ACL injury controversy: is "valgus collapse" a sex-specific mechanism? *Br J Sports Med*, [Epub ahead of print, April 15].

Renström, P., Ljungqvist, A., Arendt, E., Beynon, B., Fukubayashi, T., Garrett, W., Georgoulis, T., Hewett, T. E., Johnson, R., Krosshaug, T., Mandelbaum, B., Micheli, L., Myklebust, G., Roos, E., Roos, H., Schamasch, P., Shultz, S., Werner, S., Wojtys, E., & Engebretsen, L. (2008).

Non-contact ACL injuries in female athletes: an International Olympic Committee current concepts statement. *Br J Sports Med*, 42, 394-412.

Rønning, R., Rønning, I., Gerner, T., & Engebretsen, L. (2001). The efficacy of wrist protectors in preventing snowboarding injuries. *Am J Sports Med*, 29, 581-585.

Russell, K., Hagel, B., & Francescutti, L. H. (2007). The effect of wrist guards on wrist and arm injuries among snowboarders: a systematic review. *Clin J Sport Med*, 17, 145-150.

Shimokochi, Y., & Shultz, S. J. (2008). Mechanisms of noncontact anterior cruciate ligament injury. *J Athl Train*, 43, 396-408.

Söderman, K., Adolphson, J., Lorentzon, R., & Alfredson, H. (2001). Injuries in adolescent female players in European football: a prospective study over one outdoor soccer season. *Scand J Med Sci Sports*, 11, 299-304.

Söderman, K., Werner, S., Pietila, T., Engström, B., & Alfredson, H. (2000). Balance board training: prevention of traumatic injuries of the lower extremities in female soccer players? A prospective randomized intervention study. *Knee Surg Sports Traumatol Arthrosc*, 8, 356-363.

Soligard, T., Myklebust, G., Steffen, K., Holme, I., Silvers, H., Bizzini, M., Junge, A., Dvorak, J., Bahr, R., & Andersen, T. E. (2008). Comprehensive warm-up programme to prevent injuries in young female footballers: cluster randomised controlled trial. *BMJ*, 337, a2469.

Steffen, K., Myklebust, G., Andersen, T. E., Holme, I., & Bahr, R. (2008a). Self-reported injury history and lower limb function as risk factors for injuries in female youth soccer. *Am J Sports Med*, 36, 700-708.

Steffen, K., Myklebust, G., Olsen, O. E., Holme, I., & Bahr, R. (2008b). Preventing injuries in female youth football - a cluster-randomized controlled trial. *Scand J Med Sci Sports*, 18, 605-614.

Steffen, K., Andersen, T. E., & Bahr, R. (2007). Risk of injury on artificial turf and natural grass in young female football players. *Br J Sports Med*, 41 Suppl 1, i33-i37.

Straume-Næsheim, T. M., Andersen, T. E., Dvorak, J., & Bahr, R. (2005). Effects of heading exposure and previous concussions on neuropsychological performance among Norwegian elite footballers. *Br J Sports Med*, 39 Suppl 1, i70-i77.

Sulheim, S., Holme, I., Ekeland, A., & Bahr, R. (2006). Helmet use and risk of head injuries in alpine skiers and snowboarders. *JAMA*, 295, 919-924.

Surve, I., Schwellnus, M. P., Noakes, T., & Lombard, C. (1994). A fivefold reduction in the incidence of recurrent ankle sprains in soccer players using the Sport-Stirrup orthosis. *Am J Sports Med*, 22, 601-606.

Thompson, D. C., Rivara, F. P., & Thompson, R. (2000). Helmets for preventing head and facial injuries in bicyclists. *Cochrane Database Syst Rev*, CD001855.

- Torjussen, J., & Bahr, R. (2006). Injuries among elite snowboarders (FIS Snowboard World Cup). *Br J Sports Med*, 40, 230-234.
- Torjussen, J., & Bahr, R. (2005). Injuries among competitive snowboarders at the national elite level. *Am J Sports Med*, 33, 370-377.
- Tropp, H., Askling, C., & Gillquist, J. (1985). Prevention of ankle sprains. *Am J Sports Med*, 13, 259-262.
- Tyler, T. F., Nicholas, S. J., Campbell, R. J., & McHugh, M. P. (2001). The association of hip strength and flexibility with the incidence of adductor muscle strains in professional ice hockey players. *Am J Sports Med*, 29, 124-128.
- Verhagen, E., van der Beek, A., Twisk, J., Bouter, L., Bahr, R., & van Mechelen, W. (2004). The effect of a proprioceptive balance board training program for the prevention of ankle sprains: a prospective controlled trial. *Am J Sports Med* 32, 1385-1393.
- Verhagen, E. A., van Tulder, M., van der Beek, A. J., Bouter, L. M., & van Mechelen W. (2005). An economic evaluation of a proprioceptive balance board training programme for the prevention of ankle sprains in volleyball. *Br J Sports Med*, 39, 111-115.
- Verrall, G. M., Hamilton, I. A., Slavotinek, J. P., Oakeshott, R. D., Spriggins, A. J., Barnes, P. G., & Fon, G. T. (2005). Hip joint range of motion reduction in sports-related chronic groin injury diagnosed as pubic bone stress injury. *J Sci Med Sport*, 8, 77-84.
- von Porat, A., Roos, E. M., & Roos, H. (2004). High prevalence of osteoarthritis 14 years after an anterior cruciate ligament tear in male soccer players: a study of radiographic and patient relevant outcomes. *Ann Rheum Dis*, 63, 269-273.
- Waldén, M., Hägglund, M., & Ekstrand, J. (2006). High risk of new knee injury in elite footballers with previous anterior cruciate ligament injury. *Br J Sports Med*, 40, 158-162.
- Yu, B. & Garrett, W.E. (2007). Mechanisms of non-contact ACL injuries. *Br J Sports Med*, 41 Suppl 1, i47-i51.
- Zazulak, B. T., Hewett, T. E., Reeves, N. P., Goldberg, B., & Cholewicki, J. (2007a). Deficits in neuromuscular control of the trunk predict knee injury risk: a prospective biomechanical-epidemiologic study. *Am J Sports Med*, 35, 1123-1130.
- Zazulak, B. T., Hewett, T. E., Reeves, N. P., Goldberg, B., & Cholewicki, J. (2007b). The effects of core proprioception on knee injury: a prospective biomechanical-epidemiological study. *Am J Sports Med*, 35, 368-373.