

This file was downloaded from the institutional repository Brage NIH - brage.bibsys.no/nih

Aas, S. N., Seynnes, O. R., Benestad, H. B., Raastad, T. (2019). Strength training and protein supplementation improve muscle mass, strength, and function in mobility-limited older adults: a randomized controlled trial. *Aging Clinical and Experimental Research*, 32, 605–616.

Dette er siste tekst-versjon av artikkelen, og den kan inneholde små forskjeller fra forlagets pdf-versjon. Forlagets pdf-versjon finner du her:
<https://link.springer.com/article/10.1007%2Fs40520-019-01234-2>

This is the final text version of the article, and it may contain minor differences from the journal's pdf version. The original publication is available here:
<https://link.springer.com/article/10.1007%2Fs40520-019-01234-2>

Strength Training and Protein Supplementation Improve Muscle Mass, Strength, and Function in Mobility-Limited Older Adults: A Randomized Controlled Trial

Sigve Nyvik Aas^{1*}, Olivier Seynnes¹, Haakon B. Benestad², Truls Raastad¹

¹Department of Physical Performance, Norwegian School of Sport Sciences, Oslo, Norway

²Section of Anatomy, Institute of Basic Medical Sciences, University of Oslo, Norway

*** Correspondence:**

Sigve Nyvik Aas

s.n.aas@nih.no

Phone: 0047 414 99 074

ORCID ID of Sigve Nyvik Aas: 0000-0003-1196-0632

Keywords: resistance exercise, frailty, sarcopenia, hypertrophy, mobility-limited, skeletal muscle.

ABSTRACT

Background: Adaptation to strength training in very old mobility-limited individuals is not fully characterized. Therefore, the aim of this study was to perform a thorough investigation of the adaptation to a lower body strength training regime in this population, with particular emphasis on the relationship between changes in selected variables. **Methods:** Twenty-two mobility-limited older men and women (85 ± 6 years) were randomized to either a group performing thirty minutes of heavy-load strength training three times a week, with daily protein supplementation, for ten weeks (ST), or a control group (CON). Endpoints were leg lean mass assessed by DXA, muscle thickness assessed by ultrasound, isometric and dynamic strength, rate of torque development, and functional capacity. **Results:** Leg lean mass increased from baseline in ST (0.7 ± 0.3 kg), along with increased thickness of vastus lateralis ($4.4 \pm 3.2\%$), rectus femoris ($6.7 \pm 5.1\%$), and vastus intermedius ($5.8 \pm 5.9\%$). The hypertrophy was accompanied by improved knee extensor strength (20-23%) and functional performance (7-11%). In ST, neither the change in leg lean mass nor muscle thickness correlated with changes in muscle strength. However, a strong correlation was observed between the change in isometric strength and gait velocity ($r = 0.70$). **Conclusions:** The mismatch between gains in muscle size and strength suggest that muscle quality related adaptations contributed to the increases in strength. The correlations observed between improvements in strength and function suggests that interventions eliciting large improvements in strength may also be superior in terms of functional gains in this population.

INTRODUCTION

Muscle strength is associated with functional performance in older adults (1), and various tasks of daily living are therefore hampered by the rapid decline in strength at very high age. The Short Physical Performance Battery (SPPB) is commonly used to assess functional capacity, and individuals with a score of 10 or less out of maximum 12 may be categorized as mobility-limited. These individuals represent a group of particular interest, because small-to-moderate limitations in functional status assessed by SPPB is associated with higher odds of losing the ability to walk 400 m three years later (2).

It is established that heavy-load strength training, alone or in combination with protein supplementation, can improve muscle mass, strength and function in healthy older adults (3-5). In contrast, studies in very old individuals with certain mobility limitations are scarce and report discrepant findings. Gains in lean mass are observed by some (6), but not all (7), and information about the hypertrophic adaptation within specific muscles are lacking. Training effects on knee extension strength measured as one repetition maximum (1RM) vary greatly, from ~25 to ~200% (6-9). Isometric strength gains are substantially lower, ranging from ~5 to 37% (7, 8). Findings are also inconsistent with respect to functional improvements (7, 9). Therefore, the extent to which training-induced gains in muscle mass and size are related to improvements in strength and function is still poorly understood in this population, and studies combining different methods to assess changes in muscle mass, strength and function are therefore needed. Moreover, to improve the basis for investigating such relationships, it might be beneficial to include a protein supplement to optimize the gains in muscle mass (10).

Hence, the aim of the present investigation was to conduct a thorough examination of the adaptation to strength training and protein supplementation in very old mobility-limited individuals. By complementing leg lean mass changes with ultrasound measurements of quadriceps femoris, we aimed to precisely characterize changes in muscle mass. Moreover, both dynamic and isometric strength testing, as well as several functional tests were performed, to investigate relationships between improvements in muscle function and functional performance. We hypothesized that strength training and protein supplementation would elicit increases in both leg lean mass and muscle thickness, but that these increases would be exceeded by greater improvements in muscle strength. Besides, we hypothesized that muscle hypertrophy would show stronger correlations with changes in isometric than dynamic strength.

Lastly, we hypothesized that improvements in both dynamic and isometric strength would correlate with improvements in functional performance.

METHOD AND DESIGN

The study was conducted from August to December 2015 in Oslo, Norway. Twenty-two older men and women aged between 73 and 92 [85 ± 6 years; mean \pm standard deviation (SD)] were recruited through advertisement in the local community and visits at a local senior activity center. Upon inclusion, subjects were screened using the Short Physical Performance Battery (SPPB). The SPPB consists of timed standing balance, gait speed, and timed chair-rise assessments (11). Performance for each of these tasks is scored between 0 and 4, with a summary score of 0–12. Participants with an SPPB score ≤ 10 were included. In addition, individuals using aids such as walkers, canes or Nordic walking sticks in their daily life were included, regardless of SPPB score. Participants were excluded if they had significant cognitive impairment, unstable chronic illness, cardiac arrhythmia, had experienced a fracture of a lower extremity during the last year, were smokers, or had other musculoskeletal problems making testing and training impossible.

The study complied with the standards set by the Declaration of Helsinki. The nature and goals of the study were thoroughly explained, and all participants provided a written informed consent. The study was evaluated by the Regional Committee for Medical & Health Research Ethics, Section C, South East Norway. The study data storage methods were in accordance with The Norwegian Centre for Research Data. The study was retrospectively registered at clinicaltrials.gov as NCT03723902 (URL: <https://clinicaltrials.gov/show/NCT03723902>). After baseline assessment, stratified randomization was performed on ranked baseline knee extensor strength (normalized to body weight), five times chair-rise and stair climbing performance. Randomization was performed by a researcher not involved in the study, using an online random allocation software. Participants were assigned in a 1:1 ratio to a group performing heavy load strength training three times per week and receiving daily protein supplementation, both for ten weeks (ST), or a control group continuing their normal daily activities and dietary habits (CON) (Figure 1).

Fig. 1 Participant flow chart

Training

Training of participants in ST was carried out at a local senior activity center. The strength training protocol was a supervised progressive program, performed three times per week for ten weeks. The training was performed in groups of two to three participants, and each session took approximately thirty minutes to complete, including warm-up. Sessions were usually performed Mondays, Wednesdays and Fridays, where Mondays and Fridays were repetition maximum (RM) sessions. On Wednesdays, the load was adjusted to approximately 80-90% of RM loads. Each session included leg press, knee extension, and adapted one-leg squat. In the latter exercise, subjects stood sideways toward a wall bar for support (one hand), with the opposite leg on a step platform (initial height of 20 cm). From this position, subjects extended their knee and hip, before returning to the starting position. After a given number of repetitions, the participant turned around 180 degrees, to perform the exercise for the other leg. Progression was achieved by gradually increasing the height of the step platform, and by adding weight (dumbbell in the arm not used for support). For all three exercises, subjects were instructed to use 1-2 seconds both on the concentric and eccentric phase. The participants performed 1-3 sets of 12 repetitions at submaximal loads the first week, and from week 2-10 subjects gradually progressed from 3 sets of 12RM to 3-4 sets of 6RM (Table 1). Each set was followed by approximately 2-min rest periods. Strength training sessions were preceded by a lower extremity 3-min warm-up exercise, using a step platform.

Table 1 Strength training program performed in ST

	Week 1			Week 2-4			Week 5-7			Week 8-10		
	WO 1	WO 2	WO 3	WO 1	WO 2	WO 3	WO 1	WO 2	WO 3	WO 1	WO 2	WO 3
Leg press	2 x 12 (<RM)	2 x 12 (<RM)	2 x 12 (<RM)	3 x 12 (RM)	3 x 10 (<RM)	3 x 10 (RM)	3 x 10 (RM)	3 x 10 (<RM)	3 x 8 (RM)	3 x 8 (RM)	3 x 10 (<RM)	4 x 6 (RM)
Knee extension	3 x 12 (<RM)	3 x 12 (<RM)	3 x 12 (<RM)	3 x 12 (RM)	3 x 10 (<RM)	3 x 10 (RM)	3 x 10 (RM)	2 x 10 (<RM)	3 x 8 (RM)	4 x 8 (RM)	4 x 10 (<RM)	4 x 6 (RM)
Adapted one-leg squat	1 x 12 (<RM)	1 x 12 (<RM)	1 x 12 (<RM)	1 x 12 (RM)	1 x 10 (<RM)	1 x 10 (RM)	1 x 10 (RM)	2 x 10 (<RM)	2 x 8 (RM)	2 x 8 (RM)	2 x 10 (<RM)	2 x 6 (RM)

WO, work-out. RM: repetition maximum

Supplementation

Participants in ST received 34 g of milk protein each day, divided in two equal servings (each contained 17 g of protein, 18 g of carbohydrate, 1 g fat; giving 149 kcal [627 kJ]). Participants consumed one supplement in the morning and one in the evening. On training days, subjects were instructed to consume one of the servings within two hours after training. Participants were encouraged to continue their regular food routines in addition to the supplements.

Tests and measurements

All tests were monitored by the same investigator before and after the intervention period. Neither participants nor test personnel were blinded during testing. Analyses of ultrasound images were performed in a blinded manner. All testing took place at Norwegian School of Sport Sciences.

Diet assessment

24-hour diet recall interviews were performed to evaluate whether the supplement had changed the participants' diet. Two interviews were conducted before, and two toward the end of the intervention period.

Body composition

Dual energy x-ray absorptiometry. Fat and lean mass was assessed with a DXA scan after an overnight fast. Regional analyses were performed to investigate leg lean mass (primary outcome). The test-retest reliability for measuring leg lean mass is high in our laboratory (CV < 1.7%).

Quadriceps thickness

Muscle thickness was measured in vastus lateralis, rectus femoris, and vastus intermedius muscles of the dominant leg, with a 50-mm linear probe (5–12 MHz) connected to a Philips HD11 XE ultrasound apparatus (Royal Philips Electronics, Amsterdam, Netherlands). Scans of vastus lateralis were obtained at 40% of the distance between the knee joint and the greater trochanter, and halfway between the iliac crest and the myotendinous junction for rectus femoris. Vastus intermedius was measured on the same image as vastus lateralis, where the latter was thickest in the medio-lateral direction. Probe position was recorded for each measurement on transparent, acetate paper positioned over the thigh.

Several images were obtained for each muscle at each registration time point, and the three best images were chosen for analysis, which was performed using OsiriX v5.5.1 (Bernex, Switzerland). The shortest distance between the upper and lower aponeurosis was measured at 25, 50 and 75% of the width of the field of view. Muscle thickness was calculated as the average of these measurements across images.

Muscle strength

Maximal strength was assessed with a one-repetition maximum (1RM) in knee extension. After a general warm-up (5 min bicycling), a specific warm-up/preparation procedure was performed using loads corresponding to 50%, 70%, 80% and 90% of expected 1RM – conducting 10, 6, 3, and 1 repetitions, respectively. Optimally, 2–5 attempts were used to establish 1RM. The loads could be adjusted in steps as low as 0.5 kg. The left and right leg was tested interchangeably (unilateral), and each leg rested for approximately three minutes between attempts. The range of motion was strictly controlled.

Isometric maximal voluntary contraction (MVC) was also tested for the knee extensors. The participants were securely strapped to the testing chair, with an angle of 90° in the hip and knee joints (0° corresponding to full extension). The lever arm of the knee-extension device with an attached strength gauge (HBM U2AC2, Darmstadt, Germany) was adjusted so that the contact point with the participant's calf was just proximal to the line of the medial malleolus. The participants performed three 5-second attempts and were instructed to reach maximum torque (Newton meter - Nm) as rapidly as possible. This was done to obtain a measure of the rate of torque development (RTD), both in relative terms ($RTD_{max}/\text{maximal torque}$) and absolute terms (torque at 100 ms). The left and right leg was tested interchangeably, with sixty seconds of rest between each attempt. The best result of three trials per leg was used in the analyses.

Functional performance

Sit to stand x 5

Participants were asked to rise from a chair to a full stand five times as fast as possible. Subjects were instructed to keep their hands crossed on their chest. If they were not able to stand up without using their hands, the use of hands was allowed, and performed the same way at both registration time points. Time was measured with a pressure sensor (TC-Start Pod and Touch Pad, Brower Timing Systems, Draper, USA). The best time of two trials was used in the analyses.

Stair climbing test

Participants were asked to climb a stair with twenty steps as fast as possible, without running. Time was assessed by photocells (Brower Timing Systems, Draper, USA). Steps were 18 cm high and 30 long (horizontal length). The total height was 3.6 m and the total length was 6 m. The participants were encouraged to climb the stairs without using the handrail, but were

allowed to do so if they felt it was necessary. If so, hands were used at both registration time points. The best time of two trials was used in the analyses.

Habitual gait velocity

Participants were asked to walk 10 meters at their habitual gait velocity. Photo cells (Brower Timing Systems, Draper, USA) were placed to measure the time between 2 to 8 meters. The mean time of three trials was used in the analyses.

Statistical analyses

Leg lean mass was the primary outcome. The sample size calculation was based on values obtained from a previous study (10), and on unpublished results from our own lab. Using an online calculator (clincalc.com), power calculations indicated that 22 participants would be needed to identify a difference of 0.5 kg between the ST and CON for leg lean mass (alpha set at 0.05, beta at 0.20), with an expected standard deviation (SD) of 0.4 kg. Results for the functional tests (five times chair-rise, stair climbing and habitual gait velocity) were not normally distributed (D'Agostino and Pearson omnibus normality test), and were therefore log-transformed prior to statistical analysis. Differences at baseline were assessed with an unpaired Student's t test. A two-way ANOVA and Sidak's multiple comparison test was used to evaluate the effect of time and group for variables measured both before and after the intervention period. Differences in response between vastus lateralis, rectus femoris and vastus intermedius were analyzed with one-way ANOVA, and Tukey's multiple comparisons test. Associations between variables were investigated using Pearson's correlation coefficient. Correlations were considered weak if $r < 0.4$, moderate if $r = 0.4-0.6$, and strong if $r > 0.6$. Statistical analyses were performed using GraphPad Prism6 software (GraphPad Software, Inc., La Jolla, CA). Two-sided statistical significance was set at $P < 0.05$.

RESULTS

There were no significant differences between ST and CON in physical characteristics or protein intake at baseline, although ST tended to be older than CON (Table 2). All 22 subjects completed the study. However, one subject in CON was excluded from the per-protocol analysis, as the subject started strength training during the intervention period. Thus, 11 subjects in ST and 10 subjects in CON remained for the per-protocol analyses. Moreover, one additional subject in CON did not perform the ultrasound measurement after the intervention. Thus, for

this analysis, the number of subjects in ST and CON is 11 and 9, respectively. No adverse events occurred in any of the groups.

Table 2 Baseline participant characteristics

	ST (n = 11) N or Mean \pm SD	CON (n = 11) N or Mean \pm SD	P value
Men / women (n)	4 / 7	3 / 8	
Age	86.6 \pm 6.0	82.6 \pm 4.5	0.09
Height (m)	1.68 \pm 0.07	1.65 \pm 0.10	0.50
Body mass (kg)	73.6 \pm 12.7	71.5 \pm 9.5	0.67
Lean mass (kg)	44.8 \pm 5.6	41.5 \pm 4.5	0.14
BMI (body mass/height ²)	26.1 \pm 4.2	26.3 \pm 3.8	0.94
Protein intake (g/kg/day)	0.91 \pm 0.17	0.83 \pm 0.29	0.47
Short Physical Performance Battery (SPPB)	9.0 \pm 1.4	8.9 \pm 2.2	0.91

Baseline correlations

A strong correlation was observed between leg lean mass and knee extension 1RM at baseline ($r = 0.80$, $P < 0.001$). In addition, 1RM strength normalized to body mass showed moderate correlations with habitual gait velocity ($r = 0.46$, $P < 0.05$) and stair climbing time ($r = -0.50$, $P < 0.05$), but not with five times chair-rise ($r = -0.25$, $P = 0.26$) (Table 3).

Table 3 Correlations at baseline for all participants (n = 22). Values for leg lean mass, vastus lateralis thickness, and knee extension 1RM were normalized to body weight prior to correlation analyses

Measure	1	2	3	4	5	6
1. Leg lean mass (kg)	—					
2. Vastus lateralis thickness (mm)	.29	—				
3. Knee extension 1RM (kg)	.80***	.22	—			
4. Habitual gait velocity (m/s)	.25	.13	.46*	—		
5. Five times chair rise (s)	-.02	.02	-.25	-.54**	—	
6. Stair climbing (s)	-.40	.01	-.50*	-.69***	.53*	—

* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

Adherence

All subjects in ST performed at least 90% of the training sessions in the ten-week period (95.8 ± 2.6), and all subjects in ST were therefore included in the per-protocol analysis. There were no correlations between adherence and training effect on any variable.

Protein intake

ST and CON had a protein intake of $0.91 (\pm 0.17)$ and $0.83 (\pm 0.29)$ $\text{g}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ at baseline, respectively. As a consequence of the protein supplement, ST increased their daily protein intake to $1.22 \text{ g}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ ($P = 0.002$). No significant change was observed in CON, and the group x time interaction only tended to be significant ($P = 0.058$) (Figure 2).

Fig. 2 Change in daily protein intake in ST (A) and CON (B). $**P < 0.01$, compared to baseline within group

Body composition

Body mass tended to increase from baseline in CON (1.1 ± 1.5 kg, $P = 0.056$), but not in ST (0.7 ± 1.5 kg, $P = 0.24$). No group x time interaction was observed ($P = 0.54$). However, a group x time interaction was observed for leg lean mass ($P = 0.001$), due to an increase from baseline in ST (0.7 ± 0.3 kg, $P < 0.001$), and no change in CON (0.1 ± 0.3 kg, $P = 0.69$) (Figure 3A). For fat mass, a tendency towards an increase was observed in CON (0.6 ± 0.7 kg; $P = 0.052$), whereas a non-significant reduction was observed in ST (-0.5 ± 0.9 kg, $P = 0.15$). Thus, for fat mass, a group x time interaction was observed ($P = 0.007$) (Figure 3B).

Fig. 3 Change in leg lean mass (A) and total body fat mass (B), in ST (●), CON (▲). $\#P < 0.05$, group x time interaction. $***P < 0.001$, compared to baseline within group. Results are presented as individual values and with mean and standard deviation (SD) for each group

Muscle thickness

In ST, increased thickness of vastus lateralis ($4.4 \pm 3.2\%$), rectus femoris ($6.7 \pm 5.1\%$), and vastus intermedius ($5.8 \pm 5.9\%$) were observed ($P < 0.001$ for all muscles) (Figure 4). Muscle thickness did not change in CON, and a group x time interaction was observed for all muscles investigated ($P < 0.01$ for all muscles). There were no differences between the relative change in vastus lateralis, rectus femoris and vastus intermedius in ST ($P = 0.38$). A strong correlation was observed between change in vastus lateralis thickness and the change in leg lean mass in the training group ($r = 0.67$, $P < 0.05$).

Fig. 4 Percentage changes in ST+PRO (●) and CON (▲), measured in vastus lateralis (A), rectus femoris (B), and vastus intermedius (C). # $P < 0.01$, group x time interaction. *** $P < 0.001$, compared to baseline within group. Results are presented as individual values and with mean and standard deviation (SD) for each group

Muscle strength

ST improved KE 1RM and isometric MVC by $23 \pm 15\%$ and $20 \pm 11\%$, respectively, and a group x time interaction was observed for both tests ($P < 0.001$) (Figure 5). The change in knee extension 1RM was not correlated with the change in leg lean mass, nor with the change in thickness of vastus lateralis, rectus femoris or vastus intermedius. Force at 100 milliseconds during the MVC improved from baseline only in ST ($22 \pm 22\%$), and a group x time interaction was observed ($P = 0.039$). Relative rate of torque development ($RTD_{max}/\text{maximal torque}$) did not change in either group.

Fig. 5 Percentage change in knee extension 1RM in ST (●) and CON (▲). # $P < 0.001$, group x time interaction. *** $P < 0.001$, compared to baseline within group. Results are presented as individual values and with mean and standard deviation (SD) for each group

Functional performance

ST participants used less time on five times chair-rise and stair climbing after the intervention, with improvements of 10 ± 13 ($P = 0.05$) and $11 \pm 12\%$ ($P = 0.034$) from baseline, respectively (Figure 6). A group x time interaction was observed for stair climbing ($P = 0.019$), but not five times chair-rise ($P = 0.13$). In addition, a group x time interaction was observed for habitual gait velocity ($P = 0.028$), due to a non-significant improvement in ST ($7 \pm 10\%$; $P = 0.17$), along with a non-significant decline in CON ($-5 \pm 14\%$, $P = 0.24$) (Figure 7). No significant correlations were observed between training-induced changes in knee extension 1RM and changes in five times chair-rise time ($r = -0.46$, $P = 0.15$) or stair climbing time ($r = 0.04$, $P = 0.89$). However, the change in knee extension 1RM tended to correlate with the change in habitual gait velocity ($r = 0.54$, $P = 0.09$). Furthermore, a strong correlation was observed between the change in isometric MVC and the change in habitual gait velocity in the training group ($r = 0.70$, $P = 0.015$) (Figure 8). No significant correlations were observed between the improvement in force at 100ms during MVC, and changes in functional performance.

Fig. 6 Percentage change in time used during five times chair rise (A) and stair climbing (B) in ST (●) and CON (▲). # $P < 0.05$, group x time interaction. * $P < 0.05$, compared to baseline within group. Results are presented as individual values and with mean and standard deviation (SD) for each group. Data were log-transformed prior to statistical analyses. The group x time interaction in B was significant ($P = 0.017$) even if the outlier in CON was excluded from the analysis

Fig. 7 Percentage change in habitual gait velocity in ST (●) and CON (▲). # $P < 0.05$, group x time interaction. Results are presented as individual values and with mean and standard deviation (SD) for each group. Data were log-transformed prior to statistical analyses

Fig. 8 Relationship between the change in knee extension MVC and habitual gait velocity in ST

DISCUSSION

Thirty minutes of heavy load strength training performed three times per week in combination with protein supplementation increased leg lean mass, as well as strength and functional capacity in mobility-limited older adults. Concerning hypertrophy of vastus lateralis, rectus femoris and vastus intermedius, we found no significant differences between the muscles. Although no correlations were observed between the changes in muscle mass/thickness and changes in muscle strength, we did observe correlations between strength gains and improvements in functional capacity.

Leg lean mass and muscle thickness

The intervention was successful in eliciting muscle hypertrophy, demonstrated both by the change in leg lean mass and muscle thickness. A meta-analysis of 49 studies, representing a total of 1328 participants, examined the effect of whole-body strength training on lean body mass (LBM) in individuals > 50 years. The mean intervention duration was 20.5 weeks, and the weighted pooled estimate of lean body mass change was 1.1 kg (12). Interestingly, the meta-analysis also found a negative association between age and change in lean mass. With a mean age of 86.6 years, the subjects in our experimental group (ST) can be categorized as *very old*, and thus an increase of 0.7 kg in leg lean mass after only ten weeks of leg exercise is somewhat higher than expected. Due to the design of our study, we are not able to conclude whether the protein supplement contributed to the observed effects. The literature is equivocal with regards to the importance of protein supplementation during strength training in healthy old individuals (3, 6, 13, 14), but there are reports of an additive effect in frail elderly (10). According to the diet assessments, the experimental group had a protein intake of $0.91 \text{ g}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ at baseline. Although this is above the Recommended Dietary Allowance (RDA) of $0.8 \text{ g}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ (15), it is below the Nordic recommendations of $1.2 \text{ g}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ for older adults (16). Furthermore, the variability in protein intake at baseline was quite high (range: $0.6\text{-}1.2 \text{ g}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$), and it is therefore possible that the protein supplement had additive effects at least in the individuals with a low protein intake at baseline.

Muscle thickness of vastus lateralis, rectus femoris and vastus intermedius increased by 4.4%, 6.7%, and 5.8%, respectively. In a systematic review, Stewart and coworkers investigated the hypertrophic response to strength training in very old individuals (> 75 years) (17). Three of four studies observed increased cross-sectional area (CSA) of quadriceps femoris, reporting improvements of 2-9%. Moreover, a recent study observed a 3.4% increase in quadriceps femoris CSA following twelve weeks of heavy-load strength training and protein supplementation in very old subjects (83-94 years) (7). As we measured muscle thickness and not cross-sectional area, we cannot directly compare our results with the above findings. However, a 7.5% increase in vastus lateralis thickness assessed by ultrasound corresponded to a 5.2% increase in CSA assessed by MRI following strength training in young individuals (18). Accordingly, our observation of 4-7% increases in muscle thickness corresponds well with previous reports (6, 7, 9, 19). No significant differences between vastus lateralis, rectus femoris and vastus intermedius were observed in the present investigation. To our knowledge, no study has investigated the relative hypertrophy in different compartments of quadriceps femoris in this population previously. In somewhat younger individuals (67 ± 5 years), increases of 6% and 16% were observed for vastus lateralis and rectus femoris, respectively, after 12 weeks of strength training (20). Also in young individuals, greater increases have been observed for rectus femoris (~18%), compared to both vastus lateralis (~8%) and vastus intermedius (~9%) (21). Although direct comparisons across studies should be performed with care, the overall greater improvements in muscle thickness in young (21), as well as in individuals in their late sixties (20), imply that the hypertrophic response to strength training may indeed be reduced at very high age, in line with other reports (12, 22, 23). Additionally, our results suggest that vastus lateralis, rectus femoris and vastus intermedius in relative terms contribute equally to the strength training induced increase in quadriceps femoris area in this population.

Muscle strength and rate of torque development

Knee extension 1RM increased by 23% in the training group. This is in the lower range of previous reports in similar populations (~25 to ~200%) (6, 7, 9, 24). In the present investigation, testing and training was performed in separate machines. This is likely advantageous in order to limit learning effects, which may have contributed to the substantial 1RM strength gains in some of the previous reports. This is supported by the often-observed discrepancy between increases in dynamic (1RM) and isometric (MVC) strength (7, 8). In contrast, dynamic and isometric strength gains were in good agreement in the present study, suggesting that the 23% improvement in knee extension 1RM was mainly due to an increase in the force-generating

capacity of the quadriceps femoris muscles, and not due to learning effects. However, in line with our hypothesis, strength gains still exceeded the increases in leg lean mass (4%) and muscle thickness (4-7%). The latter imply that additional factors contributed to the improvements in strength. In this regard, increased activation level, muscle density, muscle length and single fiber specific force may all have contributed (19, 25, 26). Although not significant, we did observe associations between the change in knee extension 1RM and habitual gait velocity ($r = 0.54$, $P = 0.09$) and time used during five times chair-rise ($r = -0.46$, $P = 0.15$). Additionally, the change in isometric MVC was significantly correlated with the change in habitual gait velocity ($r = 0.70$, $P < 0.05$). These relationships suggests that the extent to which strength gains are achieved, is decisive with respect to the degree of functional improvement in this population.

The improvements of knee extension 1RM and peak MVC were paralleled by improvement of torque at 100 ms during MVC. Torque at 100 ms represents the capacity of the muscle as a whole to produce force rapidly, and is therefore an absolute measure of rate of torque development (RTD). However, if torque at 100 ms were normalized to maximal torque, no effect of training was observed. Moreover, normalized RTD_{max} , given as the rise in torque development relative to maximal torque, did not improve. Hence, both these relative measures of RTD suggest that the intrinsic properties of the knee extensors were not affected. In theory, it would not be surprising if we had observed increased relative RTD as well, due to the preferential hypertrophy of type II-fibers with strength training (27-29). Nevertheless, increasing muscle strength alone will allow the muscle to produce more force over a short time interval, and thus improve the muscle's ability to produce power. This is important, since functional performance correlates even better with muscle power than muscle strength (30, 31). In addition, improved ability to produce force rapidly may prevent falls, due to an increased ability to correct unexpected imbalances that might easily occur during daily life (e.g. walking) (32).

Functional performance

The control subjects in the present investigation displayed a non-significant reduction (5%) in gait velocity over the 10-week period, indicating that functional declines in this population are rapid. It has been shown previously that gait velocity is associated with survival in older adults (33, 34), underscoring the importance of countermeasures. The 7% improvement observed in the training group is in agreement with two meta-analyses showing improvements of

approximately 5-10% following strength training in older adults (35, 36). The training group also improved five times chair-rise and stair climbing performance by 10 and 11%, respectively, again in accordance with previous reports (5, 6, 24, 37, 38). It has been suggested that high-speed power training has some advantages over traditional strength training with regards to improvements in function in this population (39). Nevertheless, our findings demonstrate that traditional heavy-load strength training effectively improves functional performance in mobility-limited older adults. Importantly, no adverse event occurred and adherence to the training was high, demonstrating that such training can be effectively and safely implemented.

Limitations

With only two groups, where the intervention group both performed strength training *and* received protein supplementation, we are not able to discern effects from training and effects from supplementation. Including a protein only and training only group would have provided this information. Moreover, intention-to-treat analyses were not performed in the present investigation. Although we aimed to recruit 26 subjects to account for dropouts, we were only able to recruit 22 participants. Luckily, as there were no dropouts, we still reached the number needed based on the power calculation. Another limitation of the study is that subjects and test personnel were not blinded (except ultrasound analysis). The lack of information about compliance to the supplement also represent a limitation of the study.

Conclusions

In line with our hypothesis, strength gains (20-23%) exceeded increases in leg lean mass (4%) and muscle thickness (4-7%) in mobility-limited older adults following heavy load strength training. We nevertheless hypothesized that the training-induced hypertrophy would correlate with increases in isometric strength, but no such correlations were observed. Collectively these findings indicate that other factors (e.g. activation level and muscle quality) contributed to the increases in strength. In line with our hypothesis, increases in muscle strength correlated with functional improvements, suggesting that interventions eliciting large improvements in strength may also be superior in terms of functional gains in this group vulnerable to frailty and functional decline.

Perspectives

Among older adults with mobility limitations living at home, many are at the verge of being institutionalized. Increasing functional capacity through strength training may allow these

individuals to become less dependent on assistance in various daily activities, to an extent where their activity level can be better maintained, and further loss of muscle mass and strength can be blunted or even prevented. Such strategies are important both from a socioeconomic and individual perspective. Future studies should address the mechanisms underlying the often-observed mismatch between improvements in muscle strength and size. A deeper understanding of such factors will strengthen the basis for the development of optimal countermeasures.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest regarding the publication of this article.

AUTHOR CONTRIBUTIONS

SA and TR contributed to the conception and design of the research; SA performed the experiments; SA analyzed the data; SA, OS, HB and TR interpreted the results of the experiments; SA prepared the figures; SA drafted the manuscript; SA, OS, HB and TR edited and revised the manuscript; SA, OS, HB and TR approved the final version of the manuscript.

ACKNOWLEDGEMENTS

The authors would like to thank Kristin Holte for analysis of the dietary registrations, Mauritz Kaashagen for performing the ultrasound measurements, and Anne Lene Nordengen for assistance during the supervision of training. We thank TINE SA for providing the nutritional supplements. We also thank our participants for their great contributions.

FUNDING

The study was funded by the Norwegian School of Sport Sciences.

REFERENCES

1. Hayashida I, Tanimoto Y, Takahashi Y, Kusabiraki T, Tamaki J. Correlation between muscle strength and muscle mass, and their association with walking speed, in community-dwelling elderly Japanese individuals. *PloS one*. 2014;9(11):e111810.
2. Vasunilashorn S, Coppin AK, Patel KV, Lauretani F, Ferrucci L, Bandinelli S, et al. Use of the Short Physical Performance Battery Score to predict loss of ability to walk 400 meters: analysis from the InCHIANTI study. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2009;64(2):223-9.
3. Cermak NM, Res PT, de Groot LC, Saris WH, van Loon LJ. Protein supplementation augments the adaptive response of skeletal muscle to resistance-type exercise training: a meta-analysis. *The American journal of clinical nutrition*. 2012;96(6):1454-64.
4. Verdijk LB, Jonkers RA, Gleeson BG, Beelen M, Meijer K, Savelberg HH, et al. Protein supplementation before and after exercise does not further augment skeletal muscle hypertrophy after resistance training in elderly men. *The American journal of clinical nutrition*. 2009;89(2):608-16.
5. Pinto RS, Correa CS, Radaelli R, Cadore EL, Brown LE, Bottaro M. Short-term strength training improves muscle quality and functional capacity of elderly women. *Age*. 2014;36(1):365-72.
6. Chale A, Cloutier GJ, Hau C, Phillips EM, Dallal GE, Fielding RA. Efficacy of whey protein supplementation on resistance exercise-induced changes in lean mass, muscle strength, and physical function in mobility-limited older adults. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2013;68(6):682-90.
7. Bechshoft RL, Malmgaard-Clausen NM, Gliese B, Beyer N, Mackey AL, Andersen JL, et al. Improved skeletal muscle mass and strength after heavy strength training in very old individuals. *Experimental gerontology*. 2017;92:96-105.
8. Harridge SD, Kryger A, Stensgaard A. Knee extensor strength, activation, and size in very elderly people following strength training. *Muscle & nerve*. 1999;22(7):831-9.
9. Fiatarone MA, O'Neill EF, Ryan ND, Clements KM, Solares GR, Nelson ME, et al. Exercise training and nutritional supplementation for physical frailty in very elderly people. *The New England journal of medicine*. 1994;330(25):1769-75.
10. Tieland M, Dirks ML, van der Zwaluw N, Verdijk LB, van de Rest O, de Groot LC, et al. Protein supplementation increases muscle mass gain during prolonged resistance-type exercise training in frail elderly people: a randomized, double-blind, placebo-controlled trial. *Journal of the American Medical Association*. 2012;307(8):713-9.
11. Guralnik JM, Ferrucci L, Simonsick EM, Salive ME, Wallace RB. Lower-extremity function in persons over the age of 70 years as a predictor of subsequent disability. *The New England journal of medicine*. 1995;332(9):556-61.
12. Peterson MD, Sen A, Gordon PM. Influence of resistance exercise on lean body mass in aging adults: a meta-analysis. *Medicine and science in sports and exercise*. 2011;43(2):249-58.
13. Thomas DK, Quinn MA, Saunders DH, Greig CA. Protein Supplementation Does Not Significantly Augment the Effects of Resistance Exercise Training in Older Adults: A Systematic Review. *Journal of the American Medical Association*. 2016;317(10):959 e1-9.

14. Finger D, Goltz FR, Umpierre D, Meyer E, Rosa LH, Schneider CD. Effects of protein supplementation in older adults undergoing resistance training: a systematic review and meta-analysis. *Sports medicine*. 2015;45(2):245-55.
15. Volpi E, Campbell WW, Dwyer JT, Johnson MA, Jensen GL, Morley JE, et al. Is the optimal level of protein intake for older adults greater than the recommended dietary allowance? *The journals of gerontology Series A, Biological sciences and medical sciences*. 2013;68(6):677-81.
16. Nordic Nutrition Recommendations. Copenhagen: Nordic Council of Ministers; 2012.
17. Stewart VH, Saunders DH, Greig CA. Responsiveness of muscle size and strength to physical training in very elderly people: a systematic review. *Scandinavian journal of medicine & science in sports*. 2014;24(1):e1-10.
18. Franchi MV, Longo S, Mallinson J, Quinlan JJ, Taylor T, Greenhaff PL, et al. Muscle thickness correlates to muscle cross-sectional area in the assessment of strength training-induced hypertrophy. *Scandinavian journal of medicine & science in sports*. 2018;28(3):846-53.
19. Sipilä S, Suominen H. Effects of strength and endurance training on thigh and leg muscle mass and composition in elderly women. *Journal of applied physiology*. 1995;78(1):334-40.
20. Bjørnsen T, Salvesen S, Berntsen S, Hetlelid KJ, Stea TH, Lohne-Seiler H, et al. Vitamin C and E supplementation blunts increases in total lean body mass in elderly men after strength training. *Scandinavian journal of medicine & science in sports*. 2016;26(7):755-63.
21. Ema R, Wakahara T, Miyamoto N, Kanehisa H, Kawakami Y. Inhomogeneous architectural changes of the quadriceps femoris induced by resistance training. *European journal of applied physiology*. 2013;113(11):2691-703.
22. Greig CA, Gray C, Rankin D, Young A, Mann V, Noble B, et al. Blunting of adaptive responses to resistance exercise training in women over 75y. *Experimental gerontology*. 2011;46(11):884-90.
23. Raue U, Slivka D, Minchev K, Trappe S. Improvements in whole muscle and myocellular function are limited with high-intensity resistance training in octogenarian women. *Journal of applied physiology*. 2009;106(5):1611-7.
24. Seynnes O, Fiatarone Singh MA, Hue O, Pras P, Legros P, Bernard PL. Physiological and functional responses to low-moderate versus high-intensity progressive resistance training in frail elders. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2004;59(5):503-9.
25. Arnold P, Bautmans I. The influence of strength training on muscle activation in elderly persons: a systematic review and meta-analysis. *Experimental gerontology*. 2014;58:58-68.
26. Frontera WR, Hughes VA, Krivickas LS, Kim SK, Foldvari M, Roubenoff R. Strength training in older women: early and late changes in whole muscle and single cells. *Muscle & nerve*. 2003;28(5):601-8.
27. Verdijk LB, Gleeson BG, Jonkers RA, Meijer K, Savelberg HH, Dendale P, et al. Skeletal muscle hypertrophy following resistance training is accompanied by a fiber type-specific increase in satellite cell content in elderly men. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2009;64(3):332-9.

28. Kosek DJ, Kim JS, Petrella JK, Cross JM, Bamman MM. Efficacy of 3 days/wk resistance training on myofiber hypertrophy and myogenic mechanisms in young vs. older adults. *Journal of applied physiology*. 2006;101(2):531-44.
29. Kryger AI, Andersen JL. Resistance training in the oldest old: consequences for muscle strength, fiber types, fiber size, and MHC isoforms. *Scandinavian journal of medicine & science in sports*. 2007;17(4):422-30.
30. Reid KF, Fielding RA. Skeletal muscle power: a critical determinant of physical functioning in older adults. *Exercise and sport sciences reviews*. 2012;40(1):4-12.
31. Bean JF, Kiely DK, Herman S, Leveille SG, Mizer K, Frontera WR, et al. The relationship between leg power and physical performance in mobility-limited older people. *Journal of the American Geriatrics Society*. 2002;50(3):461-7.
32. Pijnappels M, Reeves ND, Maganaris CN, van Dieen JH. Tripping without falling; lower limb strength, a limitation for balance recovery and a target for training in the elderly. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology*. 2008;18(2):188-96.
33. White DK, Neogi T, Nevitt MC, Peloquin CE, Zhu Y, Boudreau RM, et al. Trajectories of gait speed predict mortality in well-functioning older adults: the Health, Aging and Body Composition study. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2013;68(4):456-64.
34. Studenski S, Perera S, Patel K, Rosano C, Faulkner K, Inzitari M, et al. Gait speed and survival in older adults. *JAMA : the journal of the American Medical Association*. 2011;305(1):50-8.
35. Van Abbema R, De Greef M, Craje C, Krijnen W, Hobbelen H, Van Der Schans C. What type, or combination of exercise can improve preferred gait speed in older adults? A meta-analysis. *BMC geriatrics*. 2015;15:72.
36. Hortobagyi T, Lesinski M, Gabler M, VanSwearingen JM, Malatesta D, Granacher U. Effects of Three Types of Exercise Interventions on Healthy Old Adults' Gait Speed: A Systematic Review and Meta-Analysis. *Sports medicine*. 2015;45(12):1627-43.
37. Schot PK, Knutzen KM, Poole SM, Mrotek LA. Sit-to-stand performance of older adults following strength training. *Research quarterly for exercise and sport*. 2003;74(1):1-8.
38. Dias CP, Toscan R, de Camargo M, Pereira EP, Griebler N, Baroni BM, et al. Effects of eccentric-focused and conventional resistance training on strength and functional capacity of older adults. *Age*. 2015;37(5):99.
39. Sayers SP, Gibson K. High-speed power training in older adults: a shift of the external resistance at which peak power is produced. *Journal of strength and conditioning research / National Strength & Conditioning Association*. 2014;28(3):616-21.