

Hingst, J. R., Bjerre, R. D., Wojtaszewski, J. F. P., Jensen, J. (2019). Rapid radiochemical filter paper assay for determination of hexokinase activity and affinity for glucose-6-phosphate. *Journal of applied physiology*, 127(3), 661-667. <http://dx.doi.org/10.1152/jappphysiol.00207.2018>

Dette er siste tekst-versjon av artikkelen, og den kan inneholde små forskjeller fra forlagets pdf-versjon. Forlagets pdf-versjon finner du her:
<http://dx.doi.org/10.1152/jappphysiol.00207.2018>

This is the final text version of the article, and it may contain minor differences from the journal's pdf version. The original publication is available here:
<http://dx.doi.org/10.1152/jappphysiol.00207.2018>

1 **A rapid radiochemical filter paper assay for determination of Hexokinase activity and affinity for**
2 **glucose-6-phosphate (G6P)**

3

4 Janne R. Hingst¹, Rie D. Bjerre¹, Jørgen F. P. Wojtaszewski¹, Jørgen Jensen^{2,*}

5

6

7

8 ¹ Section of Molecular Physiology, Department of Nutrition, Exercise and Sports, Faculty of Science,
9 University of Copenhagen, DK-2100 Copenhagen, Denmark

10 ² Department of Physical Performance, Norwegian School of Sports Sciences, Oslo, Norway

11

12

13

14 * To whom correspondence should be addressed. Department of Physical Performance, Norwegian
15 School of Sport Sciences, Sognsveien 220, PO Box 4014 Ullevål Stadion, 0806 Oslo, Norway.

16 e-mail: jorgen.jensen@nih.no

17

18

19 *Abstract*

20 Glucose phosphorylation by hexokinase (HK) is a rate-limiting step in glucose metabolism. Regulation
21 of HK includes feedback inhibition by its product glucose-6-phosphate (G6P) and mitochondria
22 binding. HK affinity for G6P is difficult to measure because its natural product (G6P) inhibits enzyme
23 activity. HK phosphorylates several hexoses and we have taken advantage of the fact that 2-
24 deoxyglucose-6-phosphate (2DG-6P) does not inhibit HK activity. By this we have developed a new
25 method for rapid radiochemical analysis of HK activity with 2-deoxyglucose (2-DG) as substrate,
26 which allows control of the concentrations of G6P to investigate HK affinity for inhibition by G6P. We
27 verified that 2DG serves as a substrate for the HK reaction with linear time and concentration
28 dependency as well as expected V_{max} and K_M . This is the first simple assay evaluating feedback
29 inhibition of HK by its product G6P and provides a unique technique for future research evaluating the
30 regulation of glucose phosphorylation under various physiological conditions.

31

32 *New & Noteworthy*

33 Traditionally, hexokinase activity has been analyzed spectrophotometrically where the product
34 formation of G6P is analyzed by an indirect reaction coupled to NADPH formation during conversion
35 of G6P to 6-P gluconolactone. By nature this assay prevents measurements of HK affinity for inhibition
36 by G6P. We have developed a rapid radiochemical filter paper assay to study HK affinity for G6P by
37 use of radiolabeled 2-DG as substrate, to study physiological regulation of HK affinity for G6P-
38 induced inhibition.

39

40 *Introduction.* Glucose uptake in various tissues is regulated by glucose delivery, glucose transport
41 across the membrane and intracellular phosphorylation (32). Although glucose phosphorylation can be
42 limiting for glucose uptake under various physiological conditions (2), the regulation of glucose
43 phosphorylation has received surprisingly little attention. Hexokinase (HK) catalyzes the irreversible
44 reaction converting glucose to glucose 6-phosphate (G6P). Glucose phosphorylation is a potential rate-
45 limiting step for glucose metabolism and Mammals express four different isoforms of hexokinase
46 (hexokinase I-IV) (37) while skeletal muscle expresses HK I and II (6, 11). HK is inhibited
47 allosterically by G6P with a K_i for G6P in the physiological range (37). It has been known for many
48 years that HK binds to mitochondria and it was more recently reported that HK II is phosphorylated by
49 PKB at Thr⁴⁷³ (26). The translocation of HK to the mitochondria reduces the ability of G6P to inhibit
50 enzyme activity. However, the fact that the product inhibits HK activity makes it difficult to study the
51 physiological regulation of HK by G6P.

52 HK activity can easily be measured spectrophotometrically or fluorometrically (28). In this
53 method G6P-dehydrogenase (G6P-DH) in the assay buffer converts the formed G6P to 6-
54 Phosphogluconolactone in a reaction coupled to conversion of NADP⁺ to NADPH; production of
55 NADPH is measured to describe HK activity (28). Inclusion of G6P-DH in these methods is required
56 for the formation of NADPH, the presence of G6P-DH also prevents the accumulation of G6P which
57 otherwise would exert a negative feedback on HK enzyme activity. Radiolabeled glucose or ATP can
58 also be used to measure HK activity where glucose and G6P can easily be separated. Sanderson et al.
59 used ¹⁴C-labelled glucose to measure HK activity and excluded G6P-DH from the assay buffer. In this
60 study the authors demonstrated less inhibition of HK activity as G6P accumulated after insulin
61 stimulation in skeletal muscles from lean but not from obese Zucker rats (28). This suggests that insulin

62 decreases HK affinity for G6P-induced inhibition, but the method does not allow to calculate K_i
63 (inhibitory constant that is reflective of the binding affinity) for G6P-induced inhibition of HK. These
64 data suggest that HK affinity for G6P is regulated in skeletal muscles and highlights that methodologies
65 to investigate HK affinity for G6P are required.

66 Hexokinase phosphorylates a number of hexoses like 2-deoxyglucose (2-DG), but only G6P
67 and a few other hexose 6-phosphates inhibit HK activity (36). Importantly, 2-deoxyglucose-6-
68 phosphate (2-DG-6P) does not inhibit HK activity (37). Thus, 2-DG serves as ideal substrate for HK
69 activity assay where the sensitivity for inhibition by G6P can be studied. Here we present a novel
70 radiochemical filter paper assay for rapid determination of HK activity and affinity for inhibition by
71 G6P. This method may not only be applicable for muscle physiology but also for cancer research as the
72 malignant phenotype of cancer cells is associated with increased hexokinase expression and glycolytic
73 flux (20).

74

75 **Materials and methods**

76 Experiments were conducted in Norway and Denmark. Experiments in Norway were approved by
77 Norwegian authorities and experimental procedures in Denmark by Danish and Animal Experimental
78 Inspectorate. All experiments were conducted in compliance with the European Union convention for
79 protection of vertebrate animals used for scientific purposes (Council of Europe 123, Strasbourg, France,
80 1985).

81 *Animals.* In Norway male Wistar rats were purchased from B & K Universal (Nittedal, Norway). In
82 Denmark male Wistar rats and female C57B6 mice were purchased from Taconic (Ejby, Denmark).

83 Animals were kept in a room with 12:12-h light-dark cycle (light on/off, 6:00 AM/6:00 PM) and had
84 free access to water and chow.

85 *Tissue preparation and muscle incubation.* Animals were anesthetized by intraperitoneal injection of
86 pentobarbital sodium (10 mg/100 g body wt) and tissues were dissected. Tissue samples were
87 homogenized directly (fresh tissue). In some studies, epitrochlearis and soleus were contracted or
88 treated with insulin *in vitro* prior to analysis of HK activity. Information about experimental conditions
89 is given in figure and table legends. For *in vitro* studies, muscles were dissected and suspended at
90 approximate resting length in an incubation system containing Krebs-Henseleit buffer (KHB) as
91 previously described (1). After 30 min preincubation, muscles were either rested for 30 min, stimulated
92 by supraphysiological insulin concentration (10 mU/ml) or forced to contract by electrical stimulation.
93 Muscle contraction was induced by electrical stimulation for 30 min consisting of 200-ms trains (100
94 Hz, 0.2 ms pulse duration, 10 V) delivered every 2 s, a protocol that previously has been shown to
95 result in a robust increase in glucose uptake (1). During the entire incubation period the buffer was
96 maintained at 30°C and oxygenated with 95% O₂ and 5% CO₂. After incubation, muscles were
97 harvested, blotted on filter paper and immediately homogenized.

98 *Homogenization.* Tissue was homogenized in ice cold buffer containing 250 mM sucrose, 20 mM
99 Sodium-pyrophosphate, 5 mM HEPES (pH 7.4), 5 mM magnesium chloride, 1 mM DTT, 5% Dextran
100 70, 20 mM β-glycerophosphate, 10 mM sodium fluoride, 2 mM PMSF, 1 mM EGTA (pH 8.0), 10
101 μg/ml Aprotinin, 10 μg/ml Leupeptin, 2 mM sodium orthovanadate and 3 mM benzamidine for 2 x 30
102 s at 30 Hz using a TissueLyzer II (Qiagen, Hilden, Germany).
103 Homogenates were analyzed for total protein concentration by use of the bicinchoninic acid method
104 (Thermo Scientific, Waltham, MA, USA).

105 A detailed list providing product no. and stock concentrations for reagents used in the homogenization
106 buffer is given in Supplemental I ([dx.doi.org/10.17504/protocols.io.xcifue](https://doi.org/10.17504/protocols.io.xcifue)).

107

108 *Filter paper HK activity assay.* Experiments were initially performed in Eppendorf tubes and the
109 reaction buffer with homogenate was spotted on DE81 anion exchanger filter paper (Whatman, UK,
110 #3685 915) and washed as described below. To optimize and make the assay more efficient, the assay
111 was changed to 96-wells PCR plates (Thermo Fisher Scientific, AB0600). Homogenates were diluted
112 to ~ 1-2 mg protein/ml in homogenization buffer. 15 µl sample was added to a 96-well PCR plate
113 placed on an ice cold aluminum plate. As controls 15 µl homogenization buffer, cold assay buffer
114 (buffer blank), a standard pool and a blank sample (sample blank) was added to every assay plate.
115 Kinase activity was started by adding 15 µl assay buffer in 1 sec intervals with a multichannel pipette
116 while the aluminum plate was placed on a 30°C heating block. The assay buffer contained 50 mM TEA
117 (pH 7.4), 5 mM ATP, 5 mM magnesium chloride, 10 mM creatine phosphate, 2.4 µCi/ml ¹⁴C-2-
118 deoxyglucose or ³H-2-deoxyglucose (Perkin Elmer) , 2 mM 2DG, 0.05% BSA, 7 U/ml creatine kinase.
119 In case HK activity was evaluated in the presence of G6P or 2DG6P, various concentrations were
120 added to the assay buffer (see results for details). Kinase reaction was stopped at 30 °C after 15-25 min
121 (depending on tissue and protein amount in the reaction) by adding 30 µl (1 vol) 90 % EtOH. The PCR
122 plate was kept under gentle agitation (manual agitation) at room temperature for 30 sec to ensure full
123 mixture and hence complete stop of kinase reaction. 20 µl were spotted in duplicates by a multichannel
124 pipette on DE81 filter paper. The filter paper was subsequently washed 3 x 20 min in 300 ml 60%
125 EtOH in a plastic box (31cm x 22 cm x 6 cm) to remove non-phosphorylated 2DG and dried in a fume
126 hood. In order to determine specific activity, assay buffer (2 x 1 µl and 2 x 2 µl) was spotted on the

127 dried filter paper and added in vials containing 4 ml scintillation fluid. Finally the dried DE81 paper
128 was wrapped in plastic wrap and placed on a STORM cassette for 48 hours and analyzed on a STORM
129 Phosphoimager. Hexokinase activity was calculated as amount of ^{14}C -2DG6 or ^3H -2-DG6 captured in
130 the DE81 filter paper and given relative to specific activity of the assay buffer. A step-by-step
131 description of the assay is given in Supplemental II ([dx.doi.org/10.17504/protocols.io.xckfiuw](https://doi.org/10.17504/protocols.io.xckfiuw)).

132 *Comparison of fluorometric and radiochemical assay for determination of HK activity.*

133 Tissue was homogenized as described above and analyzed for HK activity by the new radiochemical
134 filter paper assay and compared to the well-established fluorometric method.

135 For fluorometric determination of HK activity, 20 μl sample was loaded as triplicate on a microtiter
136 plate and 200 μl reaction mixture was added. The reaction mixture contained 200 mM Tris buffer (Tris
137 HCl, Calbiochem 648313 and Tris Base, Calbiochem 648311) (pH 8.0), 10 mM magnesium chloride
138 (Sigma M2670), 12.5 mM Dithiothreitol (DTT) (Sigma D0362), 25 mM Glucose (Sigma G8270), 1
139 mM NADP, 11 U/ml G6PDH. The reaction was started by adding 10 μl ATP (100mM) (Sigma A
140 3377) and absorbance was measured at 340 nm as described by Lowry & Passoneau (18).

141 *Effect of exercise training on HK activity:* Quadriceps muscle samples from female WT mice
142 (SV129/C57B6 mixed background) from a previous study (9) were investigated for HK activity by use
143 of the new radiochemical filter paper assay and compared to the fluorometric method. Briefly, mice
144 were housed in single cages for 4 weeks without (sedentary) or with (training) access to a running
145 wheel. Mice were euthanized by cervical dislocation and muscle tissue was frozen in liquid nitrogen
146 and stored at $-80\text{ }^{\circ}\text{C}$. We have previously investigated these samples by western blotting and reported a
147 ~ 3 fold increase in training-induced HKII protein expression (9).

148 *Results.*

149 In the new radiochemical filter paper assay glucose phosphorylated by HK to G6P will be
150 isolated by binding to the filter paper (Figure 1). Initially we compared the new radiochemical filter
151 paper assay to the established fluorometric method for determination of HK activity (Table 1).
152 Immediately after euthanization, rat brain, heart, soleus muscle and extensor digitorum longus (EDL)
153 muscle were homogenized. One portion of the homogenate was analyzed by the fluorometric assay
154 while another portion from the same homogenate sample was analyzed by the new radiochemical filter
155 paper assay. This enabled for direct comparison between these methods and revealed that the new filter
156 paper assay yields comparable results to the established fluorometric assay (Table 1).

157 Based on multiple analyses intra-assay variation and inter-assay variation were calculated to be
158 5-10% and 10-16%, respectively. These were found to be comparable to other enzymatic assay such as
159 the GS-activity assay. Collectively these data demonstrate that the filter paper assay shows sensitivity
160 for detection of radiolabeled G6P that is comparable to the established fluorometric assay.

161 The fact that HK activity is inhibited by its product G6P precludes the use of glucose as
162 substrate for determination of HK affinity for G6P. However, hexokinase phosphorylates a number of
163 hexoses that do not inhibit HK activity and hence may serve as candidates to study alterations in
164 affinity of HK activity for product inhibition by G6P (37). Therefore, we investigated 2-DG as a
165 substrate for the filter paper assay and compared it to glucose (Figure 2A). HK activity in rat heart was
166 assayed in the presence of increasing concentrations (0, 0.025, 0.05, 0.1, 0.2, 0.5, 1, 2 and 5 mM) of 2-
167 DG and glucose, respectively (Figure 2A). Kinetic parameters were calculated by Lineweaver Burk
168 Plots (Figure 2B-C) and are given in Table 2. Hexokinase enzyme reaction velocity was substantially

169 higher using 2-DG compared to glucose with a maximal velocity (V_{\max}) of 14 and 6 $\mu\text{mol/g}$
170 protein/min, respectively. K_M values derived from Lineweaver-Burk plot revealed a higher value for 2-
171 DG compared to glucose (0.67 vs. 0.16 mM). These findings do not preclude kinetic studies of HK
172 activity with 2-DG as a substrate; instead the markedly higher HK activity in light of a higher K_M
173 demonstrate that 2-DG serves as optimal substrate to study kinetic properties of HK. Furthermore,
174 product formation (2-DG6P) in rat heart increased linearly in a time- and concentration-dependent
175 manner (Figure 3A-B), validating the experimental settings of the radiochemical filter paper assay with
176 2-DG as a substrate.

177 Regular exercise training leads to a substantial increase in HK II expression and activity (35). In
178 order to verify that the new radiochemical filter paper assay is capable to detect alterations in HK
179 expression, we analyzed muscle sample from a previous mouse exercise training study (9).
180 Determination of HK by the fluorometric and filter paper assay revealed a comparable increase in HK
181 activity by exercise training (~2.7 fold increase for fluorometric assay and ~2.5 fold increase for the
182 filter paper assay) (Figure 3C). These data are in accordance with the markedly higher levels of HK
183 expression we previously have reported (9) and demonstrate that the filter paper assay shows sensitivity
184 for detecting differences in HK activity.

185 Given that majority of previous studies have investigated HK activation in frozen tissue, we
186 tested whether the radiochemical filter paper assay also can be used to investigate HK activity in frozen
187 tissue (Figure 3D). Rat soleus muscles were split in to two portions; one being homogenized as frozen
188 tissue and the other part being homogenized as fresh tissue. Homogenate from frozen and fresh tissue
189 was analyzed for HK activity by the fluorometric assay and the new filter paper assay. These data
190 demonstrate that the radiochemical filter paper assay can be used to investigate HK activity in frozen

191 tissue, albeit the HK activity generally tended to be a few percent lower in homogenate from frozen
192 tissue. Previous studies have reported HK binding to mitochondria as a potential mechanism for
193 reducing affinity for G6P-mediated inhibition (20, 26). In an attempt to maintain this interaction we
194 decided to perform subsequent analyses on fresh tissue.

195 Next, we studied HK activity in the presence of increasing concentrations (0, 0.1, 0.2, 0.5 and
196 1 mM) of 2-DG6P and G6P (Figure 4). This experiment reveals that increasing levels of G6P inhibit
197 HK activity with a 50 % reduction (K_i) at ~0.2 mM G6P, while a similar increase in 2-DG6P in the
198 assay buffer has no effect on HK activity. These findings verify that radiolabeled 2-DG can be used as
199 substrate without that the formed 2-DG6P exerts a negative feedback on HK activity.

200 When assayed in the presence of increasing G6P concentrations, the physiological regulation of HK
201 affinity for G6P may be detected. Insulin and muscle contraction serve as main physiological stimuli to
202 increase muscle glucose uptake and a subsequent rise in intracellular G6P levels (2, 14). Despite
203 elevated G6P levels skeletal glucose flux increases, suggesting that the negative feedback G6P exerts
204 on HK activity is antagonized under these physiological conditions. In order to investigate whether
205 sensitivity of HK for G6P underlies physiological regulation rat epitrochlearis muscles were stimulated
206 *ex vivo* by insulin (10 mU/ml) or forced to contract for 30 min by electrical stimulation (Figure 5).

207 Maximal hexokinase activity (~1.8 $\mu\text{mol/g protein/min}$) was observed when assayed in the absence of
208 G6P with no effect of insulin stimulation or muscle contraction. HK activity decreased with increasing
209 G6P levels in a similar pattern in all three conditions and with a half-maximal inhibition (K_i) around
210 0.2 mM G6P. Thus, under the present experimental conditions HK affinity for G6P remained
211 unaffected by insulin stimulation and muscle contraction.

212

213

214 *Discussion*

215 Hexokinase is the classical example of an enzyme that is inhibited by its product (G6P) in a feedback
216 dependent manner. However, regulation of HK activity is complex and HK translocates to
217 mitochondria and becomes phosphorylated (26). Translocation of HK to mitochondria has obtained
218 large interest, but the effect of this translocation on HK activity is poorly understood. There are
219 indications that translocation of HK to the mitochondria reduces affinity for G6P-induced inhibition,
220 but such analyses are hampered by the fact that HK is inhibited by its natural product G6P (37). Other
221 proteins are regulated by G6P; e.g. the major regulatory mechanism for glycogen synthase (GS) is by
222 changes in affinity for activation by G6P. This occurs via phosphorylation/dephosphorylation
223 mechanisms (15, 17, 25). GS activation is normally reported as fractional activity, where activity at a
224 physiological concentration of G6P is related to total activity measured with a high concentration of
225 G6P sufficient to activate GS completely (12). Our idea was to establish a comparable assay where HK
226 activities are measured without G6P (total activity) and with a physiological concentration of G6P to
227 investigate changes in HK affinity for G6P-induced inhibition. Therefore, we have developed a simple
228 and rapid method for measurement of G6P-induced inhibition of HK by using 2-deoxyglucose (2-DG)
229 as substrate to avoid inhibition by the product. This method can be used to evaluate regulation of HK
230 activity in various cell types and under various experimental conditions.

231 Here we present a rapid radiochemical filter paper assay that reveals HK activity measurements
232 with a linear dose and time dependency comparable to the well-established fluorometric assay.
233 Moreover, the use of radiolabeled 2-DG as a substrate in the filter paper assay allows to study the

234 sensitivity of HK for inhibition by its product G6P. We could demonstrate that increasing levels of G6P
235 but not 2-DG6P inhibit HK activity, providing a model to study the regulation of HK sensitivity for
236 G6P under various physiological settings.

237 Intracellular G6P serves as central regulator for skeletal muscle glycogen synthesis and should
238 be considered together with the degree of dephosphorylation of glycogen synthase when evaluating the
239 rate of glycogen synthesis (30). On the other hand G6P exerts a negative feedback on HK and the
240 intracellular concentration of G6P must therefore be balanced between activation of glycogen synthase
241 and inhibition of HK (19). We observed a half maximal inhibition (K_i) of HK activity at ~0.2 mM G6P,
242 suggesting that HK was potently inhibited by levels of G6P thought to exist in skeletal muscle *in vivo*.
243 Thus, previous studies have reported G6P concentrations within the range of 0.2 – 0.3 mM under
244 resting conditions and increase above 1 mM in response to stimulation by insulin and muscle
245 contraction (14, 19, 23, 24). The G6P concentrations applied in the current HK assay (0.1-1 mM)
246 therefore reflect what could be expected in the physiological range. Increased glucose uptake and a
247 subsequent rise in intracellular G6P associated with insulin stimulation and muscle contraction should
248 consequently increase inhibition of hexokinase further, yet an increased flux is observed *in vivo* (2, 4,
249 27). This suggests the presence of mechanisms that uncouple the negative feedback associated G6P
250 levels and HK activity *in vivo*. In contrast to our hypothesis, HK sensitivity for G6P in response to
251 insulin stimulation and muscle contractions remained unaffected under the experimental conditions in
252 the present study. Sanderson et al (28) have previously reported that insulin reduced G6P-induced
253 inhibition in skeletal muscles, and other studies have proposed association of HK with mitochondrial
254 membrane as a mechanism to regulate HK activity (3, 21, 37). Thus, once HK I and II are bound to
255 outer mitochondrial membrane inhibition of enzyme activity by G6P is antagonized. It has been

256 reported that insulin stimulation (3, 37) and muscle contraction (13, 33) increase binding of HK II to
257 the mitochondrial fraction. We used a homogenization buffer (containing 5% dextran 70 and 5 mM
258 MgCl₂) and a procedure (brief homogenization of fresh tissue) that previously has been shown to
259 maintain the integrity of the mitochondria (7, 8, 28, 34). Nevertheless we were unable to observe a
260 regulation of HK inhibition in response to insulin stimulation or muscle contraction. Insulin and
261 exercise have been shown to increase expression of HKII in human skeletal muscle whereas HKI
262 remains largely unaffected in response to these stimuli (16, 31). Notably this isoform specific
263 regulation was associated with increased HKII activity in soluble but not in the particulate fraction,
264 highlighting the importance subcellular regulation when investigating the regulation of HK activity. In
265 the present study HK enzyme activity was not investigated isoform specific in cellular sub-fractions, so
266 we cannot rule out the possibility that regulation of HKII activity may have been overseen due to the
267 crude measurement in the present study.

268 Hexokinase binds to the mitochondria (VDAC) via its N-terminal end and mutating the fifth
269 amino acid histidine to proline completely abolishes HK-2 binding to the mitochondria (5). The
270 interaction between HK2 and VDAC is not characterized, but it is tempting to speculate that the binding
271 is rather loose and may be disrupted during homogenization. However, mitochondria can be isolated
272 with attached HK (22) and it could therefore be speculated that isolation of mitochondrial fraction is
273 required to study these alterations in G6P-mediated inhibition of HK activity. Indeed, Van Houten and
274 colleagues reported an increase in HK activity in the mitochondrial fraction but not total fraction when
275 rat gastrocnemius muscles were analyzed following swimming exercise until exhaustion (13). We
276 suggest that other homogenization procedures or studies in saponised muscle bundles should be
277 investigated.

278 The interaction between HKII and mitochondria (VDAC) has large interest in cancer research
279 and emerged as a drug target for cancer therapies (29) and disruption of HKII-mitochondria interaction
280 with peptides corresponding to 15 aa of the N-terminal of HKII disrupt the interaction and triggers
281 apoptosis (38). The mechanism by which this disruption triggers apoptosis remain unknown and we
282 believe assaying HK affinity for G6P-induced inhibition will be most useful to solve this question.
283 HKII binding to mitochondria has also been suggested as key aspect in cardiac protection (10). It has
284 been demonstrated in cardiomyocytes that HK is phosphorylated on Thr⁴⁷³ by PKB, a regulatory
285 mechanisms that was found to be associated with translocation of HK to the mitochondria and was
286 associated with reduced affinity for G6P-mediated inhibition (26). This adds an additional regulatory
287 mechanism for control of HK inhibition by G6P, although the presence in intact mature tissue such as
288 skeletal muscle remains to be demonstrated. A simple method to measure HK affinity for G6P-induced
289 inhibition will be a useful tool to address such question.

290 This is the first assay that takes advantage of 2-DG as substrate to study the sensitivity of HK
291 activity for G6P. Abnormal regulation of HK activity has previously been reported in disease state such
292 as muscle insulin resistance in type 2 diabetes (28) or the increased glycolytic flux present in malignant
293 cancer cells (20). The presented method provides a valuable tool to study the regulation of HK
294 inhibition by G6P under these conditions. Future research could therefore apply this assay in order to
295 clarify the physiological regulation of HK affinity in diseased states such as T2D and cancer.

296

297 Funding sources: The study was funded by grants provided by The Lundbeck Foundation (R183-2014-
298 3990/ R180-2014-3887 to JFPW) and The Novo Nordisk Foundation (NNF 16OC0023046, NNF

299 170OC00272224 to JFPW). Jørgen Jensen received a visiting professorship by Danish Diabetes
300 Academy (DDA) funded by Novo Nordisk.

301 *Acknowledgement:* The authors acknowledge the samples from the mouse training study (9) provided
302 by J. Fentz, Section of Molecular Physiology, Department of Nutrition, Exercise and Sports, Faculty of
303 Science, University of Copenhagen.

304

305 *References*

- 306 1. **Aslesen R, Engebretsen EML, Franch J, Jensen J.** Glucose uptake and metabolic stress in rat
307 muscles stimulated electrically with different protocols. *J Appl Physiol* 91: 1237–1244, 2001.
- 308 2. **Aslesen R, Jensen J.** Effects of epinephrine on glucose metabolism in contracting rat skeletal
309 muscles. *Am J Physiol* 275: E448–E456, 1998.
- 310 3. **Bessman S, Geiger P.** Compartmentation of hexokinase and creatine phosphokinase, cellular
311 regulation, and insulin action. *Curr Top Cell Regul* 16: 55–86, 1980.
- 312 4. **Bouskila M, Hirshman MF, Jensen J, Goodyear LJ, Sakamoto K.** Insulin promotes glycogen
313 synthesis in the absence of GSK3 phosphorylation in skeletal muscle. *Am J Physiol Endocrinol*
314 *Metab* 294: E28-35, 2008.
- 315 5. **Bryan N, Raisch K.** Identification of a Mitochondrial-Binding Site on the Amino-Terminal End
316 of Hexokinase II. *Biosci. Rep.* (2015). doi: 10.1042/BSR20150047.
- 317 6. **Chang P, Jensen J, Printz R, et al.** Overexpression of Hexokinase II in Transgenic Mice. *J*
318 *Biol Chem* 271: 14834–14839, 1996.

- 319 7. **Cogswell AM, Stevens RJ, Hood DA.** Properties of skeletal muscle mitochondria from
320 subsarcolemmal and intermyofibrillar isolated regions. *Am J Physiol* 264: C383-389, 1993.
- 321 8. **Elander A, Sjöström M, Lundgren F, Scherstén T, Bylund-Fellenius AC.** Biochemical and
322 morphometric properties of mitochondrial populations in human muscle fibres. *Clin Sci (Lond)*
323 69: 153–64, 1985.
- 324 9. **Fentz J, Kjøbsted R, Kristensen CM, Hingst JR, Birk JB, Gudiksen A, Foretz M,**
325 **Schjerling P, Viollet B, Pilegaard H, Wojtaszewski JFP.** AMPK α is essential for acute
326 exercise-induced gene responses but not for exercise training-induced adaptations in mouse
327 skeletal muscle. *Am J Physiol - Endocrinol Metab* 309: ajpendo.00157.2015, 2015.
- 328 10. **Halestrap AP, Pereira GC, Pasdois P.** The role of hexokinase in cardioprotection - Mechanism
329 and potential for translation. *Br J Pharmacol* 172: 2085–2100, 2015.
- 330 11. **Halseth AE, Bracy DP, Wasserman DH.** Overexpression of hexokinase II increases insulin-
331 and exercise-stimulated muscle glucose uptake in vivo. *Am J Physiol Endocrinol Metab* 276:
332 E70-7, 1999.
- 333 12. **Højlund K, Beck-Nielsen H.** Impaired glycogen synthase activity and mitochondrial
334 dysfunction in skeletal muscle: markers or mediators of insulin resistance in type 2 diabetes?
335 *Curr Diabetes Rev* 2: 375–395, 2006.
- 336 13. **Van Houten DR, Davis JM, Meyers DM, Durstine JL.** Altered cellular distribution of
337 hexokinase in skeletal muscle after exercise. *Int J Sports Med* 13: 436–438, 1992.
- 338 14. **Jensen J, Jebens E, Brennesvik EO, Ruzzin J, Soos M a, Engebretsen EML, O’Rahilly S,**

- 339 **Whitehead JP.** Muscle glycogen inharmoniously regulates glycogen synthase activity, glucose
340 uptake, and proximal insulin signaling. *Am J Physiol Endocrinol Metab* 290: E154–E162, 2006.
- 341 15. **Kolnes AJ, Birk JB, Eilertsen E, Stuenæs JT, Wojtaszewski JFP, Jensen J.** Epinephrine-
342 stimulated glycogen breakdown activates glycogen synthase and increases insulin-stimulated
343 glucose uptake in epitrochlearis muscles. *Am J Physiol - Endocrinol Metab* 308: E231–E240,
344 2015.
- 345 16. **Koval JA, DeFronzo RA, O RM, Printz R, Ardehali H, Granner DK, Mandarino LJ,**
346 **DeFronzo RA, O’Doherty RM, Printz R, Ardehali H, Granner DK, Mandarino LJ.**
347 Regulation of hexokinase II activity and expression in human muscle by moderate exercise. *Am*
348 *J Physiol - Endocrinol Metab* 274: 304–308, 1998.
- 349 17. **Lai Y-C, Zarrinpashneh E, Jensen J.** Additive effect of contraction and insulin on glucose
350 uptake and glycogen synthase in muscle with different glycogen contents. *J Appl Physiol* 108:
351 1106–15, 2010.
- 352 18. **Lowry OH, Passonneau JV.** *A Flexible System of Enzymatic Analysis.* London: Academic
353 Press, Inc., 1972.
- 354 19. **Lueck JD, Fromm HJ.** Kinetics, mechanism, and regulation of rat skeletal muscle hexokinase.
355 *J Biol Chem* 249: 1341–1347, 1974.
- 356 20. **Mathupala SP, Ko YH, Pedersen PL.** Hexokinase II: cancer’s double-edged sword acting as
357 both facilitator and gatekeeper of malignancy when bound to mitochondria. *Oncogene* 25: 4777–
358 86, 2006.

- 359 21. **Mohan C, Geiger P, Bessman S.** The intracellular site of action of insulin: the mitochondrial
360 Krebs cycle. *Curr Top Cell Regul* 30: 105–42, 1989.
- 361 22. **Pasdois P, Parker JE, Halestrap AP.** Extent of mitochondrial hexokinase II dissociation
362 during ischemia correlates with mitochondrial cytochrome c release, reactive oxygen species
363 production, and infarct size on reperfusion. *J Am Heart Assoc* 2: 1–20, 2013.
- 364 23. **Piras R, Staneloni R.** In Vivo Regulation of Rat Muscle Glycogen Synthetase Activity.
365 *Biochemistry* 8: 2153–2160, 1969.
- 366 24. **Richter EA, Hansen SA, Hansen BF.** Mechanisms limiting glycogen storage in muscle during
367 prolonged insulin stimulation. *Am J Physiol* 255: E621–E628, 1988.
- 368 25. **Roach PJ, Larner J.** Covalent phosphorylation in the regulation of glycogen synthase activity.
369 *Mol Cell Biochem* 15: 179–200, 1977.
- 370 26. **Roberts DJ, Tan-Sah VP, Smith JM, Miyamoto S.** Akt phosphorylates HK-II at Thr-473 and
371 increases mitochondrial HK-II association to protect cardiomyocytes. *J Biol Chem* 288: 23798–
372 23806, 2013.
- 373 27. **Rothman DL, Shulman RG, Shulman GI.** 31P nuclear magnetic resonance measurements of
374 muscle glucose-6-phosphate: Evidence for reduced insulin-dependent muscle glucose transport
375 or phosphorylation activity in non-insulin-dependent diabetes mellitus. *J Clin Invest* 89: 1069–
376 1075, 1992.
- 377 28. **Sanderson a L, Radda GK, Leighton B.** Abnormal regulation of hexokinase in insulin-
378 resistant skeletal muscle. *Biochem Mol Med* 59: 80–6, 1996.

- 379 29. **Shoshan-Barmatz V, Krelin Y, Shteinfer-Kuzmine A, Arif T.** Voltage-Dependent Anion
380 Channel 1 As an Emerging Drug Target for Novel Anti-Cancer Therapeutics. *Front Oncol* 7,
381 2017.
- 382 30. **Shulman RG, Bloch G, Rothman DL.** In vivo regulation of muscle glycogen synthase and the
383 control of glycogen synthesis. *Proc Natl Acad Sci U S A* 92: 8535–42, 1995.
- 384 31. **Vogt CC, Ardehali HH, Iozzo PP, Yki-Järvinen HH, Koval JJ, Maezono KK, Pendergrass**
385 **MM, Printz RR, Granner DD, DeFronzo RR, Mandarino LL.** Regulation of hexokinase II
386 expression in human skeletal muscle in vivo. *Metabolism* 49: 814–818, 2000.
- 387 32. **Wasserman DH, Kang L, Ayala JE, Fueger PT, Lee-Young RS.** The physiological regulation
388 of glucose flux into muscle in vivo. *J Exp Biol* 214: 254–262, 2011.
- 389 33. **Weber FE, Pette D.** Changes in free and bound forms and total amount of hexokinase isozyme
390 II of rat muscle in response to contractile activity. *Eur J Biochem* 191: 85–90, 1990.
- 391 34. **Wicker U, Bücheler K, Gellerich FN, Wagner M, Kapischke M, Brdiczka D.** Effect of
392 macromolecules on the structure of the mitochondrial inter-membrane space and the regulation
393 of hexokinase. *BBA - Bioenerg* 1142: 228–239, 1993.
- 394 35. **Wilson JE.** Regulation of mammalian hexokinase activity. *Regul. Carbohydr. Metab.* .
- 395 36. **Wilson JE.** Hexokinases. *Rev Physiol Biochem Pharmacol* 126: 65–198, 1995.
- 396 37. **Wilson JE.** Isozymes of mammalian hexokinase: structure, subcellular localization and
397 metabolic function. *J Exp Biol* 206: 2049–2057, 2003.

398 38. **Woldetsadik AD, Vogel MC, Rabeh WM, Magzoub M.** Hexokinase II-derived cell-
399 penetrating peptide targets mitochondria and triggers apoptosis in cancer cells. *FASEB J* 31:
400 2168–21874, 2017.

401

402

403 **Figure legends**

404 **Figure 1: Assay reaction by the new established radiochemical filter paper assay that was**
405 **developed to study hexokinase sensitivity for inhibition by G6P.**

406 A rapid radiochemical filter paper assay for measurement of hexokinase (HK) activity and affinity for
407 G6P was developed. This assay allows for the simple separation of phosphorylated form of the
408 radiolabeled tracer ($[^{14}\text{C}]$ - Glucose-phosphate or $[^3\text{H}]$ -2-Deoxyglucose-phosphate) from the
409 unphosphorylated form ($[^{14}\text{C}]$ - Glucose and $[^3\text{H}]$ -2-deoxyglucose).

410 In this assay conversion of radiolabeled 2-Deoxyglucose (2DG) to 2-Deoxyglucose-6-phosphate (2-
411 DG-6-phosphate) is catalyzed by HK. 2DG-6-phosphate binds to DE81 ion-exchange filter paper
412 (Whatman, Maidstone, UK) and non-phosphorylated 2-DG can be separated by washing the filter paper
413 in excess 60% EtOH. Importantly, 2-DG-6-phosphate does not inhibit HK activity and 2-DG is
414 therefore a perfect substrate for studying the HK sensitivity for G6P. Thus, HK affinity can easily be
415 investigated by performing the radiochemical filter paper assay in the presence of varying G6P
416 concentration.

417

418 **Figure 2: Comparison of substrates for radiochemical filter paper assay to study hexokinase**
419 **activity.** Hexokinase activity in 6 homogenates derived from a single rat heart was measured in the
420 presence of increasing concentration (0, 0.025, 0.05, 0.1, 0.2, 0.5, 1, 2 and 5 mM) of glucose for ^{14}C -
421 glucose and 2-deoxyglucose (2-DG) for ^3H -2-deoxyglucose, respectively (A). Data are representing
422 means and variation (SEM) for 6 homogenates obtained from the same heart tissue. Lineweaver-Burk
423 Plots were used to calculate V_{\max} and K_M for ^3H -2-DG and ^{14}C -Glucose (B-C) and are given in Table 2.

424

425 **Figure 3. Sensitivity of the radiochemical filter paper assay**

426 Hexokinase activity in a single rat heart homogenate was measured over time (A) and in the presence
427 of increasing sample/protein concentration for 2 min (B) by use of the radiochemical filter paper assay
428 with 2-DG as substrate and ^3H -2DG as tracer. Product formation of 2-DG-6P was calculated. Each data
429 point represents a triplicate.

430 The effect of exercise training on HK activity was investigated in mouse muscle from a previous study
431 (9). Quadriceps muscle from sedentary mice and 4-week trained mice (access to running wheel) were
432 compared by the fluorometric and the radiochemical filter paper assay (C). Data from 4 sedentary and 4
433 trained mice are presented as means \pm SEM (n=4). For direct comparison between fresh and frozen
434 tissue, rat soleus muscles were split into two pieces and homogenized as fresh and as frozen tissue
435 respectively (D). HK activity was subsequently determined by fluorometric method and the
436 radiochemical filter paper assay (C). Data represent means \pm SEM for 4 soleus muscles that were split
437 into two pieces and investigated by by fluorometric method and the radiochemical filter paper assay
438 (n=4). T-test within fluorometric assay and filter paper assay was used to investigate for the effect of
439 exercise training and fresh/frozen muscle respectively. *** $p < 0.001$ for effect of exercise training.

440 **Figure 4. Hexokinase activity in the presence of increasing 2DG-6P and G6P concentrations.**

441 Hexokinase activity in mouse heart was analyzed by the radiochemical filter paper assay with ^{14}C -2DG
442 as a substrate and assayed in the presence of increasing 2-DG6P or G6P concentrations (0, 0.1, 0.2, 0.5,
443 1 mM), respectively. Data represent heart samples from 8 mice and are given as means \pm SEM (n=8).

444

445 **Figure 5. Physiological regulation hexokinase sensitivity for G6P-mediated inhibition.** Rat
446 epitrochlearis muscles were isolated and suspended in an *ex vivo* incubation system. Muscles were
447 rested (basal non-stimulated control), stimulated by insulin (10 mU/ml) or forced to contract for 30min.
448 Immediately after incubation, muscles were homogenized and analyzed for HK activity. Sensitivity of
449 HK activity for G6P was analyzed with use radiochemical filter paper assay with ¹⁴C-2DG as a
450 substrate assayed in the presence of increasing G6P concentrations (0, 0.1, 0.2, 0.5, 1 mM).
451 Data represent values from 8 rested control muscle, 8 insulin-stimulated muscles and 8 electrically
452 contracted muscles and are given as means+/-SEM (n=8).

453 **Table 1: Comparison of fluorometric and radiochemical assay for hexokinase activity.**

454 Rat brain, heart, soleus muscle and EDL muscle were homogenized immediately after cervical
455 dislocation. HK activity in homogenate samples was determined by the established fluorometric assay
456 (formation of NADH) and compared to the newly established radiochemical filter paper assay (¹⁴C-
457 2DG 6-phosphate). Activity is given as μmol/g protein/min.
458 Data present HK activity in different tissues from 8 Wistar rats (n=8) and are presented as means and
459 the variation among samples is given as SEM (n=8). T-tests revealed no statistical differences between
460 fluorometric and radiochemical assay for measurement of HK activity.

461 **Table 2: Kinetic parameters for 2-deoxyglucose (2DG) and Glucose**

462 Lineweaver-Burk Plots were used to calculate V_{max} and K_M for ³H-2-DG and ¹⁴C-Glucose (Figure 2B-
463 C) and are given in Table 2.

464

Figure 1

Assay reaction





Table 1

| | Fluorometric assay | Filter paper assay |
|---------------|---------------------------|---------------------------|
| Brain | 28.76 ± 2.87 | 26.01 ± 1.31 |
| Heart | 10.32 ± 1.04 | 11.63 ± 1.68 |
| Soleus | 2.31 ± 0.05 | 2.21 ± 0.17 |
| EDL | 1.23 ± 0.14 | 1.14 ± 0.29 |

Figure 2


A

Hexokinase activity - Comparison of substrates



B

³H-2DG: Lineweaver-Burk plot



C

¹⁴C-Glucose: Lineweaver-Burk plot





Table 2: Kinetic parameters for 2DG and Glucose


| | V _{max} (μmol/g protein/min) | K _M (mM) |
|-------------------------|--|------------------------|
| ³ H-2DG | 14 | 0.67 |
| ¹⁴ C-Glucose | 6 | 0.16 |

Figure 3:


A



B



C



D




Figure 4




Figure 5

