

Sagelv, E. H., Ekelund, U., Hopstock, L. A., Fimland, M. S., Løvsletten, O., Wilsgaard, T., Morseth, B. (2021) The bidirectional associations between leisure time physical activity change and body mass index gain. The Tromsø Study 1974–2016. *International Journal of Obesity*, 45(9), 1830-1843. <http://dx.doi.org/10.1038/s41366-021-00853-y>

---

Dette er siste tekst-versjon av artikkelen, og den kan inneholde små forskjeller fra forlagets pdf-versjon. Forlagets pdf-versjon finner du her: <http://dx.doi.org/10.1038/s41366-021-00853-y>

---

This is the final text version of the article, and it may contain minor differences from the journal's pdf version. The original publication is available here: <http://dx.doi.org/10.1038/s41366-021-00853-y>

---

1     **The bidirectional associations between leisure time physical activity change and body**  
2                                   **mass index gain. The Tromsø Study 1974-2016**

3     Edvard H Sagelv<sup>1\*</sup>, Ulf Ekelund<sup>2,3</sup>, Laila A Hopstock<sup>4</sup>, Marius Steiro Fimland<sup>5,6</sup>, Ola  
4     Løvsletten<sup>4</sup>, Tom Wilsgaard<sup>4</sup>, Bente Morseth<sup>1</sup>

5     **Affiliations**

6     <sup>1</sup>School of Sport Sciences, Faculty of Health Sciences, UiT The Arctic University of Norway,  
7     Tromsø, Norway

8     <sup>2</sup>Department of Sports Medicine, Norwegian School of Sport Sciences, Oslo, Norway

9     <sup>3</sup>Department of Chronic Diseases and Ageing, Norwegian Institute of Public Health, Oslo,  
10    Norway

11    <sup>4</sup>Department of Community Medicine, Faculty of Health Sciences, UiT The Arctic University  
12    of Norway, Tromsø, Norway

13    <sup>5</sup>Department of Neuromedicine and Movement Science, Faculty of Medicine and Health  
14    Sciences, Norwegian University of Science and Technology, Trondheim, Norway

15    <sup>6</sup>Unicare Helsefort Rehabilitation Centre, Rissa, Norway

16    \*Corresponding author

17    Edvard H Sagelv

18    **E-mail:** [edvard.h.sagelv@uit.no](mailto:edvard.h.sagelv@uit.no)

19    **Phone:** +47 77660236

20    **Twitter:** @edvardhsagelv

21    **Address:** UiT The Arctic University of Norway, Postboks 6050 Langnes, 9037 Tromsø,  
22    Troms, Norway

23    **Competing interests:** The authors declare that they have no competing interests.

24    **Word count:** 3797

25

26 **Abstract**

27 **Objectives:** To examine whether leisure time physical activity changes predict subsequent  
28 body mass index (BMI) changes, and conversely, whether BMI changes predict subsequent  
29 leisure time physical activity changes.

30 **Methods:** This prospective cohort study included adults attending  $\geq 3$  consecutive Tromsø  
31 Study surveys (time: T1, T2, T3) during 1974-2016 (N=10779). If participants attended  $> 3$   
32 surveys, we used the three most recent surveys. We computed physical activity change  
33 (assessed by the Saltin-Grimby Physical Activity Level Scale) from T1 to T2, categorized as  
34 Persistently Inactive (n=992), Persistently Active (n=7314), Active to Inactive (n=1167) and  
35 Inactive to Active (n=1306). We computed BMI change from T2 to T3, which regressed on  
36 preceding physical activity changes using analyses of covariance. The reverse association  
37 (BMI change from T1 to T2 and physical activity change from T2 to T3; n=4385) was  
38 assessed using multinomial regression.

39 **Results:** Average BMI increase was  $0.86 \text{ kg/m}^2$  (95% CI: 0.82 to 0.90) from T2 to T3. With  
40 adjustment for sex, birth year, education, smoking and BMI at T2, there was no association  
41 between physical activity change from T1 to T2 and BMI change from T2 to T3 (Persistently  
42 Inactive:  $0.89 \text{ kg/m}^2$  (95% CI: 0.77 to 1.00), Persistently Active:  $0.85 \text{ kg/m}^2$  (95% CI: 0.81 to  
43  $0.89$ ), Active to Inactive:  $0.90 \text{ kg/m}^2$  (95% CI: 0.79 to 1.00), Inactive to Active  $0.85 \text{ kg/m}^2$   
44 (95% CI: 0.75 to 0.95),  $p=0.84$ ). Conversely, increasing BMI was associated with Persistently  
45 Inactive (odds ratio (OR): 1.17, 95% CI: 1.08 to 1.27,  $p<0.001$ ) and changing from Active to  
46 Inactive (OR: 1.16, 95% CI: 1.07 to 1.25,  $p<0.001$ ) compared with being Persistently Active.

47 **Conclusion:** We found no association between leisure time physical activity changes and  
48 subsequent BMI changes, whereas BMI change predicted subsequent physical activity  
49 change. These findings indicate that BMI change predicts subsequent physical activity change  
50 at population level and not *vice versa*.

51 **Keywords;** <sup>1</sup>occupational physical activity, <sup>2</sup>obesity, <sup>3</sup>overweight, <sup>4</sup>adiposity, <sup>5</sup>longitudinal,  
52 <sup>6</sup>prospective, <sup>7</sup>energy expenditure, <sup>8</sup>energy balance

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

## 72 **INTRODUCTION**

73 The prevalence of overweight and obesity is continuously growing worldwide where now  
74 over 50% of the population is classified as either overweight or obese in western high-income  
75 countries (1). As overweight and obesity is associated with a substantial increased risk of non-  
76 communicable diseases and premature death (2), it is one of the greatest threats to public  
77 health in western high-income countries (1, 2).

78

79 Weight gain prevention at populational level is challenging. Obviously, excessive adiposity  
80 and weight gain are effects of an imbalanced energy intake and expenditure (3). Thus,  
81 increasing physical activity levels could potentially serve as an effective public health strategy  
82 to prevent population weight gain (3, 4). However, studies examining whether population  
83 levels of physical activity can prevent weight gain show conflicting results (4, 5), which may  
84 be attributed to methodological issues (5). For example, although current weight is a strong  
85 predictor of future weight gain (5), some studies failed to adjust for baseline weight or body  
86 mass index (BMI) (6-14). Moreover, most studies did not take the temporal reciprocal  
87 relationship between changing physical activity and weight into account (5), as they assessed  
88 the association between baseline physical activity level and future weight or BMI change (9,  
89 10, 15-22). Other studies examined the associations between change scores in both physical  
90 activity and weight or BMI (6, 8, 11-14, 23-37), which basically are cross-sectional analyses  
91 of change scores (5). Finally, the association between physical activity and weight change  
92 may be reverse as weight change may lead to physical activity change (5, 6, 22, 25, 26, 35,  
93 37-39), or this may be bidirectional (5).

94

95 Furthermore, the association between physical activity change and BMI change may be  
96 modified by other behavioural or societal factors, including sex (10), age (17, 18, 28-30),

97 smoking (23, 29), education (24), physical activity domain (*e.g.* occupation or leisure time)  
98 (29), and baseline BMI (28, 29). However, these observations are not consistent (12, 15, 26,  
99 33), which warrant further investigation.

100

101 Declines in both leisure time and occupational physical activity may contribute to population  
102 BMI gains (4). We have previously reported on the association between occupational physical  
103 activity and BMI change in a population-based cohort (The Tromsø Study) from Norway  
104 followed through repeated examinations every ~6 years. Our results suggested that  
105 occupational physical activity declines did not contribute to population BMI gains (40). As  
106 large proportions of the population are inactive during work hours (41, 42), leisure time  
107 physical activity may have greater potential to prevent weight gain. Thus, the aims of this  
108 study were to assess: 1) Whether changes in leisure time physical activity from examination 1  
109 (time (T) 1) to the next (T2) predicted subsequent changes in BMI from T2 to the next  
110 examination (T3), and 2) Whether BMI changes from T1 to T2 predicted subsequent leisure  
111 time physical activity changes from T2 to T3, with ~6 years follow up between each  
112 examination.

113

## 114 **MATERIALS AND METHODS**

### 115 **Design**

116 The study design is illustrated in Figure 1. We studied participants from the Tromsø Study, a  
117 population-based cohort study in Tromsø Municipality, Norway. There are seven repeated  
118 Tromsø Study surveys (attendance of invited participants=%): 1974 (Tromsø 1; 83%), 1979-  
119 80 (Tromsø 2; 85%), 1986-87 (Tromsø 3; 81%), 1994-95 (Tromsø 4; 77%), 2001 (Tromsø 5;  
120 79%), 2007-08 (Tromsø 6; 66%) and 2015-16 (Tromsø 7; 65%). Invited participants were

121 selected from total birth cohorts and random samples of inhabitants in Tromsø municipality  
122 (41, 43). Only men were invited to Tromsø 1 (1974), while in Tromsø 2-7 (1979-2016) both  
123 men and women were invited (details described elsewhere (41, 43)). In this study, we  
124 included participants attending at least three consecutive surveys (hereafter called T1, T2,  
125 T3). To assess the association between change in physical activity from T1 to T2 and change  
126 in BMI from T2 to T3, the inclusion criteria were information on: 1) physical activity at T1  
127 and T2, and height and weight at T2 and T3; 2) information on educational level and smoking  
128 habits at T2; and 3) not pregnant at T2 and/or T3. We also reversed the analyses to assess  
129 whether BMI change from T1 to T2 predicted physical activity change from T2 to T3. Here,  
130 inclusion criteria were: 1) height and weight at T1 to T2, physical activity at T2 to T3; 2)  
131 educational level and smoking habits at T2; and 3) not pregnant at T1 and/or T2. If the  
132 participants attended more than three consecutive surveys, we used their data from the three  
133 most recent surveys in the analyses of the overall cohort, while their data could be included in  
134 multiple period-specific samples (Tromsø 1-3: 1974-1987, Tromsø 2-4: 1979-1995, Tromsø  
135 5-7: 2001-2016).

136

137 **Insert Figure 1 about here.**

138

## 139 **Participants**

140 Participant selection for our analyses is illustrated in Figure 2. The overall cohort comprised  
141 10779 participants, which derive from the participants' three most recent Tromsø Study  
142 attendances. We also created period-specific samples where each participant may be included  
143 in multiple period-specific samples: Tromsø 1-3 (1974-1987, n=3598), Tromsø 2-4 (1979-  
144 1995, n=9691) and Tromsø 5-7 (2001-2016, n=2206). Therefore, the period-specific samples

145 do not add up to the overall cohort, which only includes participants with their three most  
146 recent consecutive surveys (Figure 2).

147

148 **Insert Figure 2 about here.**

149

150 The reversed analyses (BMI change from T1 to T2 followed by physical activity change from  
151 T2 to T3) were assessed in an overall cohort comprising 4385 participants (Figure 3). The  
152 leisure time physical activity questionnaire was not included in Tromsø 4 and only those <70  
153 years answered the questionnaire in Tromsø 5; this explains the lower sample size in the  
154 reversed analyses compared with the main analyses.

155

156 **Insert Figure 3 about here.**

157

158 All participants from Tromsø 4-7 provided written informed consent and the present study  
159 was approved by the Regional Ethics Committee for Medical Research (ref. 2016/758410).

160

### 161 **Self-reported physical activity**

162 Physical activity was measured with the Saltin-Grimby Physical Activity Level Scale  
163 (SGPALS) questionnaire (44, 45), which asks participants to rank their physical activity by  
164 four hierarchical levels for leisure- and occupational time physical activity, separately, during  
165 the last 12 months (44). The SGPALS in the Tromsø Study is slightly modified compared to  
166 the original by Saltin and Grimby (44) (Supplementary Table 1). The SGPALS is found to  
167 provide acceptable reliability (45) and validation studies have demonstrated acceptable ability  
168 to rank physical activity level when evaluated against accelerometry and cardiorespiratory  
169 fitness as the criteria (45).



170

171 Physical activity change was computed as 1) *Persistently Inactive* (reporting rank 1 at T1 and  
172 T2; n=992); 2) *Persistently Active* (rank  $\geq 2$  at T1 and T2; n=7314); 3) *Active to Inactive* (rank  
173  $\geq 2$  at T1 and rank 1 at T2; n=1167); and 4) *Inactive to Active* (rank 1 at T1 and rank  $\geq 2$  at T2;  
174 n=1306).

175

176 The leisure time SGPALS was used in all Tromsø Study surveys except Tromsø 4 (1994-95),  
177 and in Tromsø 5 (2001) not by those  $\geq 70$  years. The occupational time SGPALS was used in  
178 all surveys by participants of all ages.

179

## 180 **Body mass index and weight**

181 Weight and height were measured in light clothing and are expressed as kilograms (kg) and  
182 meters (m). BMI was calculated as  $\text{kg/m}^2$  and categorized into normal weight ( $< 25 \text{ kg/m}^2$ ),  
183 overweight ( $25\text{-}29 \text{ kg/m}^2$ ) and obese ( $\geq 30 \text{ kg/m}^2$ ) for stratified analyses. Change in BMI from  
184 T2 to T3 was calculated with height being fixed at T2 and change in BMI from T1 to T2 fixed  
185 at T1, to avoid a possible effect of height loss between the measurements.

186

## 187 **Confounders and effect modifiers**

188 We selected sex, birth year, smoking, education and BMI measured at T2 as confounders, and  
189 we also assessed potential effect modification of the confounders in addition to occupational  
190 physical activity change from T1 to T2. Data on smoking, education and occupational  
191 physical activity were retrieved from questionnaires. We categorized smoking into 1) Current  
192 smoker, 2) Previous smoker, and 3) Never smoker. The participants reported years of  
193 education in Tromsø 2 (1979-80), Tromsø 3 (1986-87) and Tromsø 5 (2001), which we  
194 categorized into 1) Primary school ( $< 10$  years), 2) High school (10-12 years), 3) University

195 <4 years (13-15 years), and 4) University  $\geq 4$  years ( $\geq 16$  years). In Tromsø 4 (1994-95) and  
196 Tromsø 6 (2007-08), the participants reported education with five response options, which  
197 included the four abovementioned groups and a fifth named “Technical school 2 years senior  
198 high” (vocational training), which we categorized as 2) High school.

199

## 200 **Availability of data and materials**

201 The data that support the findings of this study are available from the Tromsø Study but  
202 restrictions apply to the availability of these data, which were used under license for the  
203 current study, and so are not publicly available. The data are however available from the  
204 Tromsø Study upon application to the Data and Publication Committee for the Tromsø  
205 Study: [tromsous@uit.no](mailto:tromsous@uit.no).

206

## 207 **Statistical Analyses**

208 Paired t-tests were used to assess whether participants changed BMI from T2 to T3. Analyses  
209 of covariance (ANCOVA) were used to assess the association between physical activity  
210 change from T1 to T2 and BMI change from T2 to T3, adjusted for sex, birth year, smoking,  
211 education and BMI at T2. The ANCOVA was applied on the overall cohort and the period-  
212 specific samples, in total and stratified by sex, birth year, smoking, education and  
213 occupational physical activity change from T1 to T2. We interpreted the Q-Q plots of BMI  
214 change from T2 to T3 to not deviate from normal distribution. Although the Levene’s test of  
215 equality variance violated the assumption of homogeneity of variance across physical activity  
216 change groups ( $p < 0.001$ ), we considered our large sample size in all physical activity change  
217 groups to make the ANCOVA robust for this heterogeneity. Interaction effects were tested  
218 between physical activity change and potential effect modifiers (sex, birth year, smoking,  
219 education and BMI at T2, and occupational time physical activity change from T1 to T2) in

220 the overall cohort. We performed sensitivity analyses with leisure time physical activity  
221 change categorized into 6 groups; 1) *Persistently Inactive*, 2) *Persistently Active*, 3) *Active but*  
222 *decreasing* (rank 4 or 3 → 3 or 2), 4) *Active and Increasing* (rank 2 or 3 → 3 or 4), 5) *Active*  
223 *to Inactive* and 6) *Inactive to Active*. Alpha was set to 0.05 and data are shown as mean and  
224 95% confidence intervals (CI) from t-tests and ANCOVAs.

225

226 We performed multinomial logistic regressions to estimate odds ratios (OR) with 95%  
227 confidence intervals (CI) for changing leisure time physical activity from T2 to T3 per unit  
228 BMI change from T1 to T2, adjusted for sex, birth year, smoking and education at T2. The  
229 analyses were performed in the overall sample (n=4385) and stratified by sex, birth year,  
230 smoking, education and occupational physical activity change (T1 to T2). We assessed  
231 interaction effects between BMI change and potential effect modifiers (sex, birth year,  
232 smoking, education and BMI at T2, and occupational physical activity change from T1 to T2).  
233 *Persistently Active* was set as reference category. We used the Statistical Package for Social  
234 Sciences (SPSS, Version 26, IBM, Armonk, NY, United States) for all statistical analyses.

235

## 236 **RESULTS**

237 The descriptive characteristics at T2 for the overall cohort and period-specific samples  
238 (Tromsø 1-3, 1974-1987; Tromsø 2-4, 1979-1995; Tromsø 5-7, 2001-2016) are presented in  
239 Table 1. The participants increased their BMI from T2 to T3 (all  $p < 0.001$ ), except for the  
240 Tromsø 5-7 sample ( $p = 0.96$ ).

241

242 **Insert Table 1 about here**

243 **Change in BMI by preceding change in leisure time physical activity**

244 Changes in BMI by preceding leisure time physical activity change are presented in Table 2,  
245 and BMI at T2 and T3 by leisure time physical activity change are presented in  
246 Supplementary Table 2. In the overall cohort, we observed no differences in BMI change  
247 between categories of leisure time physical activity change ( $p=0.84$ ), and in general no  
248 associations in strata by sex, birth year, smoking, education and occupational physical activity  
249 change (Table 2).

250

251 **Insert Table 2 about here**

252

253 In the period-specific sample Tromsø 5-7 (2001-2016), we observed differences in BMI  
254 change between the leisure time physical activity change groups, where those changing from  
255 Active to Inactive increased their BMI more than those changing from Inactive to Active  
256 ( $p=0.01$ ). In stratified analyses, higher BMI change was observed in those changing from  
257 Active to Inactive in men ( $p=0.02$ ) but not in women ( $p=0.22$ ), and among those born  $\leq 1949$   
258 ( $p=0.05$ ). In those who never smoked, Persistently Inactive participants decreased their BMI  
259 more than those changing from Active to Inactive ( $p=0.03$ ). Finally, there were differences  
260 among the leisure time physical activity change groups among those having  $<4$  years  
261 university education; those changing from Active to Inactive increased their BMI more than  
262 all other leisure time physical activity change groups ( $p=0.003$ ) (Supplementary Table 3).

263 There were no differences in BMI increase by leisure time physical activity change in the  
264 Tromsø 1-3 (1974-1987) and Tromsø 2-4 (1979-1995) samples (Supplementary Table 4-5).

265

266 In the overall cohort, we observed no interaction for the association between leisure time  
267 physical activity change and BMI change by sex ( $p=0.62$ ), birth year ( $p=0.23$ ), smoking

268 (p=0.08) or BMI (p=0.44) at T2, or occupational physical activity change from T1 to T2  
269 (p=0.10). However, we observed that education modified the association between leisure time  
270 physical activity change and BMI change (p=0.002).

271

272 In the sensitivity analyses (in the overall cohort), with six physical activity change groups, the  
273 results were similar (Supplementary Table 6).

274

### 275 **Change in leisure time physical activity with preceding BMI change**

276 Participants who increased their BMI from T1 to T2 were more likely to be Persistently  
277 Inactive (OR: 1.17, 95% CI: 1.08 to 1.27 per 1 unit BMI (kg/m<sup>2</sup>)-increase, p<0.001) and to  
278 change from Active to Inactive (OR: 1.16, 95% CI: 1.07 to 1.25, p<0.001) from T2 to T3  
279 compared with those being Persistently Active at T2 and T3. Increasing BMI was not  
280 associated with changing from Inactive to Active (OR: 1.01, 95%CI: 0.94 to 1.08, p=0.97)  
281 compared with those being Persistently Active at T2 and T3 (Table 3).

282

### 283 **Insert Table 3 about here**

284

285 Sex, birth year, BMI, smoking and education at T2, and occupational physical activity change  
286 from T1 to T2, all modified the associations between BMI change and subsequent leisure time  
287 physical activity change (Table 3). Stratified analyses showed slight differences in ORs  
288 between strata. For example, men were more likely to be Persistently Inactive than  
289 Persistently Active per BMI-unit increase, while this was not observed in women. Those in  
290 higher birth year strata (1940-49, ≥1950) were more likely to be Persistently Inactive or  
291 changing from Active to Inactive with increasing BMI, which was not observed in those born  
292 ≤1939 (Table 3).

293

## 294 **DISCUSSION**

295 In this prospective cohort study, we found no association between leisure time physical  
296 activity changes and subsequent BMI changes, whereas BMI increases predicted subsequent  
297 low and decreasing physical activity levels.

298

299 Most previous studies assessing the prospective association between leisure time physical  
300 activity and BMI either used baseline physical activity as the predictor (9, 10, 15-23, 30),  
301 which do not take temporal changes between physical activity and BMI into account (5), or  
302 assessed associations between change scores for both physical activity and BMI (6, 8, 11-14,  
303 23-37), which basically are cross-sectional analyses that cannot provide an indication of the  
304 direction of the association (5). One study examined the association between physical activity  
305 changes from a 1<sup>st</sup> to a 2<sup>nd</sup> examination and BMI changes from the 1<sup>st</sup> to a 3<sup>rd</sup> examination and  
306 found an association between physical activity decline and BMI gain (46). Although assessing  
307 associations over three examinations are likely less influenced by confounding compared with  
308 two examinations, computing both exposure and outcome change from baseline still opens for  
309 reverse causation (*i.e.* weight gain potentially preceding physical activity decline). In the  
310 present study, we examined physical activity change from a 1<sup>st</sup> to a 2<sup>nd</sup> examination, followed  
311 by BMI change from the 2<sup>nd</sup> to a 3<sup>rd</sup> examination, which may be more suitable to assess the  
312 direction of the association, which provides an indication of causality (5).

313

314 Compared with the number of studies that examined whether physical activity is associated  
315 with BMI gain, fewer studies assessed a potential reverse association (*i.e.* BMI change predict  
316 physical activity change) (5). In those that did, high baseline BMI (6, 37) and BMI gain (22,  
317 25, 35, 37, 39) were associated with physical activity declines. In one study, baseline BMI,

318 but not BMI changes, was associated with physical activity declines (26). In a Mendelian  
319 randomization study, high body weight appeared causally associated with lower physical  
320 activity levels (38). Thus, except for one previous study (26), our study corroborates previous  
321 studies, suggesting that BMI gain leads to lower physical activity level.

322

323 Lower physical activity levels following weight gain are likely due to movement limitations.  
324 In a case-control study of normal weight and obese adolescents, physical activity measured by  
325 accelerometry was substantially lower in obese individuals compared with their normal  
326 weight peers despite similar physical activity energy expenditures (47). Similarly, this was  
327 also demonstrated in an experimental study of overfeeding with 4 MJ (1000 kilocalories  
328 (kcal)) per day over eight weeks, where free-living walking distances decreased due to lower  
329 walking velocity (*i.e.* movement limitation) in both normal weight and obese individuals  
330 following overfeeding, likely due to the increased weight (48).

331

332 Furthermore, our study contradicts a previous study, which reported that female but not male  
333 university alumni with high baseline BMI decreased their physical activity level over time (6),  
334 while we observed that both women and men were likely to decrease their physical activity  
335 with increasing BMI. This may be explained by demography (*e.g.* socioeconomic status, age)  
336 or by differences in analytical approach. Additionally, we observed that sex, birth year,  
337 baseline (T2) BMI, smoking, education and occupational physical activity change all  
338 modified the association between BMI change and subsequent physical activity change. This  
339 indicates that the effect of BMI change on physical activity change is dependent on multiple  
340 behavioural and societal factors, which warrants additional research.

341

342 A pertinent question may be whether population levels of physical activity are sufficiently  
343 high to prevent weight gain. One previous study estimated that a physical activity energy  
344 expenditure increase of ~0.4 megajoule (MJ) (*i.e.* 100 kcals) per day would be sufficient to  
345 prevent weight gain at population level (49), which could be feasible for the general  
346 population. However, highly active women who performed 60 minutes per day of moderate  
347 intensity activity (considerably higher physical activity energy expenditure than 0.4 MJ per  
348 day) seemed to still gain weight, but at a lower rate than their less active peers, indicating that  
349 such physical activity levels at best mitigates weight gain (28). Moreover, in another study,  
350 women and men being physically active at baseline had a lower baseline weight, but similar  
351 weight gain rate as those being inactive (26). Energy intake has increased with ~2 MJ (*i.e.* 500  
352 kcals) per day from the 1970s to 2000s in the United States, (50), which is similar to Western  
353 European countries from the 1960s to 2011 in a recent global study (51). About 110-150  
354 minutes of walking per day is needed to compensate for the increased energy intake of 2 MJ  
355 (50). This is seven times more than the current minimal recommendations for physical  
356 activity of 150 minutes per week (52). In Western high-income countries, one out of three fail  
357 to meet these recommendations (53). Consequently, the current physical activity levels in the  
358 general population is unlikely preventing population weight gain (5).

359

360 Our study with a comprehensive analytical approach showed no association between leisure  
361 time physical activity change and subsequent BMI change. However, we observed that BMI  
362 gains were associated with subsequent lower leisure time physical activity, which is consistent  
363 with previous studies (6, 22, 25, 26, 35, 37, 38). These observations are important, as public  
364 health initiatives aimed at weight gain prevention must acknowledge the major societal  
365 drivers for obesity in order to be successful (54, 55). As physical activity has numerous health  
366 effects independent of weight change (56), it should not be neglected, but simply



367 acknowledged in its limited potential for weight gain prevention (5). Although still ineffective  
368 (57), well-designed whole system approaches targeting multiple factors associated with  
369 population weight gain may be needed to shift the current curve of the obesity epidemic (55,  
370 57, 58).

371

## 372 **Strengths and limitations**

373 As BMI has gradually increased over decades (1), the long observation period in this study  
374 (~6 years between each examination) allowed us to examine whether physical activity change  
375 have affected the gradual long-term BMI gain (5). Further, as BMI change regressed on  
376 physical activity change, our models allowed us to interpret the direction of the association  
377 with more certainty (5). Furthermore, the merged overall cohort increased our sample size,  
378 which allowed us to assess effect modification in the association between physical activity  
379 and BMI. Finally, the Tromsø Study cohorts have high attendance of invited participants,  
380 which indicate high generalizability to high-income countries' populations (43).

381

382 There are also limitations that should be addressed. Self-reported physical activity change was  
383 categorized into crude groups; this may have introduced misclassification. Consequently,  
384 potential physical activity energy expenditure changes that could influence our results may  
385 have been missed. However, self-reported physical activity categorized into crude groups  
386 appears appropriate at population levels (59) and moreover, the SGPALS indicate predictive  
387 validity by being associated with multiple health outcomes (45). Moreover, our sensitivity  
388 analysis of six groups physical activity change showed similar results as our main analyses.  
389 Further, self-reported physical activity is likely influenced by recall and social desirability  
390 bias, which indicate that over-reporting of physical activity levels is inevitable (59). This is  
391 illustrated in our study by low variability in leisure time physical activity change, with most

392 of the included participants (68%) being classified as Persistently Active. These biases are  
393 likely to under- or overestimate the effect magnitude between physical activity and health  
394 outcomes (5) and might have influenced our results. Future long-term studies using physical  
395 activity instruments with higher accuracy (*e.g.*, device measured physical activity) are  
396 warranted to further examine whether population levels of physical activity influence weight  
397 change. Furthermore, disease onset may drive physical activity and weight change, which thus  
398 could be included as a potential confounder in our models. However, it is more likely that  
399 disease onset is a *mediator* (*i.e.* physical activity decline leads to disease, which leads to BMI  
400 change) or *ancestor* (*i.e.* disease onset leads to physical activity decline, which leads to BMI  
401 change) in the association between physical activity and BMI. Consequently, as our study's  
402 aims were to assess the total effect of physical activity change on BMI change and *vice versa*,  
403 adjusting for disease would not assess the total effect (60). Finally, our results may be  
404 influenced by residual confounding due to unavailable energy intake data.

405

## 406 **CONCLUSION**

407 In this prospective cohort study, there was no association between leisure time physical  
408 activity changes and subsequent BMI changes, whereas BMI increase was associated with  
409 subsequent consistently low and decreasing physical activity levels. These findings indicate  
410 that weight gain may lead to lower leisure time physical activity, while population levels of  
411 leisure time physical activity appears insufficient to prevent overweight and obesity.

412

413 **DECLARATIONS**

414 **Ethics approval and consent to participate**

415 All participants from Tromsø 4-7 provided written informed consent and the present study  
416 was approved by the Regional Ethics Committee for Medical Research (ref. 2016/758410).  
417

418 **Availability of data and materials**

419 The data that support the findings of this study are available from the Tromsø Study but  
420 restrictions apply to the availability of these data, which were used under license for the  
421 current study, and so are not publicly available. The data are however available from the  
422 Tromsø Study upon application to the Data and Publication Committee for the Tromsø  
423 Study: [tromsous@uit.no](mailto:tromsous@uit.no).  
424

425 **Competing interests**

426 The authors declare that they have no competing interests.  
427

428 **Funding**

429 The work of Edvard H Sagelv is funded by Population Studies in the High North  
430 (Befolkningsundersøkelser i Nord: BiN). The remaining authors are funded by their  
431 respective positions/tenures. The funders had no role in the implementation and design of the  
432 study or in writing the manuscript.  
433

434 **Authors' contributions**

435 EHS, BM, UE, LAH designed the study, EHS carried out data analysis, OL and TW provided  
436 statistical expertise, all authors interpreted the study results, EHS drafted the manuscript, and  
437 all authors contributed with manuscript revisions and approved the final version of the  
438 manuscript.

439

440 **Acknowledgements**

441 The authors would like to acknowledge Professor Bjarne Koster Jacobsen for valuable input  
442 on the study's result and for revising working manuscript drafts.

443

444 **REFERENCES**

- 445 1. NCD-RisC NRFC. Trends in adult body-mass index in 200 countries from 1975 to  
446 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million  
447 participants. *Lancet*. 2016;387(10026):1377-96.
- 448 2. Afshin A, Forouzanfar MH, Reitsma MB, Sur P, Estep K, Lee A, et al. Health Effects  
449 of Overweight and Obesity in 195 Countries over 25 Years. *N Engl J Med*. 2017;377(1):13-  
450 27.
- 451 3. Hill JO, Wyatt HR, Peters JC. Energy balance and obesity. *Circulation*.  
452 2012;126(1):126-32.
- 453 4. Jakicic JM, Powell KE, Campbell WW, Dipietro L, Pate RR, Pescatello LS, et al.  
454 Physical Activity and the Prevention of Weight Gain in Adults: A Systematic Review. *Med  
455 Sci Sports Exerc*. 2019;51(6):1262-9.
- 456 5. Jones PR, Ekelund U. Physical Activity in the Prevention of Weight Gain: the Impact  
457 of Measurement and Interpretation of Associations. *Curr Obes Rep*. 2019;8(2):66-76.

- 458 6. Basterra-Gortari FJ, Bes-Rastrollo M, Pardo-Fernández M, Forga L, Martinez JA,  
459 Martínez-González MA. Changes in weight and physical activity over two years in Spanish  
460 alumni. *Med Sci Sports Exerc.* 2009;41(3):516-22.
- 461 7. Blanck HM, McCullough ML, Patel AV, Gillespie C, Calle EE, Cokkinides VE, et al.  
462 Sedentary behavior, recreational physical activity, and 7-year weight gain among  
463 postmenopausal U.S. women. *Obesity (Silver Spring).* 2007;15(6):1578-88.
- 464 8. Brown WJ, Kabir E, Clark BK, Gomersall SR. Maintaining a Healthy BMI: Data  
465 From a 16-Year Study of Young Australian Women. *Am J Prev Med.* 2016;51(6):e165-e78.
- 466 9. Hillemeier MM, Weisman CS, Chuang C, Downs DS, McCall-Hosenfeld J, Camacho  
467 F. Transition to overweight or obesity among women of reproductive age. *J Womens Health.*  
468 2011;20(5):703-10.
- 469 10. Kaikkonen JE, Mikkilä V, Juonala M, Keltikangas-Järvinen L, Hintsanen M, Pulkki-  
470 Råback L, et al. Factors associated with six-year weight change in young and middle-aged  
471 adults in the Young Finns Study. *Scand J Clin Lab Invest.* 2015;75(2):133-44.
- 472 11. Rosenberg L, Kipping-Ruane KL, Boggs DA, Palmer JR. Physical activity and the  
473 incidence of obesity in young African-American women. *Am J Prev Med.* 2013;45(3):262-8.
- 474 12. Sjösten N, Kivimäki M, Singh-Manoux A, Ferrie JE, Goldberg M, Zins M, et al.  
475 Change in physical activity and weight in relation to retirement: the French GAZEL Cohort  
476 Study. *BMJ Open.* 2012;2(1):e000522.
- 477 13. Williams PT, Wood PD. The effects of changing exercise levels on weight and age-  
478 related weight gain. *Int J Obes.* 2006;30(3):543-51.
- 479 14. Williams PT, Thompson PD. Dose-dependent effects of training and detraining on  
480 weight in 6406 runners during 7.4 years. *Obesity.* 2006;14(11):1975-84.

- 481 15. Gebel K, Ding D, Bauman AE. Volume and intensity of physical activity in a large  
482 population-based cohort of middle-aged and older Australians: prospective relationships with  
483 weight gain, and physical function. *Prev Med.* 2014;60:131-3.
- 484 16. Gradidge PJ, Norris SA, Micklesfield LK, Crowther NJ. The Role of Lifestyle and  
485 Psycho-Social Factors in Predicting Changes in Body Composition in Black South African  
486 Women. *PLoS One.* 2015;10(7):e0132914.
- 487 17. MacInnis RJ, Hodge AM, Dixon HG, Peeters A, Johnson LE, English DR, et al.  
488 Predictors of increased body weight and waist circumference for middle-aged adults. *Public  
489 Health Nutr.* 2014;17(5):1087-97.
- 490 18. Sims ST, Larson JC, Lamonte MJ, Michael YL, Martin LW, Johnson KC, et al.  
491 Physical activity and body mass: changes in younger versus older postmenopausal women.  
492 *Med Sci Sports Exerc.* 2012;44(1):89-97.
- 493 19. Chiriboga DE, Ma Y, Li W, Olendzki BC, Pagoto SL, Merriam PA, et al. Gender  
494 differences in predictors of body weight and body weight change in healthy adults. *Obesity.*  
495 2008;16(1):137-45.
- 496 20. Barone Gibbs B, Pettee Gabriel K, Carnethon MR, Gary-Webb T, Jakicic JM, Rana  
497 JS, et al. Sedentary Time, Physical Activity, and Adiposity: Cross-sectional and Longitudinal  
498 Associations in CARDIA. *Am J Prev Med.* 2017;53(6):764-71.
- 499 21. Dugas LR, Kliethermes S, Plange-Rhule J, Tong L, Bovet P, Forrester TE, et al.  
500 Accelerometer-measured physical activity is not associated with two-year weight change in  
501 African-origin adults from five diverse populations. *PeerJ.* 2017;5:e2902.
- 502 22. Ekelund U, Kolle E, Steene-Johannessen J, Dalene KE, Nilsen AKO, Anderssen SA,  
503 et al. Objectively measured sedentary time and physical activity and associations with body  
504 weight gain: does body weight determine a decline in moderate and vigorous intensity  
505 physical activity? *Int J Obes* 2017;41(12):1769-74.

- 506 23. Botoseneanu A, Liang J. The effect of stability and change in health behaviors on  
507 trajectories of body mass index in older Americans: a 14-year longitudinal study. *J Gerontol*  
508 *A Biol Sci Med Sci.* 2012;67(10):1075-84.
- 509 24. de Munter JS, Tynelius P, Magnusson C, Rasmussen F. Longitudinal analysis of  
510 lifestyle habits in relation to body mass index, onset of overweight and obesity: results from a  
511 large population-based cohort in Sweden. *Scand J Public Health.* 2015;43(3):236-45.
- 512 25. Drenowatz C, Gribben N, Wirth MD, Hand GA, Shook RP, Burgess S, et al. The  
513 Association of Physical Activity during Weekdays and Weekend with Body Composition in  
514 Young Adults. *J Obes.* 2016;2016:8236439.
- 515 26. Hamer M, Brunner EJ, Bell J, Batty GD, Shipley M, Akbaraly T, et al. Physical  
516 activity patterns over 10 years in relation to body mass index and waist circumference: the  
517 Whitehall II cohort study. *Obesity.* 2013;21(12):E755-61.
- 518 27. Hankinson AL, Daviglus ML, Bouchard C, Carnethon M, Lewis CE, Schreiner PJ, et  
519 al. Maintaining a high physical activity level over 20 years and weight gain. *JAMA.*  
520 2010;304(23):2603-10.
- 521 28. Lee IM, Djoussé L, Sesso HD, Wang L, Buring JE. Physical activity and weight gain  
522 prevention. *JAMA.* 2010;303(12):1173-9.
- 523 29. Moholdt T, Wisløff U, Lydersen S, Nauman J. Current physical activity guidelines for  
524 health are insufficient to mitigate long-term weight gain: more data in the fitness versus  
525 fatness debate (The HUNT study, Norway). *Br J Sports Med.* 2014;48(20):1489-96.
- 526 30. Parsons TJ, Manor O, Power C. Physical activity and change in body mass index from  
527 adolescence to mid-adulthood in the 1958 British cohort. *Int J Epidemiol.* 2006;35(1):197-  
528 204.

- 529 31. Smith KJ, Gall SL, McNaughton SA, Cleland VJ, Otahal P, Dwyer T, et al. Lifestyle  
530 behaviours associated with 5-year weight gain in a prospective cohort of Australian adults  
531 aged 26-36 years at baseline. *BMC Public Health*. 2017;17(1):54.
- 532 32. Kim Y, Lee JM, Kim J, Dhurandhar E, Soliman G, Wehbi NK, et al. Longitudinal  
533 associations between body mass index, physical activity, and healthy dietary behaviors in  
534 adults: A parallel latent growth curve modeling approach. *PLoS One*. 2017;12(3):e0173986.
- 535 33. Su C, Jia XF, Wang ZH, Wang HJ, Ouyang YF, Zhang B. Longitudinal association of  
536 leisure time physical activity and sedentary behaviors with body weight among Chinese adults  
537 from China Health and Nutrition Survey 2004-2011. *Eur J Clin Nutr*. 2017;71(3):383-8.
- 538 34. Brien SE, Katzmarzyk PT, Craig CL, Gauvin L. Physical activity, cardiorespiratory  
539 fitness and body mass index as predictors of substantial weight gain and obesity: the Canadian  
540 physical activity longitudinal study. *Can J Public Health*. 2007;98(2):121-4.
- 541 35. Drenowatz C, Hill JO, Peters JC, Soriano-Maldonado A, Blair SN. The association of  
542 change in physical activity and body weight in the regulation of total energy expenditure. *Eur*  
543 *J Clin Nutr*. 2017;71(3):377-82.
- 544 36. French SA, Mitchell NR, Hannan PJ. Decrease in television viewing predicts lower  
545 body mass index at 1-year follow-up in adolescents, but not adults. *J Nutr Educ Behav*.  
546 2012;44(5):415-22.
- 547 37. Mortensen LH, Siegler IC, Barefoot JC, Grønbaek M, Sørensen TI. Prospective  
548 associations between sedentary lifestyle and BMI in midlife. *Obesity*. 2006;14(8):1462-71.
- 549 38. Richmond RC, Davey Smith G, Ness AR, den Hoed M, McMahon G, Timpson NJ.  
550 Assessing causality in the association between child adiposity and physical activity levels: a  
551 Mendelian randomization analysis. *PLoS Med*. 2014;11(3):e1001618.



- 552 39. Golubic R, Ekelund U, Wijndaele K, Luben R, Khaw KT, Wareham NJ, et al. Rate of  
553 weight gain predicts change in physical activity levels: a longitudinal analysis of the EPIC-  
554 Norfolk cohort. *Int J Obes*. 2013;37(3):404-9.
- 555 40. Sagelv EH, Ekelund U, Hopstock LA, Aars NA, Fimland MS, Jacobsen BK, et al. Do  
556 declines in occupational physical activity contribute to population gains in body mass index?  
557 Tromsø Study 1974–2016. *Occup Environ Med*. 2021;78(3):203-10.
- 558 41. Morseth B, Hopstock LA. Time trends in physical activity in the Tromsø study: An  
559 update. *PLoS ONE*. 2020;15(4):e0231581.
- 560 42. Church TS, Thomas DM, Tudor-Locke C, Katzmarzyk PT, Earnest CP, Rodarte RQ,  
561 et al. Trends over 5 decades in U.S. occupation-related physical activity and their associations  
562 with obesity. *PLoS One*. 2011;6(5):e19657.
- 563 43. Jacobsen BK, Eggen AE, Mathiesen EB, Wilsgaard T, Njølstad I. Cohort profile: the  
564 Tromsø Study. *Int J Epidemiol*. 2012;41(4):961-7.
- 565 44. Saltin B, Grimby G. Physiological analysis of middle-aged and old former athletes.  
566 Comparison with still active athletes of the same ages. *Circulation*. 1968;38(6):1104-15.
- 567 45. Grimby G, Börjesson M, Jonsdottir IH, Schnohr P, Thelle DS, Saltin B. The "Saltin-  
568 Grimby Physical Activity Level Scale" and its application to health research. *Scand J Med Sci*  
569 *Sports*. 2015;25 Suppl 4:119-25.
- 570 46. Shibata AI, Oka K, Sugiyama T, Salmon JO, Dunstan DW, Owen N. Physical  
571 Activity, Television Viewing Time, and 12-Year Changes in Waist Circumference. *Med Sci*  
572 *Sports Exerc*. 2016;48(4):633-40.
- 573 47. Ekelund U, Åman J, Yngve A, Renman C, Westerterp K, Sjöström M. Physical  
574 activity but not energy expenditure is reduced in obese adolescents: a case-control study. *Am*  
575 *J Clin Nutr*. 2002;76(5):935-41.

- 576 48. Levine JA, McCrady SK, Lanningham-Foster LM, Kane PH, Foster RC, Manohar CU.  
577 The role of free-living daily walking in human weight gain and obesity. *Diabetes*.  
578 2008;57(3):548-54.
- 579 49. Hill JO, Wyatt HR, Reed GW, Peters JC. Obesity and the environment: where do we  
580 go from here? *Science*. 2003;299(5608):853-5.
- 581 50. Swinburn B, Sacks G, Ravussin E. Increased food energy supply is more than  
582 sufficient to explain the US epidemic of obesity. *Am J Clin Nutr*. 2009;90(6):1453-6.
- 583 51. Dave D, Doytch N, Kelly IR. Nutrient intake: A cross-national analysis of trends and  
584 economic correlates. *Soc Sci Med*. 2016;158:158-67.
- 585 52. Bull FC, Al-Ansari SS, Biddle S, Borodulin K, Buman MP, Cardon G, et al. World  
586 Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sport*  
587 *Med*. 2020;54(24):1451-62.
- 588 53. Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends in insufficient physical  
589 activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1.9  
590 million participants. *Lancet Glob Health*. 2018;6(10):e1077-e86.
- 591 54. Swinburn BA, Sacks G, Hall KD, McPherson K, Finegood DT, Moodie ML, et al. The  
592 global obesity pandemic: shaped by global drivers and local environments. *Lancet*.  
593 2011;378(9793):804-14.
- 594 55. Vandevijvere S, Chow CC, Hall KD, Umali E, Swinburn BA. Increased food energy  
595 supply as a major driver of the obesity epidemic: a global analysis. *Bull World Health Organ*.  
596 2015;93(7):446-56.
- 597 56. Warburton DER, Bredin SSD. Health benefits of physical activity: a systematic review  
598 of current systematic reviews. *Curr Opin Cardiol*. 2017;32(5):541-56.

- 599 57. Bagnall AM, Radley D, Jones R, Gately P, Nobles J, Van Dijk M, et al. Whole  
600 systems approaches to obesity and other complex public health challenges: a systematic  
601 review. BMC Public Health. 2019;19(1):8.
- 602 58. Lee BY, Bartsch SM, Mui Y, Haidari LA, Spiker ML, Gittelsohn J. A systems  
603 approach to obesity. Nutr Rev. 2017;75(suppl 1):94-106.
- 604 59. Shephard R, Vuillemin A. Limits to the measurement of habitual physical activity by  
605 questionnaires. Br J Sports Med. 2003;37.
- 606 60. Aalen OO, Røysland K, Gran JM, Ledergerber B. Causality, mediation and time: a  
607 dynamic viewpoint. J R Stat Soc. 2012;175(4):831-61.

608

## 609 **FIGURE LEGENDS**

610 **Figure 1:** The study design for assessing the association between physical activity changes  
611 and future BMI change, and conversely for assessing BMI changes and physical activity  
612 changes. BMI=body mass index.

613 **Figure 2:** Flow chart of participant selection.

614 **Figure 3:** Flow chart of participant selection for the reversed analyses.

615

## 616 **TABLE LEGENDS**

617 **Table 1.** Descriptive characteristics of the overall cohort and period-specific samples. The  
618 Tromsø Study 1974-2016.

619 **Table 2.** Change in BMI from T2 to T3 by leisure physical activity change from T1 to T2.  
620 The Tromsø Study 1974-2016.

621 **Table 3.** Odds Ratio of leisure time physical activity change with body mass index change  
622 (per kg/m<sup>2</sup> increase). The Tromsø Study 1974-2016.

---

## Main analyses

---

**Time point 1**

**Time point 2**

**Time point 3**

Change Physical Activity

Adjustments:

Sex

Education

Smoking

Birth year

Change BMI

«Baseline» BMI or Weight

Tromsø 5 (2001-02)

Tromsø 2 (1979-80)

Tromsø 1 (1974)

Tromsø 6 (2007-08)

Tromsø 3 (1986-87)

Tromsø 2 (1979-80)

Tromsø 7 (2015-16)

Tromsø 4 (1994-95)

Tromsø 3 (1986-87)

---

## Reverse analyses

---

Change BMI

Adjustments:

Sex

Education

Smoking

Birth year

Change Physical Activity

Tromsø 5 (2001-02)

Tromsø 4 (1994-95)

Tromsø 1 (1974)

Tromsø 6 (2007-08)


Tromsø 5 (2001-02)


Tromsø 2 (1979-80)

Tromsø 7 (2015-16)

Tromsø 6 (2007-08)

Tromsø 3 (1986-87)





**Table 1.** Descriptive characteristics of the overall cohort and period-specific samples. The Tromsø Study 1974-2016.

Cohort	Overall cohort*		Period-specific samples**		
		Tromsø 1-7 (1974-2016)	Tromsø 1-3 (1974-1986)	Tromsø 2-4 (1979-1995)	Tromsø 5-7 (2001-2016)
<b>Total N (%)</b>		10779 (100%)	3598 (100%)	9691 (100%)	2206 (100%)
<b>BMI (kg/m<sup>2</sup>)</b>					
Time point 2	Mean	24.81	24.65	24.25	26.93
	95%CI	24.74 to 24.88	24.56 to 24.74	24.18 to 24.32	26.75 to 27.11
Time point 3	Mean	25.67	25.15	25.38	26.93
	95%CI	25.60 to 25.74	25.05 to 25.25	25.31 to 25.45	26.75 to 27.11
Change time point 2-3	Mean	0.86	0.49	1.13	-0.002
	95%CI	0.82 to 0.90	0.44 to 0.53	1.10 to 1.17	-0.09 to 0.08
<b>Baseline</b>		<b>Time point 2</b>	<b>Tromsø 2 (1979-80)</b>	<b>Tromsø 3 (1986-87)</b>	<b>Tromsø 6 (2007-08)</b>
<b>Sex</b>					
Women	n (%)	5195 (48.2%)	N/A	4834 (49.9%)	1273 (57.7%)
Men	n (%)	5584 (51.8%)	3598 (100%)	4857 (50.1%)	933 (42.3%)
<b>Age (yr)</b>	Mean	46.19	39.78	42.59	62.04
	95%CI	45.96 to 46.42	39.51 to 40.05	42.42 to 42.76	61.65 to 62.43
<b>10-year age groups</b>					
≤39 years	n (%)	3837 (35.6%)	1824 (50.7%)	3836 (39.9%)	36 (1.6%)
40-49 years	n (%)	2917 (27.1%)	1199 (33.3%)	3512 (36.2%)	289 (13.1%)
50-59 years	n (%)	2238 (20.8%)	575 (16.0%)	2110 (21.8%)	327 (14.8%)
60-69 years	n (%)	1326 (12.3%)	N/A	233 (2.4%)	1093 (49.5%)
≥70 years	n (%)	461 (4.3%)	N/A	N/A	461 (20.9%)
<b>BMI groups</b>					
Normal weight	n (%)	6276 (58.2%)	2138 (59.4%)	6255 (64.5%)	759 (34.4%)
Overweight	n (%)	3594 (33.3%)	1313 (36.5%)	2920 (30.1%)	1011 (45.8%)
Obese	n (%)	909 (8.4%)	147 (4.1%)	516 (5.3%)	436 (19.8%)
<b>Smoking</b>					
Current smoker	n (%)	4316 (40.0%)	1720 (47.8%)	4226 (43.6%)	360 (16.3%)
Previous smoker	n (%)	1715 (15.9%)	505 (14.0%)	754 (7.8%)	1019 (46.2%)
Never smoker	n (%)	4748 (44.1%)	1373 (38.2%)	4711 (48.6%)	828 (37.5%)
<b>Education</b>					
Primary school	n (%)	4555 (42.3%)	1860 (51.7%)	4331 (44.7%)	719 (32.6%)
High school	n (%)	3368 (31.2%)	1009 (28.0%)	2938 (30.3%)	772 (35.0%)
University <4 years	n (%)	1576 (14.6%)	426 (11.8%)	1381 (14.3%)	364 (16.5%)
University ≥4 years	n (%)	1280 (11.9%)	303 (8.4%)	1041 (10.7%)	351 (15.9%)
<b>Reverse analyses</b>					
<b>Total</b>	N (%)	4385 (100%)	N/A	N/A	N/A
<b>BMI (kg/m<sup>2</sup>)</b>	Mean	25.64	N/A	N/A	N/A
	95%CI	25.53 to 25.75			
<b>Sex</b>					
Women	n (%)	1307 (29.8%)	N/A	N/A	N/A
Men	n (%)	3078 (70.2%)	N/A	N/A	N/A
<b>Age (yr)</b>	Mean	50.63	N/A	N/A	N/A
	95%CI	50.16 to 51.10			
<b>10-year age groups</b>					
≤39 years	n (%)	1489 (34%)	N/A	N/A	N/A
40-49 years	n (%)	647 (14.8%)	N/A	N/A	N/A
50-59 years	n (%)	601 (13.7%)	N/A	N/A	N/A
60-69 years	n (%)	1063 (24.2%)	N/A	N/A	N/A
≥70 years	n (%)	585 (13.3%)	N/A	N/A	N/A
<b>BMI groups</b>					
Normal weight	n (%)	2131 (48.6%)	N/A	N/A	N/A
Overweight	n (%)	1746 (39.8%)	N/A	N/A	N/A
Obese	n (%)	508 (11.6%)	N/A	N/A	N/A
<b>Smoking</b>					
Current smoker	n (%)	1396 (31.8%)	N/A	N/A	N/A
Previous smoker	n (%)	1372 (31.3%)	N/A	N/A	N/A
Never smoker	n (%)	1617 (36.9%)	N/A	N/A	N/A
<b>Education</b>					
Primary school	n (%)	1731 (39.5%)	N/A	N/A	N/A
High school	n (%)	1432 (32.7%)	N/A	N/A	N/A
University <4 years	n (%)	672 (15.3%)	N/A	N/A	N/A
University ≥4 years	n (%)	550 (12.5%)	N/A	N/A	N/A

Data are shown as unadjusted mean and 95%CI or as frequency and percentage. \*The overall cohort includes participants attending ≥3 surveys and the analyses are based on their three most recent surveys. \*\*Period specific samples include all participants meeting our inclusion criteria for that period, and each participant may contribute in more than one period; therefore, these samples do not add up to the overall cohort (Tromsø 1-7). CI=confidence interval.

**Table 2.** Change in BMI from T2 to T3 by leisure physical activity change from T1 to T2. The Tromsø Study 1974-2016.

Tromsø 1-7 (1974-2016)	Change leisure physical activity T1 to T2					P <sub>equality</sub>
	Total	Persistently Inactive	Persistently Active	Active to Inactive	Inactive to Active	
<i>Total (n)</i>	10779	992	7314	1167	1306	
BMI T2 (kg/m <sup>2</sup> )*	Mean	25.25	24.73	24.64	25.05	
	95% CI	24.99 to 25.51	24.65 to 24.81	24.43 to 24.85	24.83 to 25.27	
	<b>BMI change T2 to T3</b>					
BMI change (kg/m <sup>2</sup> )	Mean	0.89	0.85	0.90	0.85	0.84
	95% CI	0.77 to 1.00	0.81 to 0.89	0.79 to 1.00	0.75 to 0.95	
<b>Sex</b>						
<i>Women (n)</i>	5195	490	3481	594	630	
BMI change (kg/m <sup>2</sup> )	Mean	1.23	1.08	1.13	1.08	0.48
	95% CI	1.05 to 1.41	1.01 to 1.15	0.96 to 1.29	0.92 to 1.24	
<i>Men (n)</i>	5584	502	3833	573	676	
BMI change (kg/m <sup>2</sup> )	Mean	0.56	0.64	0.68	0.64	0.67
	95% CI	0.42 to 0.70	0.59 to 0.69	0.55 to 0.82	0.51 to 0.76	
<b>Birth year</b>						
<i>≤1929 (n)</i>	687	56	456	84	91	
BMI change (kg/m <sup>2</sup> )	Mean	0.01	0.17	0.05	0.07	0.82
	95% CI	-0.42 to 0.43	0.02 to 0.32	-0.30 to 0.40	-0.26 to 0.41	
<i>1930-1939 (n)</i>	2868	234	2017	274	343	
BMI change (kg/m <sup>2</sup> )	Mean	0.42	0.47	0.45	0.53	0.92
	95% CI	0.18 to 0.66	0.39 to 0.55	0.22 to 0.67	0.33 to 0.73	
<i>1940-1949 (n)</i>	4115	409	2804	412	490	
BMI change (kg/m <sup>2</sup> )	Mean	1.01	0.93	0.95	0.94	0.89
	95% CI	0.82 to 1.19	0.86 to 1.00	0.77 to 1.14	0.77 to 1.10	
<i>1950-1959 (n)</i>	2821	269	1825	364	363	
BMI change (kg/m <sup>2</sup> )	Mean	1.29	1.33	1.40	1.25	0.72
	95% CI	1.08 to 1.50	1.25 to 1.41	1.22 to 1.58	1.07 to 1.43	
<i>≥1960 (n)</i>	288	24	212	33	19	
BMI change (kg/m <sup>2</sup> )	Mean	1.47	0.60	1.44	0.06	0.09
	95% CI	0.47 to 2.47	0.27 to 0.94	0.59 to 2.30	-1.07 to 1.19	
<b>BMI groups</b>						
<i>Normal weight (n)</i>	6276	524	4311	704	737	
BMI change (kg/m <sup>2</sup> )	Mean	1.04	1.01	1.02	1.04	0.96
	95% CI	0.90 to 1.18	0.96 to 1.06	0.90 to 1.14	0.93 to 1.16	
<i>Overweight (n)</i>	3594	352	2440	376	426	
BMI change (kg/m <sup>2</sup> )	Mean	0.86	0.81	0.92	0.83	0.77
	95% CI	0.66 to 1.07	0.73 to 0.89	0.72 to 1.16	0.65 to 1.02	
<i>Obese (n)</i>	909	116	563	87	143	
BMI change (kg/m <sup>2</sup> )	Mean	0.06	0.05	0.06	-0.15	0.89
	95% CI	-0.45 to 0.56	-0.19 to 0.28	-0.53 to 0.65	-0.61 to 0.32	
<b>Smoking</b>						
<i>Current smoker (n)</i>	4316	521	2570	541	684	
BMI change (kg/m <sup>2</sup> )	Mean	0.92	1.07	0.99	0.93	0.16
	95% CI	0.75 to 1.08	1.00 to 1.15	0.83 to 1.15	0.79 to 1.07	
<i>Previous smoker (n)</i>	1715	135	1224	190	166	
BMI change (kg/m <sup>2</sup> )	Mean	0.34	0.29	0.46	0.30	0.71
	95% CI	0.02 to 0.66	0.19 to 0.40	0.19 to 0.73	0.01 to 0.59	
<i>Never smoker (n)</i>	4748	336	3520	436	356	
BMI change (kg/m <sup>2</sup> )	Mean	1.07	0.88	0.98	0.96	0.15
	95% CI	0.89 to 1.26	0.82 to 0.93	0.81 to 1.15	0.80 to 1.13	
<b>Education</b>						
<i>Primary school (n)</i>	4555	465	2921	534	635	
BMI change (kg/m <sup>2</sup> )	Mean	0.82	0.83	0.73	0.69	0.32
	95% CI	0.65 to 1.00	0.76 to 0.90	0.56 to 0.89	0.54 to 0.84	
<i>High school (n)</i>	3368	317	2300	368	383	



BMI change (kg/m <sup>2</sup> )	Mean	0.93	0.91	0.87	0.96	0.92
	95% CI	0.72 to 1.13	0.83 to 0.98	0.68 to 1.06	0.78 to 1.15	
<i>University &lt;4 years (n)</i>		1576	1173	135	162	
BMI change (kg/m <sup>2</sup> )	Mean	0.90	0.87	1.53	0.95	0.001
	95% CI	0.55 to 1.25	0.76 to 0.97	1.22 to 1.84	0.66 to 1.23	
<i>University &gt;4 years (n)</i>		1280	104	920	130	126
BMI change (kg/m <sup>2</sup> )	Mean	0.99	0.80	0.95	1.16	0.08
	95% CI	0.69 to 1.30	0.69 to 0.90	0.67 to 1.22	0.88 to 1.43	

---

#### Occupational Physical Activity Change T1 to T2

---

<i>Persistently Inactive</i>		2637	340	1650	303	344	
BMI change (kg/m <sup>2</sup> )	Mean	0.81	0.85	0.93	0.92	0.79	
	95% CI	0.61 to 1.00	0.76 to 0.94	0.72 to 1.14	0.72 to 1.11		
<i>Persistently Active</i>		5014	372	3514	539	589	
BMI change (kg/m <sup>2</sup> )	Mean	1.04	1.07	1.05	1.01	0.92	
	95% CI	0.86 to 1.23	1.01 to 1.13	0.89 to 1.20	0.86 to 1.16		
<i>Active to Inactive</i>		673	62	439	96	76	
BMI change (kg/m <sup>2</sup> )	Mean	1.43	0.89	0.88	0.95	0.13	
	95% CI	1.01 to 1.86	0.73 to 1.04	0.54 to 1.22	0.57 to 1.33		
<i>Inactive to Active</i>		1277	1144	799	129	205	
BMI change (kg/m <sup>2</sup> )	Mean	1.03	0.99	0.96	0.90	0.91	
	95% CI	0.73 to 1.33	0.86 to 1.11	0.64 to 1.28	0.65 to 1.15		

---

Data are adjusted for sex, birth year, smoking, education and BMI at T2, and shown as adjusted mean BMI change with 95% CI.\*Data are shown as unadjusted mean BMI at T2 with 95% CI. CI=confidence interval, BMI=body mass index, P<sub>equality</sub>=main difference between groups, T1=time point 1, T2=time point 2, T3=time point 3.

**Table 3.** Odds Ratio of leisure time physical activity change with body mass index change (per kg/m<sup>2</sup> increase).  
The Tromsø Study 1974-2016.

<b>Tromsø 1-7 (1974-2016)</b>		<b>Persistently Inactive</b>	<b>Active to Inactive</b>	<b>Inactive to Active</b>	<b>Persistently Active</b>
Total (N)	4385	378	397	512	3098
	OR	1.14	1.16	1.02	Ref.
	95%CI	1.07 to 1.22	1.09 to 1.24	0.96 to 1.09	
<b>Sex</b>					
<i>P</i> <sub>interaction</sub>		<0.001	0.003	0.59	
Women (n)	1307	107	102	118	980
	OR	1.09	1.21	1.05	Ref.
	95%CI	0.98 to 1.22	1.09 to 1.34	0.95 to 1.16	
Men (n)	3078	271	295	394	2118
	OR	1.19	1.14	1.02	Ref.
	95%CI	1.09 to 1.30	1.04 to 1.23	0.95 to 1.10	
<b>Birth year</b>					
<i>P</i> <sub>interaction</sub>		<0.001	<0.001	0.17	
≤1939 (n)	1473	135	132	203	1003
	OR	1.11	1.01	0.94	Ref.
	95%CI	0.99 to 1.24	0.90 to 1.14	0.85 to 1.04	
1940-49 (n)	1906	162	171	205	1368
	OR	1.13	1.21	1.06	Ref.
	95%CI	1.02 to 1.25	1.10 to 1.34	0.96 to 1.16	
≥1950 (n)	1006	81	94	104	727
	OR	1.23	1.28	1.10	Ref.
	95%CI	1.06 to 1.43	1.12 to 1.45	0.96 to 1.26	
<b>BMI groups</b>					
<i>P</i> <sub>interaction</sub>		<0.001	<0.001	0.29	
Normal weight (n)	2131	153	176	227	1575
	OR	0.94	1.07	0.96	Ref.
	95%CI	0.83 to 1.06	0.95 to 1.20	0.86 to 1.07	
Overweight (n)	1746	149	163	208	1226
	OR	1.15	1.12	1.01	Ref.
	95%CI	1.04 to 1.28	1.01 to 1.24	0.92 to 1.11	
Obese (n)	508	76	58	77	297
	OR	1.17	1.21	1.00	Ref.
	95%CI	1.03 to 1.31	1.07 to 1.38	0.88 to 1.13	
<b>Smoking</b>					
<i>P</i> <sub>interaction</sub>		<0.001	<0.001	0.61	
Current smoker (n)	1396	180	168	218	830
	OR	1.14	1.14	1.00	Ref.
	95%CI	1.02 to 1.28	1.02 to 1.28	0.90 to 1.11	
Previous smoker (n)	1372	93	102	152	1025
	OR	1.18	1.23	1.0	Ref.
	95%CI	1.06 to 1.32	1.11 to 1.37	0.95 to 1.15	
Never smoker (n)	1617	105	127	142	1243
	OR	1.13	1.14	1.02	Ref.
	95%CI	0.99 to 1.30	1.00 to 1.29	0.90 to 1.15	
<b>Education</b>					
<i>P</i> <sub>interaction</sub>		0.008	<0.001	0.47	
Primary school (n)	1731	188	171	124	1129
	OR	1.17	1.11	1.04	Ref.
	95%CI	1.07 to 1.28	1.00 to 1.22	0.95 to 1.13	
High school (n)	1432	113	117	164	1038
	OR	1.15	1.12	0.96	Ref.
	95%CI	1.01 to 1.30	0.99 to 1.26	0.86 to 1.06	

University <4 years (n)	672	44	60	65	503
	OR	1.13	1.20	1.17	Ref.
	95%CI	0.89 to 1.43	0.98 to 1.46	0.96 to 1.42	
University ≥4 years (n)	550	33	49	40	428
	OR	0.95	1.38	1.04	Ref.
	95%CI	0.76 to 1.18	1.17 to 1.64	0.85 to 1.28	
<b>Occupational Physical Activity Change from T1 to T2</b>					
<i>P<sub>interaction</sub></i>		0.01	<0.001	0.81	
Persistently Inactive (n)	1125	129	119	124	753
	OR	1.24	1.12	0.96	Ref.
	95%CI	1.09 to 1.42	0.98 to 1.28	0.85 to 1.08	
Persistently Active (n)	1536	106	142	178	1110
	OR	1.17	1.17	1.06	Ref.
	95%CI	1.04 to 1.32	1.05 to 1.30	0.96 to 1.18	
Active to Inactive (n)	248	19	37	36	156
	OR	0.96	1.23	0.88	Ref.
	95%CI	0.66 to 1.41	0.93 to 1.63	0.66 to 1.16	
Inactive to Active (n)	341	33	25	53	230
	OR	1.07	1.33	0.94	Ref.
	95%CI	0.83 to 1.38	1.01 to 1.75	0.76 to 1.16	

Data are adjusted for sex, birth year, smoking and education at T2, and shown as adjusted OR with 95% CI.  $P_{interaction}$ =interaction effect for the stratified variable and BMI change on physical activity change, BMI=body mass index, CI=confidence interval, OR=Odds Ratio. T2=time point 2.